



NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Rectal Cancer

Version 3.2017 — March 13, 2017

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 3.2017 Panel Members

Rectal Cancer

[NCCN Guidelines Index](#)
[Table of Contents](#)
[Discussion](#)

* **Al B. Benson, III, MD/Chair †**
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

* **Alan P. Venook, MD/Vice-Chair † ‡**
UCSF Helen Diller Family
Comprehensive Cancer Center

Lynette Cederquist, MD †
UC San Diego Moores Cancer Center

Emily Chan, MD, PhD †
Vanderbilt-Ingram Cancer Center

Yi-Jen Chen, MD, PhD §
City of Hope Comprehensive
Cancer Center

Harry S. Cooper, MD ≠
Fox Chase Cancer Center

Dustin Deming, MD †
University of Wisconsin
Carbone Cancer Center

Paul F. Engstrom, MD †
Fox Chase Cancer Center

Peter C. Enzinger, MD †
Dana-Farber/Brigham and Women's
Cancer Center

Alessandro Fichera, MD
Fred Hutchinson Cancer Research
Center/Seattle Cancer Care Alliance

Jean L. Grem, MD †
Fred & Pamela Buffett Cancer Center

Axel Grothey, MD †
Mayo Clinic Cancer Center

Howard S. Hochster, MD †
Yale Cancer Center/Smilow Cancer
Hospital

Sarah Hoffe, MD §
Moffitt Cancer Center

Steven Hunt, MD ¶
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Ahmed Kamel, MD ¶
University of Alabama at Birmingham
Comprehensive Cancer Center

Natalie Kirilcuk, MD ¶
Stanford Cancer Institute

Smitha Krishnamurthi, MD † †
Case Comprehensive Cancer Center/
University Hospitals Seidman
Cancer Center and Cleveland
Clinic Taussig Cancer Institute

* **Wells A. Messersmith, MD †**
University of Colorado Cancer Center

Mary F. Mulcahy, MD ‡ †
Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

James D. Murphy, MD, MS §
UC San Diego Moores Cancer Center

Steven Nurkin, MD, MS ¶
Roswell Park Cancer Institute

Leonard Saltz, MD † ‡ †
Memorial Sloan Kettering Cancer Center

Sunil Sharma, MD †
Huntsman Cancer Institute
at the University of Utah

David Shibata, MD ¶
The University of Tennessee
Health Science Center

John M. Skibber, MD ¶
The University of Texas
MD Anderson Cancer Center

Constantinos T. Sofocleous, MD, PhD ¶
Memorial Sloan Kettering Cancer Center

Elena M. Stoffel, MD, MPH ≠
University of Michigan
Comprehensive Cancer Center

Eden Stotsky-Himelfarb, BSN, RN ≠
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Christopher G. Willett, MD §
Duke Cancer Institute

Christina S. Wu, MD
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

NCCN
Deborah Freedman-Cass, PhD
Kristina M. Gregory, RN, MSN, OCN

Continue

† Medical oncology	‡ Internal medicine
§ Radiotherapy/Radiation oncology	¶ Diagnostic/Interventional radiology
¶ Surgery/Surgical oncology	≠ Gastroenterology
≠ Pathology	¥ Patient advocate
‡ Hematology/Hematology oncology	*Discussion Section Writing Committee

[NCCN Guidelines Panel Disclosures](#)



[NCCN Rectal Cancer Panel Members](#)

[Summary of the Guidelines Updates](#)

Clinical Presentations and Primary Treatment:

- [Pedunculated polyp \(adenoma\) with Invasive Cancer \(REC-1\)](#)
- [Sessile Polyp \(adenoma\) with Invasive Cancer \(REC-1\)](#)
- [Rectal Cancer Appropriate for Resection \(REC-2\)](#)
 - ▶ [Treatment after Transanal Excision of cT1-2, N0 \(REC-3\)](#)
 - ▶ [Treatment after Transabdominal Resection of cT1-2, N0 \(REC-4\)](#)
 - ▶ [T3, N0 or T any, N1-2 or T4 and/or Locally Unresectable or Medically Inoperable Primary and Adjuvant Treatment \(REC-5\)](#)
 - ▶ [T any, N any, M1: Resectable Synchronous Metastases \(REC-6\)](#)
- [T any, N any, M1: Unresectable Synchronous Metastases or Medically Inoperable Treatment \(REC-7\)](#)

[Surveillance \(REC-8\)](#)

[Recurrence and Workup \(REC-9\)](#)

[Metachronous Metastases \(REC-9\)](#)

[Principles of Pathologic Review \(REC-A\)](#)

[Principles of Surgery \(REC-B\)](#)

[Principles of Adjuvant Therapy \(REC-C\)](#)

[Principles of Radiation Therapy \(REC-D\)](#)

[Chemotherapy for Advanced or Metastatic Disease \(REC-E\)](#)

[Principles of Survivorship \(REC-F\)](#)

[Staging \(ST-1\)](#)

Clinical Trials: NCCN believes that the best management for any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN Member Institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2017.



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 3.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2017 include:

[REC-8](#)

• The following footnotes removed:

- ▶ Meyerhardt JA, Mangu PB, et al. American Society of Clinical Oncology. J Clin Oncol 2013 Dec 10;31(35):4465-4470.
- ▶ CT scan may be useful for patients at high risk for recurrence (eg, lymphatic or venous invasion by tumor; poorly differentiated tumors).

[REC-10](#)

• The following footnote removed: There are limited data to support a specific treatment regimen in this setting. (also applies to REC-11)

[REC-11](#)

• Footnote “cc” modified: ~~Evidence increasingly suggests that BRAF V600E mutation makes response to panitumumab or cetuximab, as single agents or in combination with cytotoxic chemotherapy,~~ highly unlikely. (also applies REC-E 6 of 10)

[REC-C 1 of 2](#)

• The following footnote removed: NCCN recommends limiting chemotherapy orders to 24-hour units (ie, 1200 mg/m²/d NOT 2400 mg/m² over 48 hours) to minimize medication errors. (also applies to REC-E 7 of 10, REC-E 8 of 10, REC-E 9 of 10)

[REC-E 1 of 10](#)

• Initial Therapy: 5-FU/leucovorin clarified with the addition of “infusional preferred”.

[REC-E 6 of 10](#)

• The following footnotes removed:

- ▶ Discontinuation of oxaliplatin should be strongly considered from FOLFOX or CAPEOX after 3–4 months of therapy (or sooner if significant neurotoxicity develops ≥ grade 2) with other drugs maintained (fluoropyrimidine + bevacizumab) until time of tumor progression. Oxaliplatin may be reintroduced if it was discontinued previously for neurotoxicity rather than disease progression. Tournigand C, Cervantes A, Figer A, et al. OPTIMOX1: A randomized study of FOLFOX4 or FOLFOX7 with oxaliplatin in a stop-and-go fashion in advanced colorectal cancer - A GERCOR Study. J Clin Oncol 2006;24:394-400. There are no data to support the routine use of Ca/Mg infusion to prevent oxaliplatin-related neurotoxicity and therefore it should not be done.
- ▶ The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine. The relative efficacy of CAPEOX with lower starting doses of capecitabine has not been addressed in large-scale randomized trials.
- ▶ There is an increased risk of stroke and other arterial events, especially in those aged ≥65 years. The use of bevacizumab may interfere with wound healing.
- ▶ Combination therapy involving cytotoxics, anti-EGFRs, and anti-VEGFs is not recommended. Hecht JR, Mitchell E, Chidiac T, et al. A randomized phase IIIB trial of chemotherapy, bevacizumab, and panitumumab compared with chemotherapy and bevacizumab alone for metastatic colorectal cancer. J Clin Oncol 2009;27:672-80. Tol J, Koopman M, Cats A, et al. Chemotherapy, bevacizumab, and cetuximab in metastatic colorectal cancer. N Engl J Med 2009;360(6):563-572.
- ▶ There is a preponderance of data to suggest lack of activity of cetuximab and panitumumab in initial therapy for patients whose primary tumors originated on the right side of the colon.
- ▶ Infusional 5-FU is preferred.
- ▶ Patients with diminished creatinine clearance may require dose modification of capecitabine.
- ▶ EGFR testing has no demonstrated predictive value; therefore, routine EGFR testing is not recommended. No patient should be included or excluded from cetuximab or panitumumab therapy on the basis of EGFR test results.
- ▶ There are no data, nor is there a compelling rationale, to support the use of panitumumab after clinical failure on cetuximab, or the use of cetuximab after clinical failure on panitumumab. As such, the use of one of these agents after therapeutic failure on the other is not recommended.
- ▶ Single-agent or combination therapy with capecitabine, mitomycin, or gemcitabine has not been shown to be effective in this setting.
- Footnote “5” modified: Irinotecan should be used with caution ~~and with decreased doses in patients~~ with Gilbert’s disease or elevated serum bilirubin. There is a commercially available test for UGT1A1. Guidelines for use in clinical practice have not been established.

UPDATES



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 2.2017 of the NCCN Guidelines for Rectal Cancer from Version 1.2017 include:

[REC-7](#)

- Qualifier "(not recommended for T4 tumors)" add to "Short-course RT".

[REC-9](#)

- Isolated pelvic/anastomotic recurrence: Capecitabine + RT added as a treatment option.

Updates in Version 1.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2016 include:

General

- Imaging recommendations clarified with anatomy and contrast.
- Advanced or Metastatic Disease: "Chemotherapy" changed throughout to "Systemic therapy."
- T3-4, N0 or T any, N1-2, Medical contraindication to combined modality therapy (was REC-6) page was deleted.

[MS-1](#)

- The discussion section was updated to reflect the changes in the algorithm.

[REC-2](#)

- Workup: Pelvic MRI with contrast noted as "preferred."
- Workup: "Proctoscopy" removed.

[REC-3](#)

- This page now addresses adjuvant treatment after local excision.
- The option for "transabdominal resection after chemo/RT" was removed and replaced with the options of "Consider observation if complete response or transabdominal resection or consider FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine."
- The following footnotes were added: (also applies to REC-4)
 - ▶ Footnote "o": "Imaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection." (Also applies to REC-5, REC-6, REC-10, and REC-11)
 - ▶ Footnote "p": "A benefit for the addition of oxaliplatin to 5-FU/leucovorin in patients age 70 and older has not been proven."

[REC-4](#)

- This page now addresses adjuvant treatment after transabdominal resection (previously on page REC-3)
- pT3-4, N0, M0: the following treatment options were added: "Observation or Chemotherapy: FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine."

[REC-5](#)

- Adjuvant treatment following chemo/RT: FLOX removed as a treatment option.
- Footnote removed: "Postoperative therapy is indicated in all patients who receive preoperative therapy, regardless of the surgical pathology results."

[REC-6](#)

- The combination regimens with bevacizumab, cetuximab, or panitumumab were removed, along with their corresponding footnotes.
- Neoadjuvant Treatment
 - ▶ Short-course RT was added as a treatment option except for T4 tumors. Further therapy includes staged or synchronous resection (preferred) and/or local therapy for metastases and resection of rectal lesion followed by "FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine."

UPDATES



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 1.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2016 include:

[REC-6](#)

- **Primary Treatment:** Short-course RT added as a treatment option except for T4 tumors.
- **Adjuvant Treatment following Chemo/RT and resection:**
 - ▶ Recommendations changed from "Chemotherapy as per REC-E" to "FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine."
- **Footnote "u" clarified:** "Resection is preferred over locally ablative procedures (eg, image-guided ablation or SBRT). However, these local techniques can be considered for liver *or lung* oligometastases." (also applies to REC-10)

[REC-7](#)

- **Primary treatment:** "Laser recanalization" removed.
- "Short-course RT" added as a treatment option.

[REC-8](#)

- **Surveillance options differentiated based on stage.**
 - ▶ The following added: "Stage I with full surgical staging" with colonoscopic surveillance
- **Other stages:** "routinely" removed from PET/CT scan is not recommended.
- **Footnote "w":** reference updated to "Meyerhardt JA, Mangu PB, et al. American Society of Clinical Oncology. J Clin Oncol 2013 Dec 10;31(35):4465-4470."

[REC-10](#)

- **Primary Treatment**
 - ▶ **No previous chemotherapy:** Neoadjuvant chemotherapy with FLOX or Capecitabine or 5-FU/leucovorin changed from a category 2A to a category 2B.
 - ▶ **Previous chemotherapy:** Neoadjuvant chemotherapy recommendations changed from "Chemotherapy as per REC-E" to "FOLFOX (preferred) or CAPEOX (preferred) or FLOX or Capecitabine or 5-FU/leucovorin."
- **Adjuvant Treatment**
 - ▶ **No growth on neoadjuvant chemotherapy:** "Observation" added as an option.
 - ▶ **Growth on neoadjuvant chemotherapy:** "Systemic chemotherapy ± biologic therapy ([REC-E](#)) (category 2B for biologic therapy)."
 - ▶ **Previous chemotherapy:** Adjuvant therapy after resection clarified as "Systemic chemotherapy ± biologic therapy ([REC-E](#)) (category 2B for biologic therapy)."
- **Surveillance added with a link to REC-8.** (also applies to REC-11)
- **Footnote "dd" added:** "There are limited data to support a specific treatment regimen in this setting." (also applies to REC-11)

[REC-11](#)

- **Adjuvant treatment clarified as** "Systemic chemotherapy ± biologic therapy ([REC-E](#)) (category 2B for biologic therapy)."
- **For patients with deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) tumors,** nivolumab or pembrolizumab were added as treatment options in subsequent therapy for patients appropriate for intensive therapy.
- **Surveillance moved to REC-8.**



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 1.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2016 include:

[REC-A 1 of 6](#)

- Endoscopically Removed Malignant Polyps, bullet 3 modified with addition of 3rd sentence: "In several studies, tumor budding has been shown to be an adverse histological feature associated with adverse outcome and may preclude polypectomy as an adequate treatment of endoscopically removed malignant polyps."

[REC-A 2 of 6](#)

- Pathologic Stage, sub-bullet 4 modified: "Status of proximal, distal, ~~and circumferential (radial) margins~~, and mesenteric margins."
- Pathologic Stage, sub-bullet 10 modified: ~~Extranodal~~ Tumor deposits. (also applies to REC-A 3 of 6)
- Pathologic stage, sub-bullet removed: "A positive circumferential resection margin (CRM) has been defined as ≤ 1 mm."

[REC-A 4 of 6](#)

- Lymph node evaluation, first sentence modified: "The AJCC and CAP recommend examination of a minimum of 12 lymph nodes to accurately identify stage II colorectal cancer."

[REC-A 5 of 6](#)

• Microsatellite Instability (MSI) and Mismatch Repair (MMR) Testing

- ▶ Bullet 1 changed from "Lynch syndrome tumors screening (ie, IHC for MMR or PCR for MSI)* should be performed for all patients with colorectal cancer diagnosed at age ≤ 70 y and also those >70 y who meet the Bethesda guidelines. See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal" to "Universal MMR or MSI testing is recommended in all patients with a personal history of colon or rectal cancer. [See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)"
- ▶ Bullet 3 modified: "~~MMR or MSI testing should also be performed for all patients with stage II disease, because Stage II MSI-H patients may have a good prognosis and do not benefit from 5-FU adjuvant therapy.~~"
- ▶ Bullet 4 added: "MMR or MSI testing should be performed only in CLIA-approved laboratories."
- ▶ Bullet removed: "MMR or MSI testing should also be performed for all patients with metastatic disease."

[REC-B 2 of 3](#)

- Liver, bullet 7 changed from "Some institutions use arterially directed embolic therapy (category 3) in highly selected patients with chemotherapy-resistant/-refractory disease, without obvious systemic disease, with predominant hepatic metastases" to "Arterially directed catheter therapy, and in particular yttrium 90 microsphere selective internal radiation, is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases."
- Lung, bullet 5 added: "Ablative techniques may be considered alone or in conjunction with resection. All original sites of disease need to be amenable to ablation or resection."

[REC-C 1 of 2](#)

- CAPEOX dosing modified with the removal of timing for oxaliplatin over 2 hours.
- Footnote "*" modified with updated reference.
- Footnote "#" added for capecitabine: "The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine."



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 1.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2016 include:

[REC-D](#)

- **Bullet 7, first sentence modified:** "Intraoperative radiation therapy (IORT), if available, ~~should~~*may* be considered for very close or positive margins after resection, as an additional boost, especially patients with T4 or recurrent cancers."
- **Bullet 10 added:** "Arterially directed catheter therapy, and in particular yttrium 90 microsphere selective internal radiation, is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases."
- **Bullet 11, first sentence modified:** "In patients with a limited number of liver or lung metastases, radiotherapy *to the metastatic site* can be considered in highly selected cases or in the setting of a clinical trial. Radiotherapy should not be used in the place of surgical resection. Radiotherapy should be delivered in a highly conformal manner. The techniques can include 3-D conformal radiation therapy, IMRT, or stereotactic body radiation therapy (SBRT) ~~(category 3).~~"

REC-E

- The pages for the Continuum of Care - Systemic ~~Chemo~~Therapy for Advanced or Metastatic Disease reorganized to address first-line therapy and then subsequent therapy based on previous therapy.
- For patients with deficient mismatch repair (dMMR) or microsatellite instability-high (MSI-H) tumors, nivolumab or pembrolizumab were added as treatment options in subsequent therapy for patients appropriate for intensive therapy.

[REC-E 2 of 10](#)

- This page now addresses subsequent therapy options for patients receiving oxaliplatin-based therapy without irinotecan as initial therapy.

[REC-E 3 of 10](#)

- This page now addresses subsequent therapy options for patients receiving irinotecan-based therapy without oxaliplatin as initial therapy.

[REC-E 4 of 10](#)

- This page now addresses subsequent therapy options for patients receiving FOLFOXIRI as initial therapy.

[REC-E 5 of 10](#)

- This page now addresses subsequent therapy options for patients receiving fluoropyrimidine without oxaliplatin or irinotecan as initial therapy.

[REC-E 6 of 10](#)

- **Footnote "2" modified:** "~~CT with contrast or MRI is recommended. Chest/Abdominal/Pelvic CT with contrast or Chest CT and Abdominal/Pelvic MRI with contrast~~ to monitor progress of therapy. PET/CT should not be used."
- **Footnote "9" added:** "There is a preponderance of data to suggest lack of activity of cetuximab and panitumumab in initial therapy for patients whose primary tumors originated on the right side of the colon."
- **Footnote "17" modified:** "~~Cetuximab or panitumumab are recommended~~ *is indicated* in combination with irinotecan-based therapy or as a single-agent for patients who cannot tolerate irinotecan."
- **Footnote "20" modified:** "Regorafenib or trifluridine + tipiracil are treatment options for patients who have progressed through all available regimens ~~(eg, KRAS/NRAS mutant or KRAS/NRAS WT with previous exposure to anti-EGFR inhibitor.)~~"



NCCN Guidelines Version 3.2017 Updates

Rectal Cancer

Updates in Version 1.2017 of the NCCN Guidelines for Rectal Cancer from Version 2.2016 include:

[REC-E 7 of 10](#)

- Regimen added: mFOLFOX7 (Oxaliplatin 85 mg/m² IV day 1; Leucovorin 400 mg/m² IV day 1; 5-FU 1200 mg/m²/d x 2 days [total 2400 mg/m² over 46–48 hours]) IV continuous infusion. Repeat every 2 weeks.
- CAPEOX dosing modified with the removal of 850 mg for capecitabine and removal of timing for oxaliplatin over 2 hours.
- Footnote “*” modified with updated reference. (also applies to REC-E 8 of 10)
- Footnote “‡” modified with the removal of the following sentence: "The relative efficacy of CAPEOX with lower starting doses of capecitabine has not been addressed in large-scale randomized trials."

[REC-E 9 of 10](#)

- Dosing recommendations were added for pembrolizumab and nivolumab.

[REC-E 10 of 10](#)

- References 4, 30, 31 added. Reference 18 updated.


[REC-F 1 OF 2](#)

- Management of Late Sequelae of Disease or Treatment
 - ▶ Link added to the NCCN Guidelines for Survivorship.
 - ▶ "Pelvic floor rehabilitation" added as an intervention for chronic diarrhea or incontinence.
 - ▶ "Oxaliplatin-induced neuropathy" added with intervention: "Consider duloxetine for painful neuropathy."
 - ▶ "Fatigue" added with intervention: "Encourage physical activity, energy conservation measures."
 - ▶ "Potential for pelvic fractures/decreased bone density after pelvic radiation" added with intervention: "Consider bone density monitoring."
- "Survivorship Care Planning" added with the following revisions:
 - ▶ Addition: "The oncologist and primary care provider should have defined roles in the surveillance period, with roles communicated to patient."
 - ▶ Removal: "Prescription for Survivorship and Transfer of Care to Primary Care Physician (If primary physician will be assuming cancer surveillance responsibilities):"
 - ▶ Addition: "Develop survivorship care plan that includes:"
 - ◊ Sub-bullet 1 modified: ~~"Include Overall summary of treatment, including all surgeries, radiation treatments, and chemotherapy received."~~
 - ◊ Sub-bullet 2 modified: ~~"Description of possible expected time Describe possible clinical course, including expected time to resolution of acute toxicities, long-term effects of treatment, and possible late sequelae of treatment."~~
 - ◊ Sub-bullet 3 modified: ~~"Include Surveillance recommendations."~~
 - ◊ Sub-bullet 5 added: "Health behavior recommendations."
- Counseling Regarding Healthy Lifestyle and Wellness
 - ▶ Bullet 3 modified: "Consume a healthy diet with emphasis on plant sources. *Diet recommendations may be modified based on severity of bowel dysfunction.*"
 - ▶ Bullet 4 added: "Consider low-dose aspirin."



NCCN Guidelines Version 3.2017

Rectal Cancer



^aAll patients with rectal cancer should be counseled for family history. Patients with suspected Lynch syndrome, familial adenomatous polyposis (FAP), and attenuated FAP, see the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#).

^bFor melanoma histology, see the [NCCN Guidelines for Melanoma](#).

^cConfirm the presence of invasive cancer (pT1). pTis has no biological potential to metastasize.

^dIt has not been established if molecular markers are useful in treatment determination (predictive markers) and prognosis. College of American Pathologists Consensus Statement 1999. Prognostic factors in colorectal cancer. Arch Pathol Lab Med 2000;124:979-994.

^e[See Principles of Pathologic Review \(REC-A\)](#) - Endoscopically removed malignant polyp.

^fObservation may be considered, with the understanding that there is significantly greater incidence of adverse outcomes (residual disease, recurrent disease, mortality, or hematogenous metastasis, but not lymph node metastasis) than polypoid malignant polyps. [See Principles of Pathologic Review \(REC-A\)](#) - Endoscopically removed malignant polyp.

^g[See Principles of Surgery \(REC-B\)](#).


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer



^aAll patients with rectal cancer should be counseled for family history. Patients with suspected Lynch syndrome, familial adenomatous polyposis (FAP), and attenuated FAP, see the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#).

^bFor melanoma histology, see the [NCCN Guidelines for Melanoma](#).

^gSee [Principles of Surgery \(REC-B\)](#).

^hCT should be with IV and oral contrast. Consider abdominal/pelvic MRI with MRI contrast plus a non-contrast chest CT if either CT of abd/pelvis is inadequate or if patient has a contraindication to CT with IV contrast.

ⁱPET/CT does not supplant a contrast-enhanced diagnostic CT scan. PET/CT should only be used to evaluate an equivocal finding on a contrast-enhanced CT scan or in patients with strong contraindications to IV contrast.

^jT1-2, N0 should be based on assessment of endorectal ultrasound or MRI.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

FINDINGS AFTER TRANSANAL EXCISION

ADJUVANT TREATMENT^{m,n,o} (6 MO PERIOPERATIVE TREATMENT PREFERRED)^p



^gSee Principles of Surgery (REC-B).

^kHigh-risk features include positive margins, lymphovascular invasion, poorly differentiated tumors, or sm3 invasion.

^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

^mSee Principles of Adjuvant Therapy (REC-C).

ⁿSee Principles of Radiation Therapy (REC-D).

^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^pA benefit for the addition of oxaliplatin to 5-FU/leucovorin in patients age 70 and older has not been proven.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer

FINDINGS AFTER TRANSABDOMINAL RESECTION FOR cT1-2, N0

ADJUVANT TREATMENT^{m,n,o} (6 MO PERIOPERATIVE TREATMENT PREFERRED)^p

pT1-2, N0, M0

Observe

pT3-4, N0, M0

Observation
or
Chemotherapy:
FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine
or
FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine, then
capecitabine/RT (preferred) or infusional 5-FU/RT (preferred) or bolus 5-FU/leucovorin/RT,^l
then FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine
or
Infusional 5-FU/RT (preferred) or capecitabine/RT (preferred) or bolus 5-FU/leucovorin/RT^l
followed by FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine

pT1-4, N1-2

FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine, then
capecitabine/RT (preferred) or infusional 5-FU/RT (preferred) or bolus 5-FU/leucovorin/
RT,^l then FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or capecitabine
or
Infusional 5-FU/RT (preferred) or capecitabine/RT (preferred) or bolus 5-FU/leucovorin/
RT^l followed by FOLFOX (preferred) or CAPEOX (preferred) or 5-FU/leucovorin or
capecitabine

[Surveillance](#)
(See REC-8)

^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

^mSee [Principles of Adjuvant Therapy \(REC-C\)](#).

ⁿSee [Principles of Radiation Therapy \(REC-D\)](#).

^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^pA benefit for the addition of oxaliplatin to 5-FU/leucovorin in patients age 70 and older has not been proven.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer



^gSee Principles of Surgery (REC-B).

^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

^mSee Principles of Adjuvant Therapy (REC-C).

ⁿSee Principles of Radiation Therapy (REC-D).

^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^qEvaluation for short-course RT should be in a multidisciplinary setting, with a discussion of the need for down-staging and the possibility of long-term toxicity.

^rFernandez-Martos C, Pericay C, Aparicio J, et al: Phase II, randomized study of concomitant chemoradiotherapy followed by surgery and adjuvant capecitabine plus oxaliplatin (CAPOX) compared with induction CAPOX followed by concomitant chemoradiotherapy and surgery in magnetic resonance imaging-defined, locally advanced rectal cancer: Grupo cancer de recto 3 study. J Clin Oncol 2010;28:859-865.

Cercek A, Goodman KA, Hajj C, et al. Neoadjuvant chemotherapy first, followed by chemoradiation and then surgery, in the management of locally advanced rectal cancer. J Natl Compr Canc Netw 2014;12:513-519.

^sFOLFOXIRI is not recommended in this setting.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer



^gSee Principles of Surgery (REC-B).

^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

^mSee Principles of Adjuvant Therapy (REC-C).

ⁿSee Principles of Radiation Therapy (REC-D).

^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^qEvaluation for short-course RT should be in a multidisciplinary setting, with a discussion of the need for down-staging and the possibility of long-term toxicity.

[†]Determination of tumor gene status for RAS (KRAS and NRAS) and BRAF. Determination of tumor MMR or MSI status (if not previously done). See Principles of Pathologic Review (REC-A 5 of 6) - KRAS, NRAS, and BRAF Mutation Testing and Microsatellite Instability (MSI) or Mismatch Repair (MMR) Testing.

^uResection is preferred over locally ablative procedures (eg, image-guided ablation or SBRT). However, these local techniques can be considered for liver or lung oligometastases (REC-B and REC-D).

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

CLINICAL STAGE

PRIMARY TREATMENT



^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

ⁿ[See Principles of Radiation Therapy \(REC-D\)](#).

^qEvaluation for short-course RT should be in a multidisciplinary setting, with a discussion of the need for down-staging and the possibility of long-term toxicity.

[†]Determination of tumor gene status for RAS (KRAS and NRAS) and BRAF. Determination of tumor MMR or MSI status (if not previously done). [See Principles of Pathologic Review \(REC-A 5 of 6\)](#) - KRAS, NRAS, and BRAF Mutation Testing and Microsatellite Instability (MSI) or Mismatch Repair (MMR) Testing.

^v[See Systemic Therapy for Advanced or Metastatic Disease \(REC-E\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer

SURVEILLANCE

Stage I with full surgical staging →

- Colonoscopy at 1 y
 - ▶ If advanced adenoma, repeat in 1 y
 - ▶ If no advanced adenoma,^w repeat in 3 y, then every 5 y^x

Stage II, III →

- History and physical every 3–6 mo for 2 y, then every 6 mo for a total of 5 y
- CEA^y every 3–6 mo for 2 y, then every 6 mo for a total of 5 y
- Chest/abdominal/pelvic CT^h every 6–12 mo (category 2B for frequency <12 mo) for a total of 5 y
- Colonoscopy in 1 y except if no preoperative colonoscopy due to obstructing lesion, colonoscopy in 3–6 mo
 - ▶ If advanced adenoma, repeat in 1 y
 - ▶ If no advanced adenoma,^w repeat in 3 y, then every 5 y^x
- Proctoscopy (with EUS or MRI with contrast) every 3–6 mo for the first 2 y, then every 6 mo for a total of 5 y (for patients treated with transanal excision only)
- PET/CT scan is not recommended
- See [Principles of Survivorship \(REC-F\)](#)

Stage IV →

- History and physical every 3–6 mo for 2 y, then every 6 mo for a total of 5 y
- CEA^y every 3–6 mo for 2 y, then every 6 mo for a total of 5 y
- Chest/abdominal/pelvic CT^h every 3–6 mo (category 2B for frequency <6 mo) x 2 y, then every 6–12 mo for a total of 5 y
- Colonoscopy in 1 y except if no preoperative colonoscopy due to obstructing lesion, colonoscopy in 3–6 mo
 - ▶ If advanced adenoma, repeat in 1 y
 - ▶ If no advanced adenoma,^w repeat in 3 y, then every 5 y^x
- Proctoscopy (with EUS or MRI with contrast) every 3–6 mo for the first 2 y, then every 6 mo for a total of 5 y (for patients treated with transanal excision only)
- PET/CT scan is not recommended
- See [Principles of Survivorship \(REC-F\)](#)

Serial CEA elevation or documented recurrence →

→ [See Workup and Treatment \(REC-9\)](#)

^hCT should be with IV and oral contrast. Consider abdominal/pelvic MRI with MRI contrast plus a non-contrast chest CT if either CT of abd/pelvis is inadequate or if patient has a contraindication to CT with IV contrast.

^wVillous polyp, polyp >1 cm, or high-grade dysplasia.

^xRex DK, Kahi CJ, Levin B, et al. Guidelines for colonoscopy surveillance after cancer resection: a consensus update by the American Cancer Society and the US Multi-Society Task Force on Colorectal Cancer. *Gastroenterology* 2006;130(6):1865-71.

^yIf patient is a potential candidate for resection of isolated metastasis.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer



⁹See Principles of Surgery (REC-B).

^lBolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

ⁿSee Principles of Radiation Therapy (REC-D).

^tDetermination of tumor gene status for RAS (KRAS and NRAS) and BRAF. Determination of tumor MMR or MSI status (if not previously done). See Principles of Pathologic Review (REC-A 5 of 6) - KRAS, NRAS, and BRAF Mutation Testing and Microsatellite Instability (MSI) or Mismatch Repair (MMR) Testing.


^zPatients should be evaluated by a multidisciplinary team including surgical consultation for potentially resectable patients.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer



^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^uResection is preferred over locally ablative procedures (eg, image-guided ablation or SBRT). However, these local techniques can be considered for liver or lung oligometastases (REC-B and REC-D).

^{aa}Hepatic artery infusion ± systemic 5-FU/leucovorin (category 2B) is also an option at institutions with experience in both the surgical and medical oncologic aspects of this procedure.

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer

UNRESECTABLE METACHRONOUS METASTASES

PRIMARY TREATMENT

ADJUVANT TREATMENT^o (6 MO PERIOPERATIVE TREATMENT PREFERRED)

• Previous adjuvant FOLFOX/CAPEOX within past 12 months

FOLFIRI ± (bevacizumab [preferred] or ziv-aflibercept or ramucirumab)^{cc}
or
Irinotecan ± (bevacizumab [preferred] or ziv-aflibercept or ramucirumab)^{bb}
or
FOLFIRI + (cetuximab or panitumumab) (KRAS/NRAS WT gene only)^{t,cc}
or
Irinotecan ± (Cetuximab or panitumumab) (KRAS/NRAS WT gene only)^{t,cc}
or
(Nivolumab or pembrolizumab) (dMMR/MSI-H only)

• Previous adjuvant FOLFOX/CAPEOX >12 months
• Previous 5-FU/LV or capecitabine
• No previous chemotherapy

Systemic therapy (REC-E)

Re-evaluate for conversion to resectable^g every 2 mo if conversion to resectability is a reasonable goal

Converted to resectable

Resection^{aa}

Remains unresectable

Systemic therapy (REC-E)

Systemic therapy ± biologic therapy (REC-E) (category 2B for biologic therapy) or Observation

See Surveillance (REC-8)

^gSee Principles of Surgery (REC-B).

^oImaging (Chest/Abdomen/Pelvic CT with contrast) to be performed prior to adjuvant treatment to assess response to primary therapy or resection.

^tDetermination of tumor gene status for RAS (KRAS and NRAS) and BRAF. Determination of tumor MMR or MSI status (if not previously done). See Principles of Pathologic Review (REC-A 5 of 6) - KRAS, NRAS, and BRAF Mutation Testing and Microsatellite Instability (MSI) or Mismatch Repair (MMR) Testing.

^{aa}Hepatic artery infusion ± systemic 5-FU/leucovorin (category 2B) is also an option at institutions with experience in both the surgical and medical oncologic aspects of this procedure.

^{bb}Bevacizumab is the preferred anti-angiogenic agent based on toxicity and/or cost.

^{cc}BRAF V600E mutation makes response to panitumumab or cetuximab highly unlikely.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF PATHOLOGIC REVIEW (1 of 6)

Endoscopically Removed Malignant Polyps

- A malignant polyp is defined as one with cancer invading through the muscularis mucosae and into the submucosa (pT1). pTis is not considered to be a “malignant polyp.”
- Favorable histopathologic features: grade 1 or 2, no angiolymphatic invasion, and negative margin of resection. There is no consensus as to the definition of what constitutes a positive margin of resection. A positive margin has been defined as: 1) tumor <1 mm from the transected margin; 2) tumor <2 mm from the transected margin; and 3) tumor cells present within the diathermy of the transected margin.¹⁻⁴
- Unfavorable histologic features grade 3 or 4, angiolymphatic invasion, or a “positive margin.” See above for definition of a positive margin. In several studies, tumor budding has been shown to be an adverse histological feature associated with adverse outcome and may preclude polypectomy as an adequate treatment of endoscopically removed malignant polyps.
- There is controversy as to whether malignant colorectal polyps with a sessile configuration can be successfully treated by endoscopic removal. The literature seems to indicate that endoscopically removed sessile malignant polyps have a significantly greater incidence of adverse outcome (residual disease, recurrent disease, mortality, or hematogenous metastasis, but not lymph node metastasis) than do polypoid malignant polyps. However, when one closely looks at the data, configuration by itself is not a significant variable for adverse outcome, and endoscopically removed malignant sessile polyps with grade I or II histology, negative margin, and no lymphovascular invasion can be successfully treated with endoscopic polypectomy.³⁻⁷

Transanal Excision

- Favorable histopathologic features: <3 cm size, T1, grade I or II, no lymphatic or venous invasion, or negative margins.^{8,9}
- Unfavorable histopathologic features: >3 cm in size, T1, with grade III, or lymphovascular invasion, positive margin, or sm3 depth of tumor invasion.⁸⁻¹⁰

Rectal Cancer Appropriate for Resection

- Histologic confirmation of primary malignant rectal neoplasm.

[See Pathologic Stage on REC-A 2 of 6](#)

[See Lymph Node Evaluation on REC-A 4 of 6](#)

[See KRAS, NRAS, and BRAF Mutation Testing REC-A 5 of 6](#)

[See references on REC-A 6 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGIC REVIEW (2 of 6)****Pathologic Stage**• **The following parameters should be reported:**

- ▶ **Grade of the cancer**
- ▶ **Depth of penetration (T), the T stage, is based on viable tumor. Acellular mucin pools are not considered to be residual tumor in those cases treated with neoadjuvant therapy.**
- ▶ **Number of lymph nodes evaluated and number positive (N). Acellular mucin pools are not considered to be residual tumor in those cases treated with neoadjuvant therapy.**
- ▶ **Status of proximal, distal, circumferential (radial), and mesenteric margins.¹¹⁻¹²**
- ▶ **Circumferential resection margin (CRM)¹³⁻¹⁷**
- ▶ **Neoadjuvant treatment effect^{15,16,18,19}**
- ▶ **Lymphovascular invasion^{15,16,20}**
- ▶ **Perineural invasion (PNI)²¹⁻²³**
- ▶ **Tumor deposits²⁴⁻²⁵**

• **CRM - A positive CRM is defined as tumor \leq 1 mm from the margin. This assessment includes both tumor within a lymph node as well as direct tumor extension. However, if CRM positivity is based solely on intranodal tumor, it should be stated in the pathology report. A positive CRM is a more powerful predictor of local recurrence in patients treated with neoadjuvant therapy. A positive CRM secondary to lymph node metastasis in some studies has been associated with lower recurrence rates than by direct extension.¹³⁻¹⁷**

• **Neoadjuvant treatment effect - The most recent College of American Pathologists (CAP) Guidelines on examination specimens of the rectum and the AJCC Cancer Staging Manual, Seventh Edition require commenting on treatment effect after neoadjuvant therapy. The minimum requirement is:**

- ▶ **Treatment effect present.**
- ▶ **No definitive response identified.**

The system used to grade tumor response as recommended by the AJCC Cancer Staging Manual, 7th Edition and the CAP Guidelines is that as modified from Ryan R, et al. *Histopathology* 2005;47:141-146.

- ▶ **0 - Complete response: No remaining viable cancer cells.**
- ▶ **1 - Moderate response: Only small clusters or single cancer cells remaining.**
- ▶ **2 - Minimal response: Residual cancer remaining, but with predominant fibrosis.**
- ▶ **3 - Poor response: Minimal or no tumor kill; extensive residual cancer.**

According to CAP, it is optional to grade the tumor response to treatment. However, the NCCN Rectal Cancer Guidelines Panel recommends grading tumor response. Other grading systems that are used are referenced.^{15,16,18,19}

[See Pathologic Stage continued on REC-A 3 of 6](#)

[See Endoscopically Removed Malignant Polyps, Rectal Cancer Appropriate for Resection on REC-A 1 of 6](#)

[See Lymph Node Evaluation on REC-A 4 of 6](#)

[See KRAS, NRAS, and BRAF Mutation Testing REC-A 5 of 6](#)

[See references on REC-A 6 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF PATHOLOGIC REVIEW (3 of 6)

Pathologic Stage (continued)

- **Perineural invasion (PNI)** - The presence of PNI is associated with a significantly worse prognosis. In multivariate analysis, PNI has been shown to be an independent prognostic factor for cancer-specific, overall, and disease-free survival. For stage II rectal cancer, those with PNI have a significantly worse 5-year disease-free survival compared to those without PNI (29% vs. 82% [$P = .0005$]). In stage III rectal cancer, those with PNI have a significantly worse prognosis.²¹⁻²³
- **Tumor deposits** - Irregular discrete tumor deposits in pericolic or perirectal fat away from the leading edge of the tumor and showing no evidence of residual lymph node tissue, but within the lymphatic drainage of the primary carcinoma, are considered to be tumor deposits or satellite nodules and are not counted as lymph nodes replaced by tumor. Most examples are due to lymphovascular invasion or, more rarely, PNI. Because these tumor deposits are associated with reduced disease-free and overall survival, their number should be recorded in the surgical pathology report.

[See Endoscopically Removed Malignant Polyps, Rectal Cancer Appropriate for Resection on REC-A 1 of 6](#)

[See Pathologic Stage on REC-A 2 of 6](#)

[See Lymph Node Evaluation on REC-A 4 of 6](#)

[See KRAS, NRAS, and BRAF Mutation Testing REC-A 5 of 6](#)

[See references on REC-A 6 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGIC REVIEW (4 of 6)****Lymph Node Evaluation**

- The AJCC and CAP recommend examination of a minimum of 12 lymph nodes to accurately stage rectal cancer.^{11,12,26} Sampling of 12 lymph nodes may not be achievable in patients who received preoperative chemotherapy. The literature lacks consensus as to what is the minimal number of lymph nodes to accurately identify stage II cancer. The minimal number of nodes has been reported as >7, >9, >13, >20, and >30.²⁶⁻³⁴ Most of these studies have combined rectal and colon cancers and reflect those cases with surgery as the initial treatment. Two studies confined only to rectal cancer have reported 14 and >10 lymph nodes as the minimal number to accurately identify stage II rectal cancer.^{30,33} The number of lymph nodes retrieved can vary with age of the patient, gender, tumor grade, and tumor site.²⁷ For stage II (pN0) colon cancer, if fewer than 12 lymph nodes are initially identified, it is recommended that the pathologist go back to the specimen and resubmit more tissue of potential lymph nodes. If 12 lymph nodes are still not identified, a comment in the report should indicate that an extensive search for lymph nodes was undertaken. The mean number of lymph nodes retrieved from rectal cancers treated with neoadjuvant therapy is significantly less than those treated by surgery alone (13 vs. 19, $P < .05$, 7 vs. 10, $P < .001$).^{35,36} If 12 lymph nodes is considered the number needed to accurately stage stage II tumors, then only 20% of cases treated with neoadjuvant therapy had adequate lymph node sampling.³⁶ To date, the number of lymph nodes needed to accurately stage neoadjuvant-treated cases is unknown. However, it is not known what is the clinical significance of this in the neoadjuvant setting, as postoperative therapy is indicated in all patients who receive preoperative therapy regardless of the surgical pathology results.

Sentinel Lymph Node and Detection of Micrometastasis by Immunohistochemistry

- Examination of the sentinel lymph node allows an intense histologic and/or immunohistochemical investigation to detect the presence of metastatic carcinoma. Studies in the literature have been reported using multiple H & E sections and/or immunohistochemistry (IHC) to detect cytokeratin-positive cells.³⁷⁻³⁹ The AJCC Cancer Staging Manual, Seventh Edition⁴⁰ considers “tumor clusters” <0.2 mm to be isolated tumor cells (pN0) and not metastatic carcinoma. However, some investigators believe that size should not affect the diagnosis of metastatic cancer. They believe that tumor foci that show evidence of growth (eg, glandular differentiation, distension of sinus, stromal reaction) should be diagnosed as a lymph node metastasis regardless of size.^{41,42}
- Some studies have shown that the detection of IHC cytokeratin-positive cells in stage II (N0) colon cancer (defined by H & E) has a worse prognosis, while others have failed to show this survival difference. In these studies, isolated tumor cells were considered to be micrometastases.⁴³⁻⁴⁷
- At the present time the use of sentinel lymph nodes and detection of cancer cells by IHC alone should be considered investigational, and results should be used with caution in clinical management decisions.^{37-39,43-47}

[See Endoscopically Removed Malignant Polyps, Rectal Cancer Appropriate for Resection on REC-A 1 of 6](#)

[See Pathologic Stage on REC-A 2 of 6](#)

[See KRAS, NRAS, and BRAF Mutation Testing REC-A 5 of 6](#)

[See references on REC-A 6 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF PATHOLOGIC REVIEW (5 of 6)*****KRAS*, *NRAS*, and *BRAF* Mutation Testing**

- All patients with metastatic colorectal cancer should have tumor tissue genotyped for *RAS* (*KRAS* and *NRAS*) and *BRAF* mutations. Patients with any known *KRAS* mutation (exon 2 or non-exon 2) or *NRAS* mutation should not be treated with either cetuximab or panitumumab.⁴⁸⁻⁵⁰ *BRAF V600E* mutation makes response to panitumumab or cetuximab highly unlikely.⁵¹⁻⁵³
- Testing for *KRAS*, *NRAS*, and *BRAF* mutations should be performed only in laboratories that are certified under the clinical laboratory improvement amendments of 1988 (CLIA-88) as qualified to perform *high complexity* clinical laboratory (molecular pathology) testing. No specific methodology is recommended (eg, sequencing, hybridization).
- The testing can be performed on formalin-fixed paraffin-embedded tissue. The testing can be performed on the primary colorectal cancers and/or the metastasis, as literature has shown that the *KRAS*, *NRAS*, and *BRAF* mutations are similar in both specimen types.⁵⁴

Evaluation of Mesorectum (TME)

- The pathologist should evaluate the quality (completeness) of the mesorectum (only for low rectal cancer - distal 2/3).⁵⁵⁻⁵⁷

Microsatellite Instability (MSI) or Mismatch Repair (MMR) Testing

- Universal MMR* or MSI* testing is recommended in all patients with a personal history of colon or rectal cancer. [See NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal](#)
- The presence of a *BRAF V600E* mutation in the setting of MLH1 absence would preclude the diagnosis of Lynch syndrome.
- Stage II MSI-H patients may have a good prognosis and do not benefit from 5-FU adjuvant therapy.⁵⁸
- MMR or MSI testing should be performed only in CLIA-approved laboratories.

*IHC for MMR and PCR for MSI are different assays measuring the same biological effect.

[See Endoscopically Removed Malignant Polyps, Rectal Cancer Appropriate for Resection on REC-A 1 of 6](#)

[See Pathologic Stage on REC-A 2 of 6](#)

[See Lymph Node Evaluation on REC-A 4 of 6](#)

[See references on REC-A 6 of 6](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF PATHOLOGIC REVIEW (6 of 6) - References

- 1 Volk EE, Goldblum JR, Petras RE, et al. Management and outcome of patients with invasive carcinoma arising in colorectal polyps. *Gastroenterology* 1995;109:1801-1807.
- 2 Cooper HS, Deppisch LM, Gourley WK, et al. Endoscopically removed malignant colorectal polyps: clinical pathological correlations. *Gastroenterology* 1995;108:1657-1665.
- 3 Ueno H, Mochizuki H, Hashiguchi Y, et al. Risk factors for an adverse outcome in early invasive colorectal carcinoma. *Gastroenterology* 2004;127:385-394.
- 4 Seitz U, Bohnacker S, Seewald S, et al. Is endoscopic polypectomy an adequate therapy for malignant colorectal polyps? Presentation of 114 patients and review of the literature. *Dis Colon Rectum* 2004;47:1789-1797.
- 5 Morson BC, Whiteway JE, Jones EA, et al. Histopathology and prognosis of malignant colorectal polyps treated by endoscopic polypectomy. *Gut* 1984;25:437-444.
- 6 Haggitt RC, Glotzbach RE, Soffer EE, Wruble LD. Prognostic factors in colorectal carcinomas arising in adenomas: implications for lesions removed by endoscopic polypectomy. *Gastroenterology* 1985;89:328-336.
- 7 Netzer P, Binck J, Hammer B, et al. Significance of histological criteria for the management of patients with malignant colorectal polyps. *Scand J Gastroenterol* 1997;323:915-916.
- 8 Hager T, Gall FP, and Hermanek P. Local excision of cancer of the rectum. *Dis Colon Rect* 1983;26:149-151.
- 9 Willett, CG, Tepper JE, Donnelly S, et al. Patterns of failure following local excision and local excision and postoperative radiation therapy for invasive rectal adenocarcinoma. *J Clin Oncol* 1989;7:1003-1008.
- 10 Nascimbeni R, Burgart LJ, Nivatongs S, and Larson DR. Risk of lymph node metastasis in T1 carcinoma of the colon and rectum. *Dis Colon Rectum* 2002;45:2001-2006.
- 11 Compton CC and Greene FL. The staging of colorectal cancer: 204 and beyond. *Cancer J Clin* 2004;54:295-308.
- 12 Compton CC, Fielding LP, Burkhardt LJ, et al. Prognostic factors in colorectal cancer. College of American pathologists consensus statement. *Arch Pathol Lab Med* 2000;124:979-994.
- 13 Nagtegaal ID, Merijnena M, Kranenbarg EK, et al. Circumferential margin involvement is still an important predictive local occurrence in rectal carcinoma. Not one millimeter but two millimeters is the limit. *Am J Surg Pathol* 2002;26:350-357.
- 14 Wibe A, Rendedal PR, Svensson E, et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. *Br J Surgery* 2002;89 327-334.
- 15 Washington MK, Berlin J, Branton P, et al. Protocol for examination of specimens from patients with primary carcinoma of the colon and rectum. *Arch Pathol Lab Med* 2009;133:1539.
- 16 Edge SB, Byrd D, Compton C, et al (eds). *AJCC Cancer Staging Manual 7th Edition*. Springer NY, 2010.
- 17 Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? *J Clin Oncol* 2008;26:303-312.
- 18 Rodel C, Martus P, Papadopoulos T, et al. Prognostic significance of tumor regression after preoperative chemoradiotherapy for rectal cancer. *J Clin Oncol* 2005;23:8688-8696.
- 19 Gavioli M, Luppi G, Losi L, et al. Incidence and clinical impact of sterilized disease and minimal residual disease after preoperative radiochemotherapy for rectal cancer. *Dis Colon Rectum* 2005;48:1851-1857.
- 20 Nissan A, Stojadinovic A, Shia J, et al. Predictors of recurrence in patients with T2 and early T3, N0 adenocarcinoma of the rectum treated by surgery alone. *J Clin Oncol* 2006;24:4078-4084.
- 21 Liebig C, Ayala G, Wilks J, et al. Perineural invasion is an independent predictor of outcome in colorectal cancer. *J Clin Oncol* 2009;27:5131-5137.
- 22 Fujita S, Shimoda T, Yoshimura K, et al. Prospective evaluation of prognostic factors in patients with colorectal cancer undergoing curative resection. *J Surg Oncol* 2003;84:127-131.
- 23 Quah HM. Identification of patients with high risk stage II colon cancer for adjuvant therapy. *Dis Colon Rect* 2008;51:53-507.
- 24 Ueno H, Mochizuki H, Hashiguchi Y, et al. Extramural cancer deposits without nodal structure in colorectal cancer: optimal categorization for prognostic staging. *J Clin Pathol* 2007;117:287-294.
- 25 Lo DS, Pollett A, Siu LL, et al. Prognostic significance of mesenteric tumor nodules in patients with stage III colorectal cancer. *Cancer* 2008;112:50-54.
- 26 Sobin LH and Green EFL. TNM classification. Clarification of number of regional lymph nodes for PNO. *Cancer* 2001;92:452.
- 27 Sarli L, Bader G, Lusco D, et al. Number of lymph nodes examined and prognosis of TNM stage II colorectal cancer. *European Journal of Cancer* 2005;41:272-279.
- 28 Chaplin S, Scerottini G-P, Bosman FT, et al. For patients with Duke's B (TNM stage II) colorectal carcinoma, examination of six or fewer lymph nodes is related to poor prognosis. *Cancer* 1998;83:666-72.
- 29 Maurel J, Launoy G, Grosclaude P, et al. Lymph node harvest reporting in patients with carcinoma of the large bowel. A French population-based study. *Cancer* 1998;82:1482-6.
- 30 Pocard M, Panis Y, Malassagane B, et al. Assessing the effectiveness of mesorectal excision in rectal cancer. *Dis Colon Rectum* 1998;41:839-845.
- 31 Joseph NE, Sigurdson ER, Hamlin AL, et al. Accuracy of determining nodal negativity in colorectal cancer on the basis of number of nodes retrieved on resection. *Ann of Surg Oncol* 2003;10:213-218.
- 32 Goldstein NS. Lymph node recurrences from 2427 PT3 colorectal resection specimens spanning 45 years. Recommendations for a minimum number of recovered lymph nodes based on predictive probabilities. *Am J Surg Pathol* 2002;26:179-189.
- 33 Tepper JE, O'Connell MJ, Niedzwiecki D, et al. Impact of number of nodes retrieved on outcome in patients with rectal cancer. *J Clin Oncol* 2001;19:157-162.
- 34 Scott KWM and Grace RH. Detection of lymph node metastasis and colorectal carcinoma before and after fat clearance. *Br J Surg* 1989;76:1165-1167.
- 35 Wichmann MW, Mollar C, Meyer G, et al. Effect of pre-operative radiochemotherapy on lymph node retrieval after resection of rectal cancer. *Arch Surg* 2002;137:206-210.
- 36 Baxter NN, Morris AM, Rothenberger DA, and Tepper JE. Impact of pre-operative radiation for rectal cancer on subsequent lymph node evaluation: population based analysis. *Int J Radiation Oncology Biol Phys* 2005;61:426-431.
- 37 Turner RR, Nora DT, Trochas D, and Bilchik AJ. Colorectal carcinoma in nodal staging. Frequency and nature of cytokeratin positive cells in sentinel and nonsentinel lymph nodes. *Arch Pathol Lab Med* 2003;127:673-679.
- 38 Wood TF, Nora DT, Morton DL, et al. One hundred consecutive cases of sentinel node mapping in early colorectal carcinoma. Detection of missed micrometastasis. *J Gastrointest Surg* 2002;6:322-330.
- 39 Wiese DA, Sha S, Badin J, et al. Pathological evaluation of sentinel lymph nodes in colorectal carcinoma. *Arch Pathol Lab Med* 2000;124:1759-1763.
- 40 *AJCC Cancer Staging Manual*, 7th ed. Edge SB, Byrd D, Compton CC, et al. (editors) Springer, New York, 2010.
- 41 Jass JB, O'Brien MJ, Riddell RH, Snover DC, on behalf of the Association of Directors of Anatomic and Surgical Pathology. Recommendations for the reporting of surgically resected specimens of colorectal carcinoma. *Hum Pathol* 2007;38:537-545.
- 42 Hermanek P, Hutter RVP, Sobin LH, Wittekind CH. Classification of isolated tumor cells and micrometastasis. *Cancer* 1999;86:2668-73.
- 43 Noura S, Yamamoto H, Ohnishi T, et al. Comparative detection of lymph node micrometastasis of stage II colorectal cancer by reverse transcriptase polymerase chain reaction in immunohistochemistry. *J Clin Oncol* 2002;20:4232-4241.
- 44 Yasuda K, Adachi Y, Shiraishi N, et al. Pattern of lymph node micrometastasis and prognosis of patients with colorectal cancer. *Ann Surg Oncol* 2001;8:300-304.
- 45 Noura S, Yamamoto H, Miyake Y, et al. Immunohistochemical assessment of localization of frequency of micrometastasis in lymph nodes of colorectal cancer. *Clin Cancer Research* 2002;8:759-767.
- 46 Oberg A, Stenling R, Tavelin B, Lindmark G. Are lymph node micrometastasis of any clinical significance in Duke stages A and B colorectal cancer? *Dis Colon Rectum* 1998;41:1244-1249.
- 47 Greenson JK, Isenhardt TCE, Rice R, et al. Identification of occult micrometastasis in pericolic lymph nodes of Duke's B colorectal cancer. Patient's using monoclonal antibodies against cytokeratin and CC49. Correlation with long term survival. *Cancer* 1994;73:563-9.
- 48 Lievre A, Bachatte J-B, Blige V, et al. KRAS mutations as an independent prognostic factor in patients with advanced colorectal cancer treated with Cetuximab. *J Clin Oncol* 2008;26:374-379.
- 49 Amado IG, Wolf M, Peters M, et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* 2008;26:1626-1634.
- 50 Douillard JY, Oliner KS, Siena S, et al. Panitumumab--FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* 2013;369:1023-1034.
- 51 Di Nicolantonio F, Martini M, Molinari F, et al. Wild-type BRAF is required for response to panitumumab or cetuximab in metastatic colorectal cancer. *J Clin Oncol* 2008;26:5705-5712.
- 52 Bokmeyer C, Cutsem EV, Rougier P, et al. Addition of cetuximab to chemotherapy as first-line treatment for KRAS wild-type metastatic colorectal cancer: Pooled analysis of the CRYSTAL and OPUS randomised clinical trials. *Eur J Cancer* 2012;48:1466-1475.
- 53 Pietrantonio F, Petrelli F, Coiro A, et al. Predictive role of BRAF mutations in patients with advanced colorectal cancer receiving cetuximab and panitumumab: a meta-analysis. *Eur J Cancer* 2015
- 54 Etienne-Gimeldi M-C, Formenta J-L, Francoual M, et al. KRAS mutations in treatment outcome in colorectal cancer in patients receiving exclusive fluoropyrimidine. *Clin Cancer Research* 2008;14:4830-4835.
- 55 Parfitt JR and Driman KR. Total mesorectal excision specimen for rectal cancer: A review of its pathological assessment. *J Clin Pathol* 60:849-855, 2007.
- 56 Jass JR, O'Brien MJ, Riddell RH, Snover DC. On behalf of the association of Directors of Anatomic and Surgical Pathology recommendations for the reporting of surgically resected specimens in colorectal carcinoma. *Human Pathol* 38:537-545, 2007.
- 57 Nagtegaal ID, Vandevelde CJ, Derworp EV, et al. Macroscopic evaluation of the rectal cancer resection margin: Clinical significance of the pathologist in quality control. *J Clin Oncol* 20: 1729-1734, 2002.
- 58 Sargent DJ, Marsoni S, Monges G, et al. Defective mismatch repair as a predictive marker for lack of efficacy of fluorouracil-based adjuvant therapy in colon cancer. *J Clin Oncol* 2010;28:3219-3226.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF SURGERY (1 of 3)****Transanal Excision:¹****• Criteria**

- ▶ <30% circumference of bowel
- ▶ <3 cm in size
- ▶ Margin clear (>3 mm)
- ▶ Mobile, nonfixed
- ▶ Within 8 cm of anal verge
- ▶ T1 only
- ▶ Endoscopically removed polyp with cancer or indeterminate pathology
- ▶ No lymphovascular invasion or PNI
- ▶ Well to moderately differentiated
- ▶ No evidence of lymphadenopathy on pretreatment imaging

• When the lesion can be adequately identified in the rectum, transanal endoscopic microsurgery (TEM) may be used. TEM for more proximal lesions may be technically feasible.

Transabdominal Resection: Abdominoperineal resection or low anterior resection or coloanal anastomosis using total mesorectal excision

• Management principles

- ▶ The treating surgeon should perform a rigid proctoscopy before initiating treatment.
- ▶ Remove primary tumor with adequate margins.
- ▶ Treat draining lymphatics by total mesorectal excision.
- ▶ Restore organ integrity, if possible.
- ▶ Surgery should be 5–12 weeks following full-dose 5.5-week neoadjuvant chemoradiation.

• Total mesorectal excision

- ▶ Reduces positive radial margin rate.
- ▶ Extend 4–5 cm below distal edge of tumors for an adequate mesorectal excision. In distal rectal cancers (ie, <5 cm from anal verge), negative distal bowel wall margin of 1–2 cm may be acceptable; this must be confirmed to be tumor free by frozen section.
- ▶ Full rectal mobilization allows for a negative distal margin and adequate mesorectal excision.

- Some studies have shown that laparoscopy is associated with similar short- and long-term outcomes when compared to open surgery,^{2,3} whereas other studies have shown that laparoscopy is associated with higher rates of circumferential margin positivity and incomplete TME.^{4,5} Therefore, minimally invasive resection may be considered based on the following principles:
 - ▶ The surgeon should have experience performing minimally invasive proctectomy with total mesorectal excision.
 - ▶ It is not indicated for locally advanced disease with a threatened or high-risk circumferential margin based on staging. For these high-risk tumors, open surgery is preferred.
 - ▶ It is not indicated for acute bowel obstruction or perforation from cancer.
 - ▶ Thorough abdominal exploration is required.
- Lymph node dissection^{6,7}
 - ▶ Biopsy or remove clinically suspicious nodes beyond the field of resection if possible.
 - ▶ Extended resection is not indicated in the absence of clinically suspected nodes.

[See Criteria for Resectability of Metastases on REC-B 2 of 3](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF SURGERY (2 of 3)****CRITERIA FOR RESECTABILITY OF METASTASES AND LOCOREGIONAL THERAPIES WITHIN SURGERY****Liver**

- Hepatic resection is the treatment of choice for resectable liver metastases from colorectal cancer.⁸
- Complete resection must be feasible based on anatomic grounds and the extent of disease; maintenance of adequate hepatic function is required.^{9,10}
- The primary tumor must have been resected for cure (R0). There should be no unresectable extrahepatic sites of disease.¹¹⁻¹³ Plan for a debulking resection (R1/R2 resection) is not recommended.
- Patients with resectable metastatic disease and primary tumor in place should have both sites resected with curative intent. These can be resected in one operation or as a staged approach, depending on the complexity of the hepatectomy or colectomy, comorbid diseases, surgical exposure, and surgeon expertise.
- When hepatic metastatic disease is not optimally resectable based on insufficient remnant liver volume, approaches utilizing preoperative portal vein embolization or staged liver resections can be considered.
- Ablative techniques may be considered alone or in conjunction with resection.⁸ All original sites of disease need to be amenable to ablation or resection.
- Arterially directed catheter therapy, and in particular yttrium 90 microsphere selective internal radiation, is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases.
- Conformal external beam radiation therapy (category 3) may be considered in highly selected cases or in the setting of a clinical trial and should not be used indiscriminately in patients who are potentially surgically resectable.
- Re-resection can be considered in selected patients.¹⁴

Lung

- Complete resection based on the anatomic location and extent of disease with maintenance of adequate function is required.¹⁵⁻¹⁸
- The primary tumor must have been resected for cure (R0).
- Resectable extrapulmonary metastases do not preclude resection.¹⁹⁻²²
- Re-resection can be considered in selected patients.²³
- Ablative techniques may be considered alone or in conjunction with resection. All original sites of disease need to be amenable to ablation or resection.
- Ablative techniques can be considered when unresectable and amenable to complete ablation.
- Patients with resectable synchronous metastases can be resected synchronously or using a staged approach.
- Conformal external beam radiation therapy may be considered in highly selected cases or in the setting of a clinical trial and should not be used indiscriminately in patients who are potentially surgically resectable (category 3).

Evaluation for Conversion to Resectable Disease

- Re-evaluation for resection should be considered in otherwise unresectable patients after 2 months of preoperative chemotherapy and every 2 months thereafter.²⁴⁻²⁷
- Disease with a higher likelihood of being converted to resectable are those with initially convertible disease distributed within limited sites.
- When considering whether disease has been converted to resectable, all original sites need to be amenable to resection.²⁸ Preoperative chemotherapy regimens with high response rates should be considered for patients with potentially convertible disease.²⁹

[See footnotes on REC-B 3 of 3](#)**Note:** All recommendations are category 2A unless otherwise indicated.**Clinical Trials:** NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF SURGERY (3 of 3) - REFERENCES**

- ¹Nash GM, Weiser MR, Guillem JG, et al. Long-term survival after transanal excision of T1 rectal cancer. *Dis Colon Rectum* 2009;52:577-82.
- ²Bonjer HJ, Deijen CL, Abis GA, et al. A randomized trial of laparoscopic versus open surgery for rectal cancer. *N Eng J Med* 2015;372:1324-1332.
- ³Jeong S-Y, Park JW, Nam BH, et al. Open versus laparoscopic surgery for mid-rectal or low-rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): survival outcomes of an open-label, non-inferiority, randomised controlled trial. *Lancet Oncol* 2014;15:767-774.
- ⁴Fleshman J, Branda M, Sargent DJ, et al. Effect of laparoscopic-assisted resection vs open resection of stage II or III rectal cancer on pathological outcomes. The ACOSOG Z6051 randomized clinical trial. *JAMA* 2015;314:1346-1355.
- ⁵Stevenson ARL, Solomon MJ, Lumley JW, et al. Effect of laparoscopic-assisted resection vs open resection on pathological outcomes in rectal cancer. The ALacaRT randomized clinical trial. *JAMA* 2015;314:1356-1363.
- ⁶Gunderson LL, Sargent DJ, Tepper JB, et al. Impact of T and N stage and treatment on survival and relapse in adjuvant rectal cancer: a pooled analysis. *J Clin Oncol* 2004;22(10):1785-1796.
- ⁷Greene FL, Stewart AK, Norton HJ. New tumor-node-metastasis staging strategy for node-positive (stage III) rectal cancer: an analysis. *J Clin Oncol* 2004;22(10):1778-1784.
- ⁸Abdalla EK, Vauthey JN, Ellis LM, et al. Recurrence and outcomes following hepatic resection, radiofrequency ablation, and combined resection/ablation for colorectal liver metastases. *Ann Surg* 2004;239:818-825; discussion 825-7.
- ⁹Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of indications for resection. *Registry of Hepatic Metastases. Surgery* 1988;103:278-288.
- ¹⁰Hughes KS, Simon R, Songhorabodi S, et al. Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of patterns of recurrence. *Surgery* 1986;100:278-284.
- ¹¹Fong Y, Cohen AM, Fortner JG, et al. Liver resection for colorectal metastases. *J Clin Oncol* 1997;15:938-946.
- ¹²Nordlinger B, Quilichini MA, Parc R, Hannoun L, Delva E, Huguet C. Surgical resection of liver metastases from colo-rectal cancers. *Int Surg* 1987;72:70-72.
- ¹³Fong Y, Fortner J, Sun RL, Brennan MF, Blumgart LH. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: analysis of 1001 consecutive cases. *Ann Surg* 1999;230:309-318; discussion 318-321.
- ¹⁴Adam R, Bismuth H, Castaing D, et al. Repeat hepatectomy for colorectal liver metastases. *Ann Surg* 1997;225:51-62.
- ¹⁵McAfee MK, Allen MS, Trastek VF, Ilstrup DM, Deschamps C, Pairolero PC. Colorectal lung metastases: results of surgical excision. *Ann Thorac Surg* 1992;53:780-785; discussion 785-786.
- ¹⁶Regnard JF, Grunenwald D, Spaggiari L, et al. Surgical treatment of hepatic and pulmonary metastases from colorectal cancers. *Ann Thorac Surg* 1998;66:214-218; discussion 218-219.
- ¹⁷Inoue M, Kotake Y, Nakagawa K, Fujiwara K, Fukuhara K, Yasumitsu T. Surgery for pulmonary metastases from colorectal carcinoma. *Ann Thorac Surg* 2000;70:380-383.
- ¹⁸Sakamoto T, Tsubota N, Iwanaga K, Yuki T, Matsuoka H, Yoshimura M. Pulmonary resection for metastases from colorectal cancer. *Chest* 2001;119:1069-1072.
- ¹⁹Rena O, Casadio C, Viano F, et al. Pulmonary resection for metastases from colorectal cancer: factors influencing prognosis. Twenty-year experience. *Eur J Cardiothorac Surg* 2002;21:906-912.
- ²⁰Irshad K, Ahmad F, Morin JE, Mulder DS. Pulmonary metastases from colorectal cancer: 25 years of experience. *Can J Surg* 2001;44:217-221.
- ²¹Ambiru S, Miyazaki M, Ito H, et al. Resection of hepatic and pulmonary metastases in patients with colorectal carcinoma. *Cancer* 1998;82:274-278.
- ²²Yano T, Hara N, Ichinose Y, Yokoyama H, Miura T, Ohta M. Results of pulmonary resection of metastatic colorectal cancer and its application. *J Thorac Cardiovasc Surg* 1993;106:875-879.
- ²³Hendriks JM, Romijn S, Van Putte B, et al. Long-term results of surgical resection of lung metastases. *Acta Chir Belg* 2001;101:267-272.
- ²⁴Adam R, Avisar E, Ariche A, et al. Five-year survival following hepatic resection after neoadjuvant therapy for nonresectable colorectal. *Ann Surg Oncol* 2001;8:347-353.
- ²⁵Rivoire M, De Cian F, Meeus P, Negrier S, Sebban H, Kaemmerlen P. Combination of neoadjuvant chemotherapy with cryotherapy and surgical resection for the treatment of unresectable liver metastases from colorectal carcinoma. *Cancer* 2002;95:2283-2292.
- ²⁶Vauthey JN, Pawlik TM, Ribero D, et al. Chemotherapy regimen predicts steatohepatitis and an increase in 90-day mortality after surgery for hepatic colorectal metastases. *J Clin Oncol*. 2006 May 1;24(13):2065-72.
- ²⁷Pawlik TM, Olino K, Gleisner AL, et al. Preoperative chemotherapy for colorectal liver metastases: impact on hepatic histology and postoperative outcome. *J Gastrointest Surg*. 2007 Jul;11(7):860-8.
- ²⁸Benoist S, Brouquet A, Penna C, et al. Complete response of colorectal liver metastases after chemotherapy: does it mean cure? *J Clin Oncol*. 2006 Aug 20;24(24):3939-45.
- ²⁹Bartlett DL, Berlin J, Lauwers GY, et al. Chemotherapy and regional therapy of hepatic colorectal metastases: expert consensus statement. *Ann Surg Oncol*. 2006;13:1284-92.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF ADJUVANT THERAPY (1 of 2)**

Adjuvant therapy for rectal cancer consists of regimens that include both concurrent chemotherapy/RT and adjuvant chemotherapy. A total of approximately 6 months of perioperative treatment is preferred.

Postoperative Adjuvant Chemotherapy:• **mFOLFOX 6^{1,2,3}**

Oxaliplatin 85 mg/m² IV, day 1,* leucovorin 400 mg/m² IV day 1,** 5-FU 400 mg/m² IV bolus on day 1, then 1200 mg/m²/d x 2 days (total 2400 mg/m² over 46–48 hours) continuous infusion. Repeat every 2 weeks to a total of 6 mo perioperative therapy.

• **Simplified biweekly infusional 5-FU/LV (sLV5FU2)⁴**

Leucovorin 400 mg/m² IV day 1, followed by 5-FU bolus 400 mg/m² and then 1200 mg/m²/d x 2 days (total 2400 mg/m² over 46–48 hours) continuous infusion. Repeat every 2 weeks to a total of 6 mo perioperative therapy.

• **Capecitabine⁵**

Capecitabine[#] 1000–1250 mg/m² twice daily days 1–14 every 3 weeks to a total of 6 months perioperative therapy.

• **CAPEOX^{6,7}**

Oxaliplatin 130 mg/m² day 1.* Capecitabine[#] 1000 mg/m² twice daily days 1–14 every 3 weeks. Repeat every 3 weeks to a total of 6 months perioperative therapy.

• **5-FU 500 mg/m² IV bolus weekly x 6 + leucovorin 500 mg/m² IV weekly x 6, each 8-week cycle. Repeat every 8 weeks to a total of 6 months perioperative therapy.⁸****Dosing Schedules for Concurrent Chemotherapy/RT:**• **XRT + continuous infusion 5-FU⁹**

5-FU 225 mg/m² over 24 hours 5 or 7 days/week during XRT

• **XRT + Capecitabine^{11,12}**

Capecitabine[#] 825 mg/m² twice daily 5 d/wk + XRT x 5 weeks

• **XRT + 5-FU/leucovorin^{10‡}**

5-FU 400 mg/m² IV bolus + leucovorin 20 mg/m² IV bolus for 4 days during week 1 and 5 of XRT

[See footnotes on REC-C 2 of 2](#)

*Oxaliplatin may be given either over 2 hours, or may be infused over a shorter time at a rate of 1 mg/m²/min. Leucovorin infusion should match infusion time of oxaliplatin.

Cercek A, Park V, Yaeger R, et al. Faster FOLFOX: oxaliplatin can be safely infused at a rate of 1 mg/m²/min. J Oncol Pract 2016;12:e548-553.

**Leucovorin 400 mg/m² is the equivalent of levoleucovorin 200 mg/m².

‡Bolus 5-FU/leucovorin/RT is an option for patients not able to tolerate capecitabine or infusional 5-FU.

#The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF ADJUVANT THERAPY (2 of 2) - REFERENCES

- ¹Andre T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 2004;350:2343-2351.
- ²Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Br J Cancer* 2002;87:393-399.
- ³Maindault-Goebel F, deGramont A, Louvet C, et al. Evaluation of oxaliplatin dose intensity in bimonthly leucovorin and 48-hour 5-fluorouracil continuous infusion regimens (FOLFOX) in pretreated metastatic colorectal cancer. *Ann Oncol* 2000;11:1477-1483.
- ⁴Andre T, Louvet C, Maindault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
- ⁵Twelves C, Wong A, Nowacki MP, et al. Capecitabine as adjuvant treatment for stage III colon cancer. *N Engl J Med* 2005;352(26):2696-2704.
- ⁶Schmoll HJ, Cartwright T, Tabernero J, et al. Phase III trial of capecitabine plus oxaliplatin as adjuvant therapy for stage III colon cancer: a planned safety analysis in 1,864 patients. *J Clin Oncol* 2007;25:102-109.
- ⁷Haller DG, Tabernero J, Maroun J, et al. Capecitabine Plus Oxaliplatin Compared With Fluorouracil and Folinic Acid As Adjuvant Therapy for Stage III Colon Cancer. *J Clin Oncol* 2011;29:1465-1471.
- ⁸Petrelli N, Douglass Jr HO, Herrare L, et al. The modulation of fluorouracil with leucovorin in metastatic colorectal carcinoma: a prospective randomized phase III trial. *J Clin Oncol* 1989;7:1419-1426.
- ⁸O'Connell MJ, Martenson JA, Wieand HS, et al. Improving adjuvant therapy for rectal cancer by combining protracted-infusion fluorouracil with radiation therapy after curative surgery. *N Engl J Med* 1994; 331:502-507.
- ¹⁰Tepper JE, O'Connell M, Niedzwiecki D, et al. Adjuvant therapy in rectal cancer: analysis of stage, sex, and local control--final report of Intergroup 0114. *J Clin Oncol* 2002;20:1744-1750.
- ¹¹O'Connell MJ, Colangelo LH, Beart RW, et al. Capecitabine and oxaliplatin in the preoperative multimodality treatment of rectal cancer: surgical end points from National Surgical Adjuvant Breast and Bowel Project trial R-04. *J Clin Oncol* 2014;32:1927-1934.
- ¹²Hofheinz R, Wenz FK, Post S, et al. Chemoradiotherapy with capecitabine versus fluorouracil for locally advanced rectal cancer: A randomized, multicentre, non-inferiority, phase 3 trial. *Lancet Oncol* 2012;13:579-588.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**PRINCIPLES OF RADIATION THERAPY**

- Radiation therapy fields should include the tumor or tumor bed, with a 2–5 cm margin, the presacral nodes, and the internal iliac nodes. The external iliac nodes should also be included for T4 tumors involving anterior structures.
- Multiple radiation therapy fields should be used (generally a 3- or 4-field technique). Positioning and other techniques to minimize the volume of small bowel in the fields should be encouraged.
- For postoperative patients treated by abdominoperineal resection, the perineal wound should be included within the fields.
- Intensity-modulated radiation therapy (IMRT) should only be used in the setting of a clinical trial or in unique clinical situations such as reirradiation of previously treated patients with recurrent disease or unique anatomical situations.
- Radiation doses:
 - ▶ 45–50 Gy in 25–28 fractions to the pelvis.
 - ▶ For resectable cancers, after 45 Gy a tumor bed boost with a 2-cm margin of 5.4 Gy in 3 fractions could be considered for preoperative radiation and 5.4–9.0 Gy in 3–5 fractions for postoperative radiation.
 - ▶ Small bowel dose should be limited to 45 Gy.
- Short-course radiation therapy (25 Gy in 5 fractions) with surgery within 1 to 2 weeks of completion of therapy can also be considered for patients with ultrasound or pelvic MRI stage T3 rectal cancer.¹
- Intraoperative radiation therapy (IORT), if available, may be considered for very close or positive margins after resection, as an additional boost, especially for patients with T4 or recurrent cancers. If IORT is not available, 10–20 Gy external beam radiation and/or brachytherapy to a limited volume could be considered soon after surgery, prior to adjuvant chemotherapy.
- For unresectable cancers, doses higher than 54 Gy may be required, if technically feasible.
- Fluoropyrimidine–based chemotherapy should be delivered concurrently with radiation therapy.
- Arterially directed catheter therapy, and in particular yttrium 90 microsphere selective internal radiation, is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases.
- In patients with a limited number of liver or lung metastases, radiotherapy to the metastatic site can be considered in highly selected cases or in the setting of a clinical trial. Radiotherapy should not be used in the place of surgical resection. Radiotherapy should be delivered in a highly conformal manner. The techniques can include 3-D conformal radiation therapy, IMRT, or stereotactic body radiation therapy (SBRT).
- Side effect management:
 - Female patients should be considered for vaginal dilators and instructed on the symptoms of vaginal stenosis.
 - Male patients should be counseled on infertility risks and given information regarding sperm banking.
 - Female patients should be counseled on infertility risks and given information regarding oocyte, egg, or ovarian tissue banking prior to treatment.

¹Ngan SY, Burmeister B, Fisher RJ, et al. Randomized trial of short-course radiotherapy versus long-course chemoradiation comparing rates of local recurrence in patients with T3 rectal cancer: Trans-Tasman Radiation Oncology Group trial 01.04. J Clin Oncol 2012;30:3827-3833.

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017

Rectal Cancer

CONTINUUM OF CARE - SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 1 of 10)



[See footnotes on REC-E 6 of 10](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

CONTINUUM OF CARE - SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 2 of 10)

Subsequent Therapy



*if neither previously given
**if not previously given

[See footnotes on REC-E 6 of 10](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

CONTINUUM OF CARE - SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 3 of 10)

Subsequent Therapy



*if neither previously given

**if not previously given

[See footnotes on REC-E 6 of 10](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

CONTINUUM OF CARE - SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 4 of 10)

Subsequent Therapy



[See footnotes on REC-E 6 of 10](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 3.2017

Rectal Cancer

CONTINUUM OF CARE - SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE:¹ (PAGE 5 of 10)

Subsequent Therapy



*if neither previously given
**if not previously given

[See footnotes on REC-E 6 of 10](#)

Note: All recommendations are category 2A unless otherwise indicated.
Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE (PAGE 6 of 10)

¹For chemotherapy references, [see Chemotherapy Regimens and References \(REC-E 7-10\)](#).

²Chest/Abdominal/Pelvic CT with contrast or Chest CT and Abdominal/Pelvic MRI with contrast to monitor progress of therapy. PET/CT should not be used.

³[See Principles of Pathologic Review \(REC-A 5 of 6\)](#).

⁴BRAF V600E mutation makes response to panitumumab or cetuximab highly unlikely.

⁵Irinotecan should be used with caution in patients with Gilbert's disease or elevated serum bilirubin. There is a commercially available test for UGT1A1. Guidelines for use in clinical practice have not been established.

⁶A treatment option for patients not able to tolerate oxaliplatin or irinotecan.

⁷The use of single-agent capecitabine after progression on a fluoropyrimidine-containing regimen has been shown to be ineffective; therefore, this is not recommended.

⁸Bevacizumab is the preferred anti-angiogenic agent based on toxicity and/or cost.

⁹There are no data to suggest activity of FOLFIRI-ziv-aflibercept or FOLFIRI-ramucirumab in a patient who has progressed on FOLFIRI-bevacizumab, or vice versa. Ziv-aflibercept and ramucirumab have only shown activity when given in conjunction with FOLFIRI in FOLFIRI-naïve patients.

¹⁰Cetuximab or panitumumab are recommended in combination with irinotecan-based therapy or as single-agent therapy for patients who cannot tolerate irinotecan.

¹¹Regorafenib or trifluridine + tipiracil are treatment options for patients who have progressed through all available regimens.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE - CHEMOTHERAPY REGIMENS (PAGE 7 of 10)****mFOLFOX 6^{1,2,3†}****Oxaliplatin 85 mg/m² IV day 1*****Leucovorin 400 mg/m² IV day 1******5-FU 400 mg/m² IV bolus on day 1, then 1200 mg/m²/d x 2 days (total 2400 mg/m² over 46–48 hours) IV continuous infusion****Repeat every 2 weeks****mFOLFOX7⁴****Oxaliplatin 85 mg/m² IV day 1*****Leucovorin 400 mg/m² IV day 1******5-FU 1200 mg/m²/d x 2 days (total 2400 mg/m² over 46–48 hours) IV continuous infusion****Repeat every 2 weeks****FOLFOX + bevacizumab⁵****Bevacizumab 5 mg/kg IV, day 1****Repeat every 2 weeks****FOLFOX + panitumumab⁶ (KRAS/NRAS WT only)****Panitumumab 6 mg/kg IV over 60 minutes, day 1****Repeat every 2 weeks****FOLFOX + cetuximab⁷ (KRAS/NRAS WT only)****Cetuximab 400 mg/m² IV over 2 hours first infusion, then 250 mg/m² IV over 60 minutes weekly****or Cetuximab 500 mg/m² IV over 2 hours, day 1, every 2 weeks****CAPEOX⁸****Oxaliplatin 130 mg/m² IV day 1*****Capecitabine 1000[‡] mg/m² twice daily PO for 14 days****Repeat every 3 weeks****CAPEOX + bevacizumab^{8†}****Oxaliplatin 130 mg/m² IV day 1*****Capecitabine 1000[‡] mg/m² PO twice daily for 14 days****Bevacizumab 7.5 mg/kg IV day 1****Repeat every 3 weeks**[See References on REC-E 10 of 10](#)*Oxaliplatin may be given either over 2 hours, or may be infused over a shorter time at a rate of 1 mg/m²/min. Leucovorin infusion should match infusion time of oxaliplatin.Cercek A, Park V, Yaeger R, et al. Faster FOLFOX: oxaliplatin can be safely infused at a rate of 1 mg/m²/min. J Oncol Pract 2016;12:e548-553.**Leucovorin 400 mg/m² is the equivalent of levoleucovorin 200 mg/m².‡The majority of safety and efficacy data for this regimen have been developed in Europe, where a capecitabine starting dose of 1000 mg/m² twice daily for 14 days, repeated every 21 days, is standard. Evidence suggests that North American patients may experience greater toxicity with capecitabine (as well as with other fluoropyrimidines) than European patients, and may require a lower dose of capecitabine.

†Bevacizumab may be safely given at a rate of 0.5 mg/kg/min (5 mg/kg over 10 minutes and 7.5 mg/kg over 15 minutes).

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE - CHEMOTHERAPY REGIMENS (PAGE 8 of 10)****FOLFIRI^{9,10}**

Irinotecan 180 mg/m² IV over 30–90 minutes, day 1
 Leucovorin** 400 mg/m² IV infusion to match duration of irinotecan infusion, day 1
 5-FU 400 mg/m² IV bolus day 1, then 1200 mg/m²/d x 2 days (total 2400 mg/m² over 46–48 hours) continuous infusion
 Repeat every 2 weeks

FOLFIRI + bevacizumab^{11,††}

Bevacizumab 5 mg/kg IV, day 1
 Repeat every 2 weeks

FOLFIRI + cetuximab (KRAS/NRAS WT only)

Cetuximab 400 mg/m² IV over 2 hours first infusion, then 250 mg/m² IV over 60 minutes weekly¹²
 or Cetuximab 500 mg/m² IV over 2 hours, day 1, every 2 weeks¹³

FOLFIRI + panitumumab¹⁴ (KRAS/NRAS WT only)

Panitumumab 6 mg/kg IV over 60 minutes, day 1
 Repeat every 2 weeks

FOLFIRI + ziv-aflibercept¹⁵

Ziv-aflibercept 4 mg/kg IV over 60 minutes, day 1
 Repeat every 2 weeks

FOLFIRI + ramucirumab¹⁶

Ramucirumab 8 mg/kg over 60 minutes, day 1
 Repeat every 2 weeks

FOLFOXIRI¹⁷

Irinotecan 165 mg/m² IV day 1, oxaliplatin 85 mg/m² IV day 1,*
 leucovorin 400** mg/m² day 1, fluorouracil 1600 mg/m²/d x 2 days (total 3200 mg/m² over 48 hours) continuous infusion starting on day 1.
 Repeat every 2 weeks
 The dose of 5-FU listed here was used in European studies. U.S. patients have been shown to have poorer tolerance for 5-FU. A starting dose of 5-FU consistent with the dose recommended in FOLFOX or FOLFIRI should be strongly considered for U.S. patients.

FOLFOXIRI + bevacizumab¹⁸

Bevacizumab 5 mg/kg IV, day 1
 Repeat every 2 weeks

IROX¹⁹

Oxaliplatin 85 mg/m² IV*,
 followed by irinotecan 200 mg/m² over 30–90 minutes every 3 weeks

[See References on REC-E 10 of 10](#)

*Oxaliplatin may be given either over 2 hours, or may be infused over a shorter time at a rate of 1 mg/m²/min. Leucovorin infusion should match infusion time of oxaliplatin.

Cercek A, Park V, Yaeger R, et al. Faster FOLFOX: oxaliplatin can be safely infused at a rate of 1 mg/m²/min. J Oncol Pract 2016;12:e548-553.

**Leucovorin 400 mg/m² is the equivalent of levoleucovorin 200 mg/m².

††Bevacizumab may be safely given at a rate of 0.5 mg/kg/min (5 mg/kg over 10 minutes and 7.5 mg/kg over 15 minutes).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE - CHEMOTHERAPY REGIMENS (PAGE 9 of 10)****Bolus or infusional 5-FU/leucovorin****Roswell Park regimen²⁰**

Leucovorin 500 mg/m² IV over 2 hours, days 1, 8, 15, 22, 29, and 36
5-FU 500 mg/m² IV bolus 1 hour after start of leucovorin,
days 1, 8, 15, 22, 29, and 36
Repeat every 8 weeks

Simplified biweekly infusional 5-FU/LV (sLV5FU2)⁹

Leucovorin** 400 mg/m² IV over 2 hours on day 1,
followed by 5-FU bolus 400 mg/m² and then 1200 mg/m²/d x 2 days
(total 2400 mg/m² over 46–48 hours) continuous infusion
Repeat every 2 weeks

Weekly

Leucovorin 20 mg/m² IV over 2 hours on day 1, 5-FU 500 mg/m² IV
bolus injection 1 hour after the start of leucovorin. Repeat weekly.²¹
5-FU 2600 mg/m² by 24-hour infusion plus leucovorin 500 mg/m²
Repeat every week²¹

Capecitabine⁸

Capecitabine 850–1250 mg/m² PO twice daily, days 1–14
Repeat every 3 weeks

Capecitabine + Bevacizumab^{22,†}

Bevacizumab 7.5 mg/kg IV, day 1
Repeat every 3 weeks

Irinotecan

Irinotecan 125 mg/m² IV over 30–90 minutes, days 1 and 8
Repeat every 3 weeks^{23,24}
or Irinotecan 180 mg/m² IV over 30–90 minutes, day 1

Repeat every 2 weeks

or Irinotecan 300–350 mg/m² IV over 30–90 minutes, day 1

Repeat every 3 weeks**Irinotecan + cetuximab (KRAS/NRAS WT only)**

Cetuximab 400 mg/m² first infusion, then 250 mg/m² IV weekly²⁵
or Cetuximab 500 mg/m² IV over 2 hours, day 1, every 2 weeks¹³

Cetuximab (KRAS/NRAS WT only)

Cetuximab 400 mg/m² first infusion, then 250 mg/m² IV weekly²⁵
or Cetuximab 500 mg/m² IV over 2 hours, day 1, every 2 weeks¹³

Panitumumab²⁶ (KRAS/NRAS WT only)

Panitumumab 6 mg/kg IV over 60 minutes every 2 weeks

Regorafenib²⁷

Regorafenib 160 mg[§] PO daily days 1–21
Repeat every 28 days

Trifluridine + tipiracil²⁸

Trifluridine + tipiracil 35 mg/m² up to a maximum dose of 80 mg per
dose (based on the trifluridine component)
PO twice daily days 1–5 and 8–12
Repeat every 28 days

Pembrolizumab²⁹

Pembrolizumab 2 mg/kg every 3 weeks

Nivolumab³⁰

Nivolumab 3 mg/kg every 2 weeks
or Nivolumab 240 mg IV every two weeks

[See References on REC-E 10 of 10](#)

**Leucovorin 400 mg/m² is the equivalent of levoleucovorin 200 mg/m².

†Bevacizumab may be safely given at a rate of 0.5 mg/kg/min (5 mg/kg over 10 minutes and 7.5 mg/kg over 15 minutes).

§It is common practice to start at a lower dose of regorafenib (80 or 120 mg) and escalate, as tolerated.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

**SYSTEMIC THERAPY FOR ADVANCED OR METASTATIC DISEASE - REFERENCES (PAGE 10 of 10)**

- ¹deGramont A, Figer A, Seymour M, et al. Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced rectal cancer. *J Clin Oncol* 2000;18:2938-2947.
- ²Cheeseman SL, Joel SP, Chester JD, et al. A 'modified de Gramont' regimen of fluorouracil, alone and with oxaliplatin, for advanced colorectal cancer. *Br J Cancer* 2002;87:393-399.
- ³Maindrault-Goebel F, deGramont A, Louvet C, et al. Evaluation of oxaliplatin dose intensity in bimonthly leucovorin and 48-hour 5-fluorouracil continuous infusion regimens (FOLFOX) in pretreated metastatic colorectal cancer. *Ann Oncol* 2000;11:1477-1483.
- ⁴Hochster HS, Grothey A, Hart L, et al. Improved time to treatment failure with an intermittent oxaliplatin strategy: results of CONcePT. *Ann Oncol* 2014;25:1172-1178.
- ⁵Emmanouilides C, Sfakiotaki G, Androulakis N, et al. Front-line bevacizumab in combination with oxaliplatin, leucovorin and 5-fluorouracil (FOLFOX) in patients with metastatic colorectal cancer: a multicenter phase II study. *BMC Cancer* 2007;7:91.
- ⁶Douillard JY, Siena S, Cassidy J, et al. Randomized, phase III trial of panitumumab with infusional fluorouracil, leucovorin, and oxaliplatin (FOLFOX4) versus FOLFOX4 alone as first-line treatment in patients with previously untreated metastatic colorectal cancer: the PRIME study. *J Clin Oncol* 2010;28:4697-4705.
- ⁷Venook AP, Niedzwiecki D, Lenz H-J, et al. CALGB/SWOG 80405: Phase III trial of irinotecan/5-FU/leucovorin (FOLFIRI) or oxaliplatin/5-FU/leucovorin (mFOLFOX6) with bevacizumab or cetuximab for patients with KRAS wild-type untreated metastatic adenocarcinoma of the colon or rectum [abstract]. *ASCO Meeting Abstracts* 2014;32:LBA3.
- ⁸Saltz LB, Clarke S, Diaz-Rubio E, et al. Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in metastatic colorectal cancer: a randomized phase III study. *J Clin Oncol* 2008;26:2013-2019.
- ⁹Andre T, Louvet C, Maindrault-Goebel F, et al. CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer. *Eur J Cancer* 1999;35(9):1343-7.
- ¹⁰Fuchs CS, Marshall J, Mitchell E, et al. Randomized, controlled trial of irinotecan plus infusional, bolus, or oral fluoropyrimidines in first-line treatment of metastatic colorectal cancer: results from the BICC-C Study. *J Clin Oncol* 2007;25:4779-4786.
- ¹¹Heinemann V, von Weikersthal LF, Decker T, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomized, open-label, phase 3 trial. *Lancet Oncol* 2014.
- ¹²Cunningham D, Humblet Y, Siena S, et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* 2004;351:337-345.
- ¹³Martín-Martorell P, Roselló S, Rodríguez-Braun E, et al. Biweekly cetuximab and irinotecan in advanced colorectal cancer patients progressing after at least one previous line of chemotherapy: results of a phase II single institution trial. *Br J Cancer* 2008;99:455-458.
- ¹⁴Peeters M, Price TJ, Cervantes A, et al. Randomized phase III study of panitumumab with fluorouracil, leucovorin, and irinotecan (FOLFIRI) compared with FOLFIRI alone as second-line treatment in patients with metastatic colorectal cancer. *J Clin Oncol* 2010;28:4706-4713.
- ¹⁵Van Cutsem E, Tabernero J, Lakomy R, et al. Addition of Afibercept to Fluorouracil, Leucovorin, and Irinotecan Improves Survival in a Phase III Randomized Trial in Patients With Metastatic Colorectal Cancer Previously Treated With an Oxaliplatin-Based Regimen. *J Clin Oncol* 2012;30:3499-3506.
- ¹⁶Tabernero J, Yoshino T, Cohn AL, et al. Ramucirumab versus placebo in combination with second-line FOLFIRI in patients with metastatic colorectal carcinoma that progressed during or after first-line therapy with bevacizumab, oxaliplatin, and a fluoropyrimidine (RAISE): a randomized, double-blind, multicentre, phase 3 study. *Lancet Oncol* 2015;16:499-508.
- ¹⁷Falcone A, Ricci S, Brunetti I, et al. Phase III trial of infusional fluorouracil, leucovorin, oxaliplatin, and irinotecan (FOLFOXIRI) compared with infusional fluorouracil, leucovorin, and irinotecan (FOLFIRI) as first-line treatment for metastatic colorectal cancer: The Gruppo Oncologico Nord Ovest. *J Clin Oncol* 2007;25(13):1670-1676.
- ¹⁸Cremolini C, Loupakis F, Antoniotti C, et al. FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. *Lancet Oncol* 2015;16:1306-1315.
- ¹⁹Haller DG, Rothenberg ML, Wong AO, et al. Oxaliplatin plus irinotecan compared with irinotecan alone as second-line treatment after single agent fluoropyrimidine therapy for metastatic colorectal carcinoma. *J Clin Oncol* 2008;26:4544-4550.
- ²⁰Wolmark N, Rockette H, Fisher B, et al. The benefit of leucovorin-modulated fluorouracil as postoperative adjuvant therapy for primary colon cancer: results from National Surgical Adjuvant Breast and Bowel Protocol C-03. *J Clin Oncol* 1993;11:1879-1887.
- ²¹Jäger E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. *J Clin Oncol* 1996;14:2274-2279.
- ²²Cunningham D, Lang I, Marcuello E, et al. Bevacizumab plus capecitabine versus capecitabine alone in elderly patients with previously untreated metastatic colorectal cancer (AVEX): an open-label, randomised phase 3 trial. *Lancet Oncol* 2013;14:1077-1085.
- ²³Cunningham D, Pyrhonen S, James R, et al. Randomised trial of irinotecan plus supportive care versus supportive care alone after fluorouracil failure for patients with metastatic colorectal cancer. *The Lancet* 1998;352:1413-1418.
- ²⁴Fuchs CS, Moore MR, Harker G, et al. Phase III comparison of two irinotecan dosing regimens in second-line therapy of metastatic colorectal cancer. *J Clin Oncol* 2003;21:807-814.
- ²⁵Van Cutsem E, Tejpar S, Vanbeckevoort D, et al. Inpatient Cetuximab Dose Escalation in Metastatic Colorectal Cancer According to the Grade of Early Skin Reactions: The Randomized EVEREST Study. *J Clin Oncol* 2012;30:2861-2868.
- ²⁶Van Cutsem E, Peeters M, Siena S, et al. Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol* 2007;25:1658-1664.
- ²⁷Grothey A, Van Cutsem E, Sobrero A, et al. Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. *Lancet* 2013;381:303-312.
- ²⁸Mayer RJ, Van Cutsem E, Falcone A, et al. Randomized Trial of TAS-102 for Refractory Metastatic Colorectal Cancer (RECORSE). *N Engl J Med* 2015;372:1909-19.
- ²⁹Le DT, Uram JN, Wang H, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509-2520.
- ³⁰Overman MJ, Kopetz S, McDermott RS, et al. Nivolumab {+/-} ipilimumab in treatment of patients with metastatic colorectal cancer (mCRC) with and without high microsatellite instability (MSI-H): CheckMate-142 interim results [abstract]. *ASCO Meeting Abstracts* 2016;34:3501.

Note: All recommendations are category 2A unless otherwise indicated.**Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.**

**PRINCIPLES OF SURVIVORSHIP - Colorectal Long-term Follow-up Care****Colorectal Cancer Surveillance:**

- See [REC-8](#)
- Long-term surveillance should be carefully managed with routine good medical care and monitoring, including cancer screening, routine health care, and preventive care.
- Routine CEA monitoring and routine CT scanning are not recommended beyond 5 years.

Management of Late Sequelae of Disease or Treatment:¹⁻⁵**[See NCCN Guidelines for Survivorship](#)**

- Chronic diarrhea or incontinence
 - ▶ Consider anti-diarrheal agents, bulk-forming agents, diet manipulation, pelvic floor rehabilitation, and protective undergarments.
- For oxaliplatin-induced neuropathy
 - ▶ Consider duloxetine for painful neuropathy
- For fatigue
 - ▶ Encourage physical activity, energy conservation measures
- Urogenital dysfunction after resection and/or pelvic radiation^{6,7}
 - ▶ Screen for sexual dysfunction, erectile dysfunction, dyspareunia, and vaginal dryness
 - ▶ Screen for urinary incontinence, frequency, and urgency
 - ▶ Consider referral to urologist or gynecologist for persistent symptoms.
- Potential for pelvic fractures/decreased bone density after pelvic radiation
 - ▶ Consider bone density monitoring

Survivorship Care Planning:

The oncologist and primary care provider should have defined roles in the surveillance period, with roles communicated to patient.⁸

- Develop survivorship care plan that includes:
 - ▶ Overall summary of treatment, including all surgeries, radiation treatments, and chemotherapy received.
 - ▶ Description of possible expected time to resolution of acute toxicities, long-term effects of treatment, and possible late sequelae of treatment.

- ▶ Surveillance recommendations.
- ▶ Delineate appropriate timing of transfer of care with specific responsibilities identified for primary care physician and oncologist.
- ▶ Health behavior recommendations.

Cancer Screening Recommendations:

These recommendations are for average-risk patients.

Recommendations for high-risk individuals should be made on an individual basis.

- Breast Cancer: [NCCN Guidelines for Breast Cancer Screening](#)
- Prostate Cancer: [NCCN Guidelines for Prostate Early Detection](#)

Counseling Regarding Healthy Lifestyle and Wellness:⁹**[See NCCN Guidelines for Survivorship](#)**

- Maintain a healthy body weight throughout life.
- Adopt a physically active lifestyle (At least 30 minutes of moderate-intensity activity on most days of the week). Activity recommendations may require modification based on treatment sequelae (ie, ostomy, neuropathy).
- Consume a healthy diet with an emphasis on plant sources. Diet recommendations may be modified based on severity of bowel dysfunction.
- Consider low-dose aspirin
- Limit alcohol consumption.
- Seek smoking cessation counseling as appropriate.

Additional health monitoring and immunizations should be performed as indicated under the care of a primary care physician. Survivors are encouraged to maintain a therapeutic relationship with a primary care physician throughout their lifetime.

[See NCCN Guidelines for Survivorship](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.

PRINCIPLES OF SURVIVORSHIP - Colorectal Long-term Follow-up Care

References

- ¹Schneider EC, Malin JL, Kahn KL, et al. Surviving colorectal cancer. *Cancer* 2007;110: 2075-2082.
- ²Sprangers MAG, Taal BG, Aaronson NK, et al. Quality of life in colorectal cancer: stoma vs. nonstoma patients. *Dis Colon Rectum* 1995;38:361-369.
- ³Gami B, Harrington K, Blake P, et al. How patients manage gastrointestinal symptoms after pelvic radiotherapy. *Aliment Pharmacol Ther* 2003;18:987-994.
- ⁴DeSnoo L, Faithfull S. A qualitative study of anterior resection syndrome: the experiences of cancer survivors who have undergone resection surgery. *Eur J Cancer Care* 2006;15:244-251.
- ⁵McGough C, Baldwin C, Frost C, Andreyev HJN. Role of nutritional intervention in patients treated with radiotherapy for pelvic malignancy. *Br J Cancer* 2004;90:2278-2287.
- ⁶Lange MM, Mass CP, Marijnen CAM, et al. Risk factors for sexual dysfunction after rectal cancer treatment. *Eur J Cancer* 2009;45:1578-88.
- ⁷Lange MM, Mass CP, Marijnen CAM, et al. Urinary dysfunction after rectal cancer treatment is mainly caused by surgery. *Brit J Cancer* 2008;95:1020-28.
- ⁸Hewitt M, Greenfield S, Stovall E. *From Cancer Patient to Cancer Survivor: Lost in Transition*. Washington, D.C.:The National Academies Press;2006.
- ⁹Kushi LH, Byers T, Doyle C, et al and The American Cancer Society 2006 Nutrition and Physical Activity Guidelines Advisory Committee. *American Cancer Society Guidelines on Nutrition and Physical Activity for Cancer Prevention: Reducing the Risk of Cancer With Healthy Food Choices and Physical Activity* *CA Cancer J Clin* 2006;56:254-281.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any patient with cancer is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 3.2017 Staging Rectal Cancer

Table 1. Definitions for T, N, M**Primary Tumor (T)**

TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
Tis	Carcinoma in situ: intraepithelial or invasion of lamina propria ^a
T1	Tumor invades submucosa
T2	Tumor invades muscularis propria
T3	Tumor invades through the muscularis propria into the pericolorectal tissues
T4a	Tumor penetrates to the surface of the visceral peritoneum ^b
T4b	Tumor directly invades or is adherent to other organs or structures ^{b,c}

Regional Lymph Nodes (N)

NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in 1-3 regional lymph nodes
N1a	Metastasis in one regional lymph node
N1b	Metastasis in 2-3 regional lymph nodes
N1c	Tumor deposit(s) in the subserosa, mesentery, or nonperitonealized pericolic or perirectal tissues without regional nodal metastasis
N2	Metastasis in four or more regional lymph nodes
N2a	Metastasis in 4-6 regional lymph nodes
N2b	Metastasis in seven or more regional lymph nodes

Distant Metastasis (M)

M0	No distant metastasis
M1	Distant metastasis
M1a	Metastasis confined to one organ or site (eg, liver, lung, ovary, nonregional node)
M1b	Metastases in more than one organ/site or the peritoneum

^aTis includes cancer cells confined within the glandular basement membrane (intraepithelial) or mucosal lamina propria (intramucosal) with no extension through the muscularis mucosae into the submucosa.

^bDirect invasion in T4 includes invasion of other organs or other segments of the colorectum as a result of direct extension through the serosa, as confirmed on microscopic examination (for example, invasion of the sigmoid colon by a carcinoma of the cecum) or, for cancers in a retroperitoneal or subperitoneal location, direct invasion of other organs or structures by virtue of extension beyond the muscularis propria (ie, respectively, a tumor on the posterior wall of the descending colon invading the left kidney or lateral abdominal wall; or a mid or distal rectal cancer with invasion of prostate, seminal vesicles, cervix, or vagina).

^cTumor that is adherent to other organs or structures, grossly, is classified cT4b. However, if no tumor is present in the adhesion, microscopically, the classification should be pT1-4a depending on the anatomical depth of wall invasion. The V and L classifications should be used to identify the presence or absence of vascular or lymphatic invasion whereas the PN site-specific factor should be used for perineural invasion.

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.

Table 2. Anatomic Stage/Prognostic Groups

Stage	T	N	M	Dukes*	MAC*
0	Tis	N0	M0	-	-
I	T1	N0	M0	A	A
	T2	N0	M0	A	B1
IIA	T3	N0	M0	B	B2
IIB	T4a	N0	M0	B	B2
IIC	T4b	N0	M0	B	B3
IIIA	T1-T2	N1/N1c	M0	C	C1
	T1	N2a	M0	C	C1
IIIB	T3-T4a	N1/N1c	M0	C	C2
	T2-T3	N2a	M0	C	C1/C2
IIIC	T1-T2	N2b	M0	C	C1
	T4a	N2a	M0	C	C2
	T3-T4a	N2b	M0	C	C2
IVB	T4b	N1-N2	M0	C	C3
	Any T	Any N	M1a	-	-
IVA	Any T	Any N	M1b	-	-

Note: cTNM is the clinical classification, pTNM is the pathologic classification.

The y prefix is used for those cancers that are classified after neoadjuvant pretreatment (eg, ypTNM). Patients who have a complete pathologic response are ypT0N0cM0 that may be similar to Stage Group 0 or I. The r prefix is to be used for those cancers that have recurred after a disease-free interval (rTNM).

*Dukes B is a composite of better (T3 N0 M0) and worse (T4 N0 M0) prognostic groups, as is Dukes C (Any TN1 M0 and Any T N2 M0). MAC is the modified Astler-Coller classification.



Discussion

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Table of Contents

Overview	MS-2
Literature Search Criteria and Guidelines Update Methodology	MS-2
Risk Assessment	MS-3
TNM Staging	MS-5
Pathology	MS-5
The Role of Vitamin D in Colorectal Cancer	MS-8
Clinical Presentation and Treatment of Nonmetastatic Disease	MS-8
Management of Polypoid Cancer.....	MS-8
Management of Localized Rectal Cancer.....	MS-9
Clinical Evaluation/Staging.....	MS-9
Surgical Approaches.....	MS-11
Neoadjuvant and Adjuvant Therapy for Resectable Nonmetastatic Disease	MS-15
Recommendations for Patients with T1 and T2 Lesions.....	MS-25
Recommendations for Patients with T3-4 Lesions, Nodal Involvement, Locally Unresectable Disease, or Who Are Medically Inoperable	MS-26

Principles of the Management of Metastatic Disease.....	MS-26
Surgical Management of Colorectal Metastases	MS-27
Local Therapies for Metastases	MS-28
Hepatic Arterial Infusion	MS-28
Arterially Directed Embolic Therapy	MS-29
Liver- or Lung-Directed Radiation.....	MS-29
Tumor Ablation	MS-30
Peritoneal Carcinomatosis.....	MS-31
Determining Resectability.....	MS-32
Conversion to Resectability	MS-32
Neoadjuvant and Adjuvant Therapy for Resectable Metastatic Disease.....	MS-34
Perioperative Bevacizumab for Resectable Metastatic Disease	MS-36
Perioperative Cetuximab and Panitumumab for Resectable Metastatic Disease.....	MS-36
Systemic Therapy for Advanced or Metastatic Disease	MS-37
Predictive Markers for Cetuximab and Panitumumab.....	MS-37
Predictive Marker for Pembrolizumab and Nivolumab: MSI Status	MS-42
Recommendations for Treatment of Resectable Synchronous Metastases	MS-43
Recommendations for Treatment of Unresectable Synchronous Metastases	MS-43
Recommendations for Treatment of Metachronous Metastases	MS-44
Endpoints for Advanced Colorectal Cancer Clinical Trials.....	MS-45
Post-Treatment Surveillance	MS-46
Surveillance for Locoregional Disease	MS-46
Surveillance for Metastatic Disease	MS-48
Managing an Increasing CEA Level	MS-48
Treatment of Locally Recurrent Disease.....	MS-49
Survivorship	MS-49
Healthy Lifestyles for Survivors of Colorectal Cancer	MS-50
Secondary Chemoprevention for Colorectal Cancer Survivors	MS-52
Summary.....	MS-52
References.....	MS-54

Overview

Colorectal cancer is the fourth most frequently diagnosed cancer and the second leading cause of cancer death in the United States. In 2016, an estimated 39,220 new cases of rectal cancer will occur in the United States (23,110 cases in men; 16,110 cases in women). During the same year, it is estimated that 49,190 people will die from rectal and colon cancer combined.¹ Despite these statistics, the incidence per 100,000 population of colon and rectal cancers decreased from 60.5 in 1976 to 46.4 in 2005.² In fact, the incidence of colorectal cancer decreased at a rate of approximately 3% per year or greater between 2003 and 2012.¹ The incidence rate for colorectal cancer reported by the CDC for 2011 is 40.0 per 100,000 persons.³ In addition, mortality from colorectal cancer decreased by almost 35% from 1990 to 2007,⁴ and is currently down by about 50% from peak mortality rates.¹ These improvements in incidence of and mortality from colorectal cancer are thought to be a result of cancer prevention and earlier diagnoses through screening and of better treatment modalities.

Despite the observed improvements in the overall colorectal cancer incidence rate, a retrospective cohort study of the SEER CRC registry found that the incidence of colorectal cancer in patients younger than 50 years has been increasing.⁵ The authors estimate that the incidence rates for colon and rectal cancers will increase by 90.0% and 124.2%, respectively, for patients 20 to 34 years by 2030. The cause of this trend is currently unknown.

This Discussion summarizes the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines) for managing rectal cancer. These guidelines begin with the clinical presentation of the patient to the primary care physician or gastroenterologist and address diagnosis, pathologic staging, neoadjuvant treatment, surgical management,

adjuvant treatment, management of recurrent and metastatic disease, patient surveillance, and survivorship. These guidelines overlap considerably with the NCCN Guidelines for Colon Cancer, especially in the treatment of metastatic disease. The recommendations in these guidelines are classified as category 2A except where noted. The panel unanimously endorses patient participation in a clinical trial over standard or accepted therapy, especially for cases of advanced disease and for patients with locally aggressive colorectal cancer who are receiving combined modality treatment.

Literature Search Criteria and Guidelines Update Methodology

Prior to the update of this version of the NCCN Guidelines for Rectal Cancer, an electronic search of the PubMed database was performed to obtain key literature in the field of colorectal cancer published between June 12, 2015 and June 12, 2016, using the following search terms: (colon cancer) OR (colorectal cancer) OR (rectal cancer). The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes only peer-reviewed biomedical literature.⁶

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase III; Clinical Trial, Phase IV; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic Reviews; and Validation Studies.

The PubMed search resulted in 375 citations, and their potential relevance was examined. The data from key PubMed articles and articles from additional sources deemed as relevant to these Guidelines and discussed by the panel have been included in this version of the Discussion section (eg, e-publications ahead of print, meeting



abstracts). Recommendations for which high-level evidence is lacking are based on the panel's review of lower-level evidence and expert opinion.

The complete details of the Development and Update of the NCCN Guidelines are available on the NCCN website (www.NCCN.org).

Risk Assessment

Approximately 20% of cases of colorectal cancer are associated with familial clustering, and first-degree relatives of patients with colorectal adenomas or invasive colorectal cancer⁷⁻¹¹ are at increased risk for colorectal cancer. Genetic susceptibility to colorectal cancer includes well-defined inherited syndromes, such as Lynch syndrome (also known as hereditary nonpolyposis colorectal cancer [HNPCC])^{12,13} and familial adenomatous polyposis (FAP).¹⁴ Therefore, it is recommended that all patients with colorectal cancer be queried regarding their family history and considered for risk assessment, as detailed in the NCCN Guidelines for Colorectal Cancer Screening (to view the most recent version of these guidelines, visit the NCCN website at www.NCCN.org). Results from a recent randomized controlled trial suggest that most individuals without a personal history of colorectal cancer and with one first-degree relative with colorectal cancer diagnosed before age 50 years or two first-degree relatives with colorectal cancer diagnosed at any age can safely be screened with colonoscopy every 6 years.¹⁵

Lynch Syndrome

Lynch syndrome is the most common form of genetically determined colorectal cancer predisposition, accounting for 2% to 4% of all colorectal cancer cases.^{12,13,16,17} This hereditary syndrome results from germline mutations in DNA mismatch repair (MMR) genes (*MLH1*, *MSH2*, *MSH6*, and *PMS2*). Although identifying a germline mutation in

an MMR gene through sequencing is definitive for Lynch syndrome, patients usually undergo selection by considering family history and performing an initial test on tumor tissue before sequencing. One of two different initial tests can be performed on colorectal cancer specimens to identify individuals who might have Lynch syndrome: immunohistochemical analysis for MMR protein expression, which is often diminished because of mutation, or analysis for microsatellite instability (MSI), which results from MMR deficiency and is detected as changes in the length of repetitive DNA elements in tumor tissue caused by the insertion or deletion of repeated units.¹⁸ Testing the *BRAF* gene for mutation is indicated when immunohistochemical analysis shows that *MLH1* expression is absent in the tumor. The presence of a *BRAF* mutation indicates that *MLH1* expression is down-regulated through somatic methylation of the promoter region of the gene and not through a germline mutation.¹⁸

Many NCCN Member Institutions and other comprehensive cancer centers now perform immunohistochemistry and sometimes MSI testing on all newly diagnosed colorectal and endometrial cancers regardless of family history to determine which patients should have genetic testing for Lynch syndrome.¹⁹⁻²² The cost effectiveness of this approach, referred to as universal or reflex testing, has been confirmed for colorectal cancer, and this approach has been endorsed by the Evaluation of Genomic Applications in Practice and Prevention (EGAPP) working group at the CDC.²³⁻²⁵ The US Multi-Society Task Force on Colorectal Cancer also recommends universal genetic testing of tumors of all patients with newly diagnosed colorectal cancer, as does the American Gastroenterological Association.^{26,27} The Cleveland Clinic recently reported on its experiences implementing such a screening approach.²⁸



The NCCN Colon/Rectal Cancer Panel endorses universal MMR or MSI testing of all patients with a personal history of colon or rectal cancer to identify individuals with Lynch syndrome. This testing is also relevant for treatment selection in stage IV disease (see *Predictive Marker for Pembrolizumab and Nivolumab: MSI Status*, below). Furthermore, MMR/MSI status may be a prognostic marker for rectal cancer, with 5-year rectal cancer-specific survival rates of 100% reported for patients with stage I/II dMMR tumors.²⁹ An infrastructure needs to be in place to handle the screening results in either case. A more detailed discussion is available in the NCCN Guidelines for Colorectal Cancer Screening (available online at www.NCCN.org).

Molecular Classification of Colon and Rectal Cancer

Colorectal cancer is a heterogeneous disease. An international consortium has recently reported a molecular classification, defining four different subtypes: CMS1 (MSI Immune), hypermutated, microsatellite unstable (see *Lynch Syndrome*, above), with strong immune activation; CMS2 (Canonical), epithelial, chromosomally unstable, with marked WNT and MYC signalling activation; CMS3 (Metabolic), epithelial, with evident metabolic dysregulation; and CMS4 (Mesenchymal), prominent transforming growth factor β activation, stromal invasion, and angiogenesis.³⁰ However, this classification is not yet recommended in clinical practice.

Other Risk Factors for Colorectal Cancer

It is well recognized that individuals with inflammatory bowel disease (ie, ulcerative colitis, Crohn's disease) are at an increased risk for colorectal cancer.³¹⁻³³ Other possible risk factors for the development of colorectal cancer include smoking, the consumption of red and processed meats, alcohol consumption, diabetes mellitus, low levels of physical activity, metabolic syndrome, and obesity/high body mass

index (BMI).^{32,34-54} In fact, in the EPIC cohort of almost 350,000 individuals, those who adhered to 5 healthy lifestyle factors (healthy weight, physical activity, non-smoking, limited alcohol consumption, healthy diet) had a hazard ratio (HR) for the development of colorectal cancer of 0.63 (95% CI, 0.54–0.74) compared with those who adhered to ≤ 1 of the factors. Other large studies support the conclusion that adherence to healthy lifestyle factors can reduce the risk of colorectal cancer.^{55,56}

Some data suggest that consumption of dairy may lower risk for the development of colorectal cancer.^{57,58} However, a recent systematic review and meta-analysis of 15 cohort studies (>900,000 subjects; >5200 cases of colorectal cancer) only found an association between risk for colon cancer in men and the consumption of nonfermented milk.⁵⁹ No association was seen for rectal cancer in men or for colon or rectal cancer in women, and no association was seen for either cancer in either gender with consumption of solid cheese or fermented milk. Large cohort studies and meta-analyses suggest that other dietary factors may also lower the risk for colorectal cancer, including the consumption of fish and legumes.⁶⁰⁻⁶² Furthermore, the use of aspirin or nonsteroidal anti-inflammatory drugs may also decrease the risk for colorectal cancer.⁶³⁻⁶⁸

In addition, some data suggest that smoking, metabolic syndrome, obesity, and red/processed meat consumption is associated with a poor prognosis.^{37,69-72} Conversely, post-diagnosis fish consumption may be associated with a better prognosis.⁷³ A family history of colorectal cancer increases risk while improving prognosis.⁷⁴ Data on the effect of dairy consumption on prognosis after diagnosis of colorectal cancer are conflicting.^{75,76}

The relationship between diabetes and colorectal cancer is complex. Whereas diabetes and insulin use may increase the risk of developing colorectal cancer, treatment with metformin appears to decrease risk, at least in women.⁷⁷⁻⁸⁴ Results of a small randomized study suggest that 1 year of low-dose metformin in non-diabetic patients with previously resected colorectal adenomas or polyps may reduce the likelihood of subsequent adenomas or polyps.⁸⁵ In addition, although patients with colorectal cancer and diabetes appear to have a worse prognosis than those without diabetes,⁸⁶ patients with colorectal cancer treated with metformin seem to have a survival benefit.⁸⁷ The data regarding the effects of metformin on colorectal cancer incidence and mortality, however, are not completely consistent, with some studies seeing no effect.^{88,89}

TNM Staging

The NCCN Guidelines for Rectal Cancer adhere to the current TNM (tumor, node, metastases) staging system of the AJCC Cancer Staging Manual (Table 1 of the guidelines).⁹⁰ The TNM categories reflect very similar survival outcomes for rectal and colon cancer; these diseases therefore share the same staging system.

The 8th edition of the AJCC Staging Manual was released in 2016, with implementation scheduled for January 1, 2018.⁹⁰ In this edition, T1 tumors involve the submucosa; T2 tumors penetrate through the submucosa into the muscularis propria; T3 tumors penetrate through the muscularis propria; T4a tumors directly penetrate to the surface of the visceral peritoneum; and T4b tumors directly invade or are adherent to other organs or structures.⁹⁰

Regional lymph node classification includes N1a (1 positive lymph node); N1b (2–3 positive lymph nodes), N2a (4–6 positive nodes); and N2b (7 or more positive nodes). In addition, tumor deposit(s) in the

subserosa, mesentery, or non-peritonealized pericolic or perirectal tissues without regional nodal metastasis (ie, satellite tumor nodules) have been classified as N1c. Within each T stage, survival is inversely correlated with N stage (N0, N1a, N1b, N2a, and N2b).⁹⁰

Metastatic disease is classified as M1a when metastases are to only one site/solid organ (including to lymph nodes outside the primary tumor regional drainage area). M1b is used for metastases to multiple distant sites or solid organs, exclusive of peritoneal carcinomatosis. The 8th edition of the AJCC Cancer Staging Manual includes the M1c category for peritoneal carcinomatosis with or without blood-borne metastasis to visceral organs.⁹⁰ Patients with peritoneal metastases have a shorter PFS and OS than those without peritoneal involvement.⁹¹

The prefixes “p” and “yp” used in TNM staging denote pathologic staging and pathologic staging following neoadjuvant therapy, respectively.⁹⁰

Pathology

Pathologic staging information is provided by examination of the surgical specimen. Some of the information that should be detailed in the report of the pathologic evaluation of rectal cancer includes: 1) gross description of the tumor and specimen; 2) grade of the cancer; 3) depth of penetration and extension to adjacent structures (T); 4) number of regional lymph nodes evaluated; 5) number of positive regional lymph nodes (N); 6) the presence of distant metastases to other organs or sites including non-regional lymph nodes (M); 7) the status of proximal, distal, circumferential (radial), and mesenteric margins⁹²⁻⁹⁶; 8) neoadjuvant treatment effect^{97,98}; 9) lymphovascular invasion (LVI)⁹⁹; 10) perineural invasion (PNI)¹⁰⁰⁻¹⁰²; and 11) the number of tumor deposits.¹⁰³⁻¹⁰⁷



Margins

The 8th edition of the AJCC Cancer Staging Manual includes the suggestion that the surgeon mark the area of the specimen with the deepest tumor penetration so that the pathologist can directly evaluate the status of the resection margins.⁹⁰

The circumferential margin or circumferential resection margin (CRM) is an important pathologic staging parameter in rectal cancer.¹⁰⁸ The radial margin for resected segments of the colon that are completely encased by a peritonealized (serosal) surface is also referred to as the peritoneal margin. The CRM is very important in segments of the colon or rectum that are either not encased or only partially encased in peritoneum.¹⁰⁸

The CRM is the closest radial margin between the deepest penetration of the tumor and the edge of resected soft tissue around the rectum (ie, the retroperitoneal or subperitoneal aspect of the tumor) or from the edge of a lymph node and should be measured in millimeters.

Identification of the CRM is determined through evaluation of the outer circumference of the rectal and mesorectal specimen that often requires inking of the outer surfaces and “bread-loaf” slicing of the specimen.¹⁰⁹

The panel defines a positive CRM as tumor within 1 mm from the transected margin.^{94,96,110}

Accurate pathologic assessment of the CRM of resected rectal tumor specimens is very important since the CRM has been shown to be a strong predictor of both local recurrence and overall survival (OS),^{108,110-112} including in patients undergoing neoadjuvant therapy,^{95,113} and is an important consideration when postoperative treatment decisions are made. Furthermore, in a retrospective study of over 17,000 patients with rectal cancer, CRM was found to be a better predictor of local recurrence for patients undergoing surgery as initial therapy than for those who had received preoperative therapy.⁹⁵ CRM positivity based

solely on intranodal tumor should be noted as such; some studies have shown that positive intranodal CRM is associated with lower recurrence rates than a positive CRM by direct tumor extension. Additional components of the pathologic evaluation of the surgical specimen following a total mesorectal excision (TME) are described under *Surgical Approaches*, below.

Lymph Nodes

The AJCC and the College of American Pathologists (CAP) recommend evaluation of 12 lymph nodes to accurately identify early-stage colorectal cancers, respectively.^{90,108,114} The number of lymph nodes that can be retrieved varies with age and gender of the patient and on tumor grade or site.¹¹⁵ The literature lacks consensus regarding the minimal number of lymph nodes needed to accurately identify early-stage rectal cancer.¹¹⁶ Most of these studies have combined rectal and colon cancers with surgery as the initial treatment. Two studies confined only to rectal cancer have reported 14 and >10 lymph nodes as the minimal number to accurately identify stage II rectal cancer.^{117,118} A more recent analysis of patients with stage I or II rectal cancer in the SEER database found that OS improved with greater numbers of lymph nodes retrieved.¹¹⁹ Furthermore, the mean number of lymph nodes retrieved from rectal cancers treated with neoadjuvant therapy is significantly less than those treated by surgery alone (13 vs. 19, $P < .05$; 7 vs. 10, $P \leq .0001$).¹²⁰⁻¹²² In fact, retrieval of fewer lymph nodes may be a marker of a higher tumor response and better prognosis following neoadjuvant treatment.^{123,124}

Results of studies evaluating the sentinel node for micrometastatic disease through use of hematoxylin and eosin (H&E) staining to identify small foci of tumor cells and the identification of particular tumor antigens through immunohistochemical analysis have been



reported.^{125,126} Although results of some of these studies seem promising, there is no uniformity in the definition of “true” clinically relevant metastatic carcinoma. Some studies have considered detection of single cells by immunohistochemistry or by H&E, so-called isolated tumor cells (ITCs), to be micrometastasis.^{126,127} In addition, results of one study demonstrated that, following neoadjuvant radiotherapy for rectal cancer, the sensitivity for the sentinel node procedure was only 40%.¹²⁸ Furthermore, in a recent study involving 156 patients with colon cancer and 44 patients with rectal cancer, this “ultra-staging” of lymph nodes only changed the staging for 1% of patients.¹²⁹ Others have noted that micrometastasis found in node-negative patients did not predict outcome.¹³⁰ In contrast, a recent meta-analysis found that the presence of micrometastases increases the likelihood of disease recurrence, whereas the presence of ITCs does not.¹³¹

There is also potential benefit of assessing regional lymph nodes for ITCs. One study of 312 consecutive patients with pN0 disease found that positive cytokeratin staining was associated with a higher risk of recurrence.¹³² Relapse occurred in 14% of patients with positive nodes compared to 4.7% of those with negative nodes (HR, 3.00; 95% CI, 1.23–7.32; $P = .013$). A recent systematic review and meta-analysis came to a similar conclusion, finding decreased survival in patients with pN0 disease with immunohistochemical or reverse transcriptase polymerase chain reaction (RT-PCR) evidence of tumor cells in regional nodes.¹³³

Response to Treatment

The most recent CAP Guidelines require that the pathology report comment on treatment effects of neoadjuvant therapy.¹¹⁴ The tumor response should be graded on a scale of 0 (complete response – no

viable cancer cells observed) to 3 (poor response – minimal or no tumor kill; extensive residual cancer).^{97,98,114,134}

Perineural Invasion

Several studies have demonstrated that the presence of PNI is associated with a significantly worse prognosis.^{100-102,135-137} For example, one retrospective analysis of 269 consecutive patients who had colorectal tumors resected at one institution found a 4-fold greater 5-year survival in patients without PNI versus patients whose tumors invaded nearby neural structures.¹⁰¹ Multivariate analysis of patients with stage II rectal cancer showed that patients with PNI have a significantly worse 5-year disease-free survival (DFS) compared to those without PNI (29% vs. 82%; $P = .0005$).¹⁰² Similar results were seen for patients with stage III disease.¹⁰⁰ A meta-analysis that included 58 studies and 22,900 patients also found that PNI is associated with a worse 5-year OS (Relative risk [RR], 2.09; 95% CI, 1.68–2.61) and 5-year DFS (RR, 2.35; 95% CI, 1.66–3.31).¹³⁶ PNI is therefore included as a high-risk factor for systemic recurrence.

Tumor Deposits

Tumor deposits, or satellite nodules, are irregular discrete tumor deposits in the perirectal fat that are away from the leading edge of the tumor and show no evidence of residual lymph node tissue, but that are within the lymphatic drainage of the primary tumor. They are not counted as lymph nodes replaced by tumor. Most of these tumor deposits are thought to be due to LVI or occasionally PNI. The number of tumor deposits should be recorded in the pathology report, since they have been shown to be associated with reductions in DFS and OS.^{103-107,137} Multivariate survival analysis in one study showed that patients with pN0 tumors without satellite nodules had a 91.5% 5-year survival rate compared to 37.0% for patients with pN0 tumors and the presence



of satellite nodules ($P < .0001$).¹⁰⁷ Another retrospective study found a similar difference in 5-year OS rates (80.3% vs. 34.9%, respectively; $P < .001$).¹³⁸ The association of tumor deposits with decreased survival also holds in patients with rectal cancer who had neoadjuvant chemoradiation (chemoRT).^{139,140} Tumor deposits are classified as pN1c.⁹⁰

The Role of Vitamin D in Colorectal Cancer

Prospective studies have suggested that vitamin D deficiency may contribute to colorectal cancer incidence and/or that vitamin D supplementation may decrease colorectal cancer risk.¹⁴¹⁻¹⁴⁵

Furthermore, several prospective studies have shown that low vitamin D levels are associated with increased mortality of patients with colorectal cancer.¹⁴⁶⁻¹⁴⁹ In fact, a systematic review and meta-analysis of 5 studies totaling 2330 patients with colorectal cancer compared the outcomes of patients in the highest and lowest categories of vitamin D levels and found better OS (HR, 0.71; 95% CI, 0.55–0.91) and disease-specific mortality (HR, 0.65; 95% CI, 0.49–0.86) in those with higher vitamin D levels.¹⁵⁰ Another meta-analysis determined that the relationship between vitamin D levels and mortality is linear.¹⁵¹

Results of a recent randomized, double-blind, placebo-controlled trial, however, showed that supplementation with vitamin D and/or calcium had no effect on the recurrence of colorectal adenomas within 3 to 5 years after removal of adenomas in 2259 participants.¹⁵² Furthermore, no study has yet examined whether vitamin D supplementation improves outcomes in patients with colorectal cancer. In a 2010 report, the Institute of Medicine concluded that data supporting a role for vitamin D were only conclusive in bone health, not in cancer and other diseases.¹⁵³ Citing this report and the lack of level 1 evidence, the panel does not currently recommend routine screening for vitamin D

deficiency or supplementation of vitamin D in patients with colorectal cancer.

Clinical Presentation and Treatment of Nonmetastatic Disease

Management of Polypoid Cancer

Before making a decision about surgical resection for an endoscopically resected adenomatous polyp or villous adenoma, physicians should review pathology¹⁵⁴ and consult with the patient. A malignant rectal polyp is defined as one with cancer invading through the muscularis mucosae and into the submucosa (pT1).¹⁵⁵ Conversely, polyps classified as carcinoma in situ (pTis) have not penetrated into the submucosa and are therefore incapable of regional nodal metastasis.¹⁰⁸ The panel recommends marking the cancerous polyp site at the time of colonoscopy or within 2 weeks if deemed necessary by the surgeon.

In patients with pedunculated or sessile polyps (adenomas), no additional surgery is required if the polyp has been completely resected with favorable histologic features.^{154,156} Favorable histologic features include lesions of grade 1 or 2 without angiolymphatic invasion and with a negative resection margin.¹⁵⁴ For patients with a completely removed, single-specimen, sessile polyp (pT1) with favorable histologic features and clear margins, observation may be considered, with the understanding that there is significantly greater incidence of adverse outcomes (residual disease, recurrent disease, mortality, and hematogenous metastasis, but not lymph node metastasis) than with polypoid malignant polyps. Also see the section on *Endoscopically Removed Malignant Polyps* in *Principles of Pathologic Review* in the algorithm. Rectal surgery is also an option for these patients.

Rectal surgery is also recommended for patients with polyps with unfavorable histologic features or when the specimen is fragmented or



margins cannot be assessed. Unfavorable histologic features for adenomas are grade 3 or 4, angiolymphatic invasion, or a positive margin of resection. In such cases, risk of nodal involvement is higher. It should be noted that no consensus currently exists as to the definition of what constitutes a positive margin of resection. A positive margin for an endoscopically removed polyp has been defined as the presence of tumor within 1 to 2 mm from the transected margin or by the presence of tumor cells within the diathermy of the transected margin.^{154,157-159} In addition, several studies have shown that tumor budding is an adverse histological feature associated with adverse outcome and may preclude polypectomy as an adequate treatment of endoscopically removed malignant polyps.¹⁶⁰⁻¹⁶³

For a polyp with fragmented specimen or margins that cannot be assessed, either a transanal excision or a transabdominal resection is recommended. In patients with unfavorable pathologic features, transabdominal resection should be considered in order to include lymphadenectomy. Results from a preoperative endoscopic ultrasound (EUS) evaluation may provide additional information to guide choice of surgical approach, although the accuracy of this method to detect residual cancer is limited (see section on *Clinical Evaluation/Staging*, below).¹⁶⁴ All patients who have malignant polyps removed by transanal excision or transabdominal resection should undergo total colonoscopy to rule out other synchronous polyps and should undergo surveillance as described in the guidelines.

Management of Localized Rectal Cancer

Rectal cancer has been defined as a cancerous lesion located within 12 cm of the anal verge by proctoscopy.¹⁶⁵ Some support for this definition comes from the study by Kapiteijn et al,¹⁶⁶ which included a subgroup analysis of the risk of recurrence of rectal cancer based on tumor

location. Univariate analyses indicated that local recurrence rates were low for patients who had tumors with an inferior margin of 10.1 cm or more from the anal verge, and that no significant differences between patients in this group receiving radiotherapy and surgery were observed when they were compared to those undergoing surgery alone.¹⁶⁶ A retrospective review of patients with rectal or rectosigmoid cancer demonstrated that treatment options were impacted by whether the location of the rectal lesion was characterized by proctoscopy or colonoscopy.¹⁶⁷

The determination of an optimal treatment plan for an individual patient with rectal cancer is a complex process. In addition to decisions relating to the intent of rectal cancer surgery (ie, curative or palliative), consideration must also be given to the likely functional results of treatment, including the probability of maintaining or restoring normal bowel function/anal continence and preserving genitourinary functions. For patients with distal rectal cancer, in particular, the simultaneous achievement of the goals of cure and of minimal impact on quality of life can be challenging.¹⁶⁸ Furthermore, the risk of pelvic recurrence is higher in patients with rectal cancer compared to those with colon cancer, and locally recurrent rectal cancer has frequently been associated with a poor prognosis.¹⁶⁹⁻¹⁷¹ Careful patient selection with respect to particular treatment options and the use of sequenced multimodality therapy that combines chemoRT with operative treatment for selected patients is recommended.¹⁷²

Clinical Evaluation/Staging

The initial clinical workup of patients with rectal cancer provides important preoperative information on the clinical stage of disease. Since the clinical stage is used to direct decisions regarding choice of primary treatment, including surgical intent (eg, curative or palliative) and approaches and whether to recommend preoperative chemoRT,



the implications of either clinically understaging or overstaging rectal cancer can be substantial.

Patients who present with rectal cancer appropriate for resection require a complete staging evaluation, including total colonoscopy to evaluate for synchronous lesions or other pathologic conditions of the colon and rectum. They also require a complete physical examination, including carcinoembryonic antigen (CEA) determination and assessment of performance status to determine operative risk. In addition, the accessibility of rectal cancer to evaluation by pelvic MRI with contrast (preferred) or endorectal ultrasound makes possible preoperative assessments of depth of tumor penetration and the presence of local lymph nodal metastases.¹⁷³

Additional information regarding the extent of disease and the occurrence of distant metastases can be determined preoperatively through CT scans. Thus, pelvic MRI (preferred) or endorectal ultrasound and CT scans of the chest, abdomen, and pelvis are recommended for the preoperative staging of rectal cancer. CT should be with IV and oral contrast, and if the CT of the abdomen and pelvis is inadequate or if CT with IV contrast is contraindicated, an abdominal/pelvic MRI with contrast plus a non-contrast chest CT should be considered. The chest CT can identify lung metastases, which occur in approximately 4% to 9% of patients with colon and rectal cancer.¹⁷⁴⁻¹⁷⁶ A series of 378 patients found that resection of pulmonary metastases resulted in 3-year recurrence-free survival of 28% and 3-year OS of 78%.¹⁷⁷

The consensus of the panel is that a PET scan is not indicated. PET/CT, if done, does not supplant a contrast-enhanced diagnostic CT scan. PET/CT should only be used to evaluate an equivocal finding on

a contrast-enhanced CT scan or in patients with a strong contraindication to IV contrast.

Results from a meta-analysis of 90 studies involving the accuracy of EUS, MRI, and CT for preoperative staging of rectal cancer demonstrated that EUS and MRI have similarly high sensitivities for evaluating the depth of tumor penetration into the muscularis propria (94%), although EUS was found to be more specific than MRI in the evaluation of local tumor invasion (86% vs. 69%).¹⁷⁸ Only a very limited number of studies using CT for the purpose of T-staging have been performed, and it is not currently considered to be an optimal method for staging the extent of tumor penetration.^{178,179} Accurate assessment of nodal status is one of the greatest challenges in the preoperative staging of rectal cancer. In the meta-analysis of Bipat et al,¹⁷⁸ the sensitivities and specificities of the 3 imaging modalities for accurately evaluating lymph node involvement were comparable: CT (55% and 74%); EUS (67% and 78%); and MRI (66% and 76%). However, only CT and MRI can evaluate iliac and mesenteric or retroperitoneal nodes.¹⁷⁸ Results from another recent meta-analysis of 84 articles indicated that none of the 3 imaging modalities were significantly superior to another method with respect to an accurate determination of tumor N-stage.¹⁸⁰ A disadvantage of EUS is a high degree of operator dependence.¹⁷⁸ An advantage of MRI is its ability to provide accurate images of soft tissue structures in the mesorectum, including the mesorectal fascia so as to provide information useful in the prediction of the CRM prior to radical surgery.¹⁷⁹⁻¹⁸² Recently published 5-year follow-up results of the MERCURY trial show that high-resolution MRI can accurately assess the CRM preoperatively, differentiating patients with low-risk and high-risk disease.¹⁸³ Patients with MRI-clear CRM had a 5-year OS of 62.2% compared with 42.2% in patients with MRI-involved CRM (HR, 1.97; 95% CI, 1.27–3.04; $P < .01$). The preoperative MRI



imaging also predicted DFS (HR, 1.65; 95% CI, 1.01–2.69; $P < .05$) and local recurrence (HR, 3.50; 95% CI, 1.53–8.00; $P < .05$). A group of experts developed consensus guidelines for standardized imaging of rectal cancer by MRI.¹⁸⁴

Clinical staging is also based on histopathologic examination of the specimen obtained via biopsy or local excision (eg, excised polyps). Endoscopic biopsy specimens of the lesion should undergo careful pathology review for evidence of invasion into the muscularis mucosa. If removal of the rectum is contemplated, early consultation with an enterostomal therapist is recommended for preoperative marking of the site and patient teaching purposes.

Restaging / Assessing Treatment Response

Restaging after neoadjuvant treatment is done to plan the surgical approach and, increasingly, to determine if additional therapy or resection can be avoided for select patients. Future and ongoing trials will help to answer these questions (see *Wait-and-See Nonoperative Approach for Clinical Complete Responders* and *Preoperative Chemotherapy Without Chemoradiation*, below). MRI, CT, and EUS are the most commonly used modalities for restaging after neoadjuvant treatment, but the accuracy of these techniques for determining T stage and lymph node involvement is limited.¹⁸⁵⁻¹⁹³ Advanced functional MRI techniques (eg, dynamic contrast-enhanced MRI, diffusion-weighted MRI) allow for the measurement of microcirculation, vascular permeability, and tissue cellularity and thus may be useful for determining response to neoadjuvant treatment and restaging patients with rectal cancer.^{192,194-196} FDG-PET/CT is also being investigated for its ability to accurately determining response to neoadjuvant treatment.^{195,197}

Surgical Approaches

A variety of surgical approaches, depending on the location and extent of disease, are used to treat primary rectal cancer lesions.^{198,199} These methods include local procedures, such as polypectomy, transanal excision, and transanal endoscopic microsurgery (TEM), and more invasive procedures involving a transabdominal resection (eg, low anterior resection [LAR], proctectomy with TME and coloanal anastomosis, abdominoperineal resection [APR]).^{198,199}

Transanal Excision

Transanal excision is only appropriate for selected T1, N0 early-stage cancers. Small (<3 cm), well to moderately differentiated tumors that are within 8 cm of the anal verge and limited to less than 30% of the rectal circumference and for which there is no evidence of nodal involvement can be approached with transanal excision with negative margins.²⁰⁰

TEM can facilitate excision of small tumors through the anus when lesions can be adequately identified in the rectum. TEM may be technically feasible for more proximal lesions. Although data are limited, a 2015 meta-analysis found that TEM may achieve superior oncologic outcomes compared with transanal excision.²⁰¹ Both transanal excision and TEM involve a full-thickness excision performed perpendicularly through the bowel wall into the perirectal fat. Negative (>3 mm) deep and mucosal margins are required, and tumor fragmentation should be avoided.

The locally excised specimen should be oriented and pinned before fixation and brought to the pathologist by the surgeon to facilitate an oriented histopathologic evaluation of the specimen. If pathologic examination reveals adverse features such as positive margins, LVI,



poor differentiation, or invasion into the lower third of the submucosa (sm3 level),^{202,203} a more radical resection is recommended.

Data are limited on long-term patient outcomes, including risk of local recurrence, for patients undergoing local excision for T2 tumors.²⁰⁴ Results of a multi-institutional, single-arm, open-label, non-randomized, phase II trial suggest that chemoradiotherapy with CapeOx followed by local excision may be a safe alternative to transabdominal resection in patients with T2N0 distal rectal cancer.²⁰⁵ A meta-analysis also suggests that this approach of neoadjuvant chemoRT followed by local excision may be a safe and effective alternative for patients with any T and any N stage of rectal cancer who refuse or are unfit for transabdominal resection.²⁰⁶ Further studies in this area are needed.

Advantages of a local procedure include minimal morbidity (eg, a sphincter-sparing procedure) and mortality and rapid postoperative recovery.^{168,204} Limitations of a local excision include the absence of pathologic staging of nodal involvement. Further, evidence indicates that lymph node micrometastases are both common in early rectal lesions and unlikely to be identified by endorectal ultrasound.²⁰⁷ These observations may underlie the findings that patients undergoing local excision have a higher local recurrence rate than those undergoing radical resection.^{204,208,209} A retrospective study of 282 patients undergoing either transanal excision or radical resection for T1 rectal cancer from 1985 to 2004 showed respective local recurrence rates of 13.2% and 2.7% for these 2 groups ($P = .001$).²⁰⁹ A similar retrospective study of 2124 patients showed local recurrence rates of 12.5% and 6.9% for patients undergoing local excision versus standard resection, respectively ($P = .003$).²⁰⁴ More recently, an analysis of >164,000 individuals from the National Cancer Data Base with resected, invasive, nonmetastatic rectal cancer diagnosed from 1998 to 2010 found that positive margins were more likely after local excision compared to

transabdominal excision in both the T1 and T2 populations (95% vs. 76% in T1/T2 combined; $P < .001$).²¹⁰ In the T1N0 population, a small but significant decrease in OS was also noted in the local excision group. Interestingly, limited data suggest that TEM might have superior oncologic outcomes in patients with stage I rectal cancer compared with radical resection,^{208,211} although not all studies have seen such results.²¹²

Thus, careful patient selection for local excision of T1N0 rectal cancer is important, as is the careful examination of the resection specimen with subsequent transabdominal resection in patients found to have T2 disease or high-risk features, as described above.

Transabdominal Resection

Patients with rectal cancer who do not meet requirements for local surgery should be treated with a transabdominal resection. Organ-preserving procedures that maintain sphincter function are preferable, but not possible in all cases. Preoperative chemoRT may result in tumor downsizing and a decrease in tumor bulk (see section on *Neoadjuvant and Adjuvant Therapy for Resectable Nonmetastatic Disease*, below); sphincter preservation may become possible in cases where initial tumor bulk prevented consideration of such surgery and exposure to the tumor is improved by chemoRT.

In transabdominal resections, TME is recommended. A TME involves an en bloc removal of the mesorectum, including associated vascular and lymphatic structures, fatty tissue, and mesorectal fascia as a “tumor package” through sharp dissection and is designed to spare the autonomic nerves.^{168,199,213} The lymphatic drainage regions of rectal tumors are influenced by their position in the rectum. More distal tumors are more likely to be characterized by both upward and lateral lymphatic drainage, whereas the likelihood of only upward mesorectal drainage is much higher for more proximal tumors.²¹⁴ The TME approach is



designed to radically remove lymphatic drainage regions of tumors located above the level of the levator muscles.²¹⁵ The panel does not recommend extension of nodal dissection beyond the field of resection (eg, into the distribution of iliac lymph nodes) unless these nodes are clinically suspicious. In cases where anal function is intact and distal clearance is adequate, the TME may be followed by creation of a coloanal anastomosis.

For lesions in the mid to upper rectum, an LAR extended 4 to 5 cm below the distal edge of the tumor using TME, followed by creation of a colorectal anastomosis, is the treatment of choice. Where creation of an anastomosis is not possible, colostomy is required. Wide TME is recommended in order to facilitate adequate lymphadenectomy and improve the probability of achieving negative circumferential margins.

An APR with TME should be performed when the tumor directly involves the anal sphincter or the levator muscles. An APR is also necessary in cases where a margin-negative resection of the tumor would result in loss of anal sphincter function and incontinence. An APR involves en bloc resection of the rectosigmoid, the rectum, and the anus, as well as the surrounding mesentery, mesorectum (TME), and perianal soft tissue, and it necessitates creation of a colostomy.²¹⁶ In the NSABP R-04 trial, patients who had an APR reported worse body image, worse micturition symptoms, and less sexual enjoyment at 1-year post surgery than those who had sphincter-sparing surgery.²¹⁷ An extralevator APR may have benefits over a conventional APR approach, including lower rates of CRM involvement and local recurrence.²¹⁸

Pathologists play a key role following TME in evaluating the surgical specimen, including a macroscopic assessment of both its external appearance/completeness and the CRM.^{219,220} The panel defines a

positive CRM as tumor within 1 mm from the transected margin (see *Pathology*, above).^{94,96,110} Detailed descriptions of how the quality of the mesorectal specimens should be scored were provided in the Dutch Rectal Cancer Trial, and these guidelines are endorsed by the NCCN Panel.⁹⁴

Recent retrospective comparisons of the outcomes of patients undergoing an APR versus an LAR in the treatment of rectal cancer have shown that those treated with an APR have worse local control and OS.^{221,222} Whether these differences can be attributed to the surgical procedure alone, to patient- and tumor-related characteristics, or some combination of these factors is presently unclear. However, results from a recent retrospective study of 3633 patients with T3-4 rectal cancer tumors included in 5 large European trials suggest that there is an association between the APR procedure itself and the increased risks of recurrence and death.²²¹ Importantly, quality of life between patients with or without a permanent colostomy appears to be fairly comparable.^{223,224}

Laparoscopic Resection

Data from randomized studies evaluating use of laparoscopic surgery in the treatment of patients with rectal cancer have been maturing in recent years.²²⁵⁻²²⁸

One large prospective multicentre study, which included 4405 patients with rectal cancer but was not randomized, found no differences in recurrence or survival, although complications and other measures of quality indicated a benefit to the laparoscopic approach.²²⁹ The phase III COLOR II trial, powered for non-inferiority, also randomized patients with localized rectal cancer to laparoscopic or open surgery. Short-term secondary endpoints were met, with patients in the laparoscopic arm losing less blood, having shorter hospital stays, and having a quicker



return of bowel function, but with longer operation times.²³⁰ No differences were seen in completeness of resection, percentage of patients with positive CRM, morbidity, or mortality between the arms. The primary endpoint of locoregional recurrence at 3 years was identical in the 2 groups, at 5.0%, and no statistically significant differences were seen in DFS or OS.²²⁵

In the CLASICC trial comparing laparoscopically assisted resection to open resection, nearly half of the 794 patients were diagnosed with rectal cancer.²³¹ No significant differences in local recurrence, DFS, or OS were observed between the 2 groups of patients with colon or rectal cancer based on surgical approach. A 5-year follow-up of the CLASICC trial showed that this lack of difference in local recurrence, DFS, or OS was maintained for patients with rectal cancer, despite a trend towards better 5-year OS after laparoscopic surgery (52.9% and 60.3% for open and laparoscopic surgery, respectively; $P = .132$).²³²

The COREAN trial randomized patients with stage II or III low- to mid-rectal cancer to an open or laparoscopic resection, with short-term benefits seen to the laparoscopic approach.²³³ The primary endpoint, 3-year DFS, did not differ between the 2 groups at 72.5% (95% CI, 65.0–78.6) for open surgery and 79.2% (95% CI, 72.3–84.6) for the laparoscopic group.²²⁶ Factors that may confound conclusions drawn from randomized studies comparing open surgery to laparoscopically assisted surgery for colorectal cancer have been described,²³⁴ and longer-term outcomes from laparoscopic rectal surgery have not been reported.

Two other trials, ACOSOG Z6051 and ALaCaRT, have reported pathologic outcomes.^{227,228} In Z6051, the primary endpoint was a composite of CRM >1 mm, negative distal margin, and TME completeness.²²⁷ No significant differences were observed between the

arms in these 3 measures or in the composite of successful resection. For example, complete or nearly complete TME was achieved in 92.1% (95% CI, 88.7–95.5) in the laparoscopic resection arm and 95.1% (95% CI, 92.2–97.9) in the open resection arm, for a difference of -3.0 (95% CI, -7.4 – 1.5 ; $P = .20$). However, the criteria for non-inferiority of the laparoscopic approach were not met. In ALaCaRT, the primary endpoint was also a composite of resection quality measures.²²⁸ Successful resections were achieved in 82% of the laparoscopic resection arm and 89% of the open resection arm, for a difference of -7.0% (95% CI, -12.4% to infinity). A negative CRM was achieved in 93% and 97%, respectively (risk difference, -3.7% ; 95% CI, -7.6% – 0.1% ; $P = .06$). As in Z6051, the criteria for non-inferiority of the laparoscopic approach were not met in ALaCaRT. Longer follow-up with oncologic outcomes of these trials are needed.

Reviews and meta-analyses including these and additional small trials have also been published.^{226,235-248} They consistently find the laparoscopic approach to be safe and feasible. In addition, an analysis of results from >18,000 individuals in the National Cancer Data Base undergoing LAR for rectal cancer found short-term oncologic outcomes to be similar between the open and laparoscopic approaches.²⁴⁹

In conclusion, some studies have shown that laparoscopy is associated with similar short- and long-term outcomes when compared to open surgery,^{225,226} whereas other studies have shown the laparoscopic approach to be associated with higher rates of CRM positivity and incomplete TME.^{227,228} Therefore, the panel defined principles by which laparoscopic resection of rectal cancer can be considered: the procedure can be considered by an experienced surgeon, should include thorough abdominal exploration, and should be limited to lower-risk tumors, as outlined in the guidelines. An international group of



experts has defined standards for the technical details of laparoscopic TME.²⁵⁰

Neoadjuvant and Adjuvant Therapy for Resectable Nonmetastatic Disease

Neoadjuvant/adjuvant therapy of stage II (T3-4, node-negative disease with tumor penetration through the muscle wall) or stage III (node-positive disease without distant metastasis) rectal cancer often includes locoregional treatment due to the relatively high risk of locoregional recurrence. This risk is associated with the close proximity of the rectum to pelvic structures and organs, the absence of a serosa surrounding the rectum, and technical difficulties associated with obtaining wide surgical margins at resection. In contrast, adjuvant treatment of colon cancer is more focused on preventing distant metastases since this disease is characterized by lower rates of local recurrence.

Although radiation therapy (RT) has been associated with decreased rates of local recurrence of rectal cancer, it is also associated with increased toxicity (eg, radiation-induced injury, hematologic toxicities) relative to surgery alone.^{109,251,252} It has been suggested that some patients with disease at lower risk of local recurrence (eg, proximal rectal cancer staged as T3, N0, M0, characterized by clear margins and favorable prognostic features) may be adequately treated with surgery and adjuvant chemotherapy.^{109,253,254} However, 22% of 188 patients clinically staged with T3, N0 rectal cancer by either EUS or MRI who subsequently received preoperative chemoRT had positive lymph nodes following pathologic review of the surgical specimens according to results of a retrospective multicenter study,²⁵⁵ suggesting that many patients are under-staged and would benefit from chemoRT. Therefore, the guidelines recommend preoperative chemoRT for patients with T3, N0 disease.

Combined-modality therapy consisting of surgery, concurrent fluoropyrimidine-based chemotherapy with ionizing radiation to the pelvis (chemoRT), and chemotherapy is recommended for the majority of patients with stage II or stage III rectal cancer. Use of perioperative pelvic RT in the treatment of patients with stage II/III rectal cancer continues to evolve. In these patients, the current guidelines recommend 2 possible sequences of therapy: 1) chemoRT preoperatively and chemotherapy postoperatively; or 2) chemotherapy followed by chemoRT followed by resection. The total duration of perioperative therapy, including chemoRT and chemotherapy, should not exceed 6 months.

Preoperative Versus Postoperative Radiation

Several studies have compared the administration of radiation preoperatively versus postoperatively for stage II/III rectal cancer.^{256,257} A large prospective, randomized trial from the German Rectal Cancer Study Group (the CAO/ARO/AIO-94 trial) compared preoperative versus postoperative chemoRT in the treatment of clinical stage II/III rectal cancer.²⁵⁶ Results of this study indicated that preoperative therapy was associated with a significant reduction in local recurrence (6% vs. 13%; $P = .006$) and treatment-associated toxicity (27% vs. 40%; $P = .001$), although OS was similar in the 2 groups. Long-term follow-up of this trial was later published.²⁵⁸ The improvement in local control persisted, with the 10-year cumulative incidence of local recurrence at 7.1% and 10.1% in the preoperative and postoperative treatment arms, respectively ($P = .048$). OS at 10 years was again similar between the groups (59.6% and 59.9%, respectively; $P = .85$), as was DFS and the occurrence of distant metastases.

Interestingly, a recent SEER database analysis of 4724 patients with T3N0 rectal cancer found that radiation given after resection was associated with a significant decrease in risk for cancer death compared



to surgery without any radiation (HR, 0.69; 95% CI, 0.58–0.82; $P < .001$), while radiation given before resection was not (HR, 0.86; 95% CI, 0.72–1.04; $P = .13$).²⁵⁹

Putative advantages to preoperative radiation, as opposed to radiation given postoperatively, are related to both tumor response and preservation of normal tissue.^{256,257,260} First of all, reducing tumor volume may facilitate resection and increase the likelihood of a sphincter-sparing procedure. Although some studies have indicated that preoperative radiation or chemoRT is associated with increased rates of sphincter preservation in patients with rectal cancer,^{256,257} this conclusion is not supported by 2 meta-analyses of randomized trials involving preoperative chemoRT in the treatment of rectal cancer.^{261,262} Second, irradiating tissue that is surgery-naïve and thus better oxygenated may result in increased sensitivity to RT. Third, preoperative radiation can avoid the occurrence of radiation-induced injury to small bowel trapped in the pelvis by post-surgical adhesions. Finally, preoperative radiation that includes structures that will be resected increases the likelihood that an anastomosis with healthy colon can be performed (ie, the anastomosis remains unaffected by the effects of RT because irradiated tissue is resected).

One disadvantage of using preoperative RT is the possibility of over-treating early-stage tumors that do not require adjuvant radiation.^{256,263} Improvements in preoperative staging techniques, such as MRI or CT scans, have allowed for more accurate staging, but the risk of over-staging disease has not been eliminated.²⁵⁵ Weighing these advantages and disadvantages, the panel recommends preoperative chemoRT for patients with stage II/III rectal cancer. Postoperative chemoRT is recommended when stage I rectal cancer is upstaged to stage II or III after pathologic review of the surgical specimen. Postoperative chemoRT regimens commonly employ a “sandwich” approach –

whereby chemotherapy (typically 5-FU–based) is administered before and after the chemoRT regimen.^{254,264,265}

Concurrent Chemotherapy with Radiation

A number of randomized trials have evaluated the effectiveness of the addition of chemotherapy to radiation administered either preoperatively following clinical evaluation/staging (eg, T3-4 by EUS) or postoperatively following pathologic staging of rectal cancer as pT3 and/or N1-2.²⁶⁶ Putative benefits of the addition of chemotherapy concurrent with either pre- or postoperative RT include local RT sensitization and systemic control of disease (ie, eradication of micrometastases). Preoperative chemoRT also has the potential to increase rates of pathologic complete response and sphincter preservation.

In a study of patients with T3-4 rectal cancer without evidence of distant metastases who were randomly assigned to receive either preoperative RT alone or preoperative concurrent chemoRT with 5-FU/LV, no difference in OS or sphincter preservation was observed in the 2 groups, although patients receiving chemoRT were significantly more likely to exhibit a pathologic complete response (11.4% vs. 3.6%; $P < .05$) and grade 3/4 toxicity (14.6% vs. 2.7%; $P < .05$) and less likely to exhibit local recurrence of disease (8.1% vs. 16.5%; $P < .05$).²⁶⁶

Preliminary results of a phase III trial that included an evaluation of the addition of chemotherapy to preoperative RT in patients with T3-4 resectable rectal cancer demonstrated that use of 5-FU/LV chemotherapy enhanced the tumoricidal effect of RT when the 2 approaches were used concurrently.²⁶⁷ Significant reductions in tumor size, pTN stage, and lymphatic, vascular, and PNI rates were observed with use of combined-modality therapy compared with use of RT and surgery without chemotherapy.²⁶⁷ More mature results from this trial,



which included 4 treatment groups (preoperative RT; preoperative chemoRT; preoperative RT plus postoperative chemotherapy; and preoperative chemoRT plus postoperative chemotherapy), however, indicated that no significant differences in OS were associated with adding 5-FU-based chemotherapy preoperatively or postoperatively.²⁶⁸

The conclusions of these trials have been supported in a 2009 systematic review that included 4 randomized controlled trials.²⁶⁹ In addition, a recent Cochrane review of 6 randomized controlled trials found that chemotherapy added to preoperative radiation in patients with stage III, locally advanced rectal cancer reduced the risk of local recurrence, but had no effect on OS, 30-day mortality, sphincter preservation, and late toxicity.²⁷⁰ Similarly, a separate Cochrane review of stage II and III resectable disease found that the addition of chemotherapy to preoperative radiation enhances pathologic response and improves local control, but has no effect on DFS or OS.²⁷¹ Another recent meta-analysis of 5 randomized controlled trials comparing neoadjuvant chemoRT with neoadjuvant radiotherapy came to similar conclusions.²⁵²

With respect to the type of chemotherapy administered concurrently with RT,²⁵⁴ the equivalence of bolus 5-FU/LV and infusional 5-FU in concurrent chemoRT for rectal cancer is supported by the results of a phase III trial (median follow-up of 5.7 years) in which similar outcomes with respect to OS and relapse-free survival were observed when an infusion of 5-FU or bolus 5-FU plus LV was administered concurrently with postoperative RT, although hematologic toxicity was greater in the group of patients receiving bolus 5-FU.²⁶⁵ On the other hand, results from an earlier trial from the North Central Cancer Treatment Group (NCCTG) showed that postoperative administration of infusional 5-FU during pelvic irradiation was associated with longer OS when compared to bolus 5-FU.²⁶⁴ Most of the patients in this study had node-positive

disease. The panel considers bolus 5-FU/LV/RT as an option for patients not able to tolerate capecitabine or infusional 5-FU (both preferred in the chemoRT setting).

Recent studies have shown that capecitabine is equivalent to 5-FU in perioperative chemoRT therapy.^{272,273} The randomized NSABP R-04 trial compared the preoperative use of infusional 5-FU with or without oxaliplatin to capecitabine with or without oxaliplatin in 1608 patients with stage II or III rectal cancer.^{273,274} No differences in locoregional events, DFS, OS, complete pathologic response, sphincter-saving surgery, or surgical downstaging were seen between the regimens, while toxicity was increased with the inclusion of oxaliplatin.

Similarly, a phase III randomized trial in which 401 patients with stage II or III rectal cancer received capecitabine– or 5-FU–based chemoRT either pre- or postoperatively showed that capecitabine was non-inferior to 5-FU with regard to 5-year OS (capecitabine 75.7% vs. 5-FU 66.6%; $P = .0004$), with capecitabine showing borderline significance for superiority ($P = .053$).²⁷² Furthermore, in this trial capecitabine demonstrated a significant improvement in 3-year DFS (75.2% vs. 66.6%; $P = .034$).²⁷² Because of these studies, capecitabine given concurrently with RT is now listed in the guidelines as a category 2A recommendation. The panel feels that capecitabine is an acceptable alternative to infusional 5-FU in those patients who are able to manage the responsibilities inherent in self-administered, oral chemotherapy.

Addition of oxaliplatin: In attempts to improve on the outcomes achieved with neoadjuvant 5-FU/RT or capecitabine/RT, several large randomized phase III trials (ACCORD 12, STAR-01, R-04, CAO/ARO/AIO-04, and FOWARC) addressed the addition of oxaliplatin to the regimens. In a planned interim report of primary tumor response in the STAR-01 trial, grade 3 and 4 adverse events occurred more



frequently in patients receiving infusional 5-FU/oxaliplatin/RT than in those receiving infusional 5-FU/RT (24% vs. 8%, $P < .001$), while there was no difference in pathologic response between the arms of the study (16% pathologic complete response in both arms).²⁷⁵ Recently reported results of the NSABP R-04 trial also showed that the addition of oxaliplatin did not improve clinical outcomes including the endpoints of locoregional events, DFS, OS, pathologic complete response, sphincter-saving surgery, and surgical downstaging, while it increased toxicity.^{273,274} Further follow-up of these trials is necessary to see if there is a difference in local recurrence rates and progression-free survival (PFS) over time. The primary endpoints of OS for the STAR-01 trial will be reported in the future.

Similar results were seen in the ACCORD 12/0405-ProDIGe 2 trial, in which capecitabine/RT (45 Gy) was compared to CapeOx/RT (50 Gy) and the primary endpoint was pathologic complete response.²⁷⁶ The pathologic complete response rates were similar at 19.2% and 13.9% ($P = .09$) for the oxaliplatin-containing arm and the control arm, respectively. Although patients treated with oxaliplatin and the higher radiation dose in the ACCORD 12 trial had an increased rate of minimal residual disease at the time of surgery (39.4% vs. 28.9%, $P = .008$), this did not translate to improved local recurrence rates, DFS, or OS at 3 years.

Results of the German CAO/ARO/AIO-04 trial have been published.^{277,278} This trial also assessed the addition of oxaliplatin to a fluorouracil RT regimen. In contrast to STAR-01, R-04, and ACCORD 12, higher rates of pathologic complete response were seen in the oxaliplatin arm (17% vs. 13%, $P = .038$)²⁷⁸, but this result could be because of differences in the fluorouracil schedule between the arms.²⁷⁹ The primary endpoint of this trial, the 3-year DFS rate, was 75.9% (95% CI, 72.4%–79.5%) in the oxaliplatin arm versus 71.2% (95% CI, 67.6%–

74.9%) in the control group ($P = .03$).²⁷⁷ Importantly, oxaliplatin was also added to the adjuvant therapy in the AIO-04 trial but not in the other trials, so cross-trial comparisons are limited.

In line with CAO/ARO/AIO-04, the Chinese FOWARC phase III open-label, multicenter trial, which randomized patients with locally advanced rectal cancer to neoadjuvant treatment consisting of infusional 5-FU/LV-RT, FOLFOX-RT, or FOLFOX, found that FOLFOX-RT resulted in higher rates of pathologic complete response and downstaging than the other regimens.²⁸⁰

Based on the results available to date, the addition of oxaliplatin to neoadjuvant chemoRT is not recommended at this time.

Another randomized multicenter phase III trial looked at the addition of oxaliplatin during concurrent capecitabine chemoRT in the adjuvant setting for pathologic stage II/III disease.²⁸¹ Interim analysis showed no significant difference in 3-year DFS, OS, local recurrences, or distant metastases, with an increase in grade 3/4 acute toxicity in the CapeOx-RT group.

Addition of targeted agents: The randomized phase II EXPERT-C trial assessed complete response rate with the addition of cetuximab to radiation treatment in 165 patients.²⁸² Patients in the control arm received CapeOx followed by capecitabine/RT, then surgery followed by CapeOx. Patients randomized to the cetuximab arm received the same therapy with weekly cetuximab throughout all phases. A significant improvement in OS was seen in patients with *KRAS* exon 2/3 wild-type tumors treated with cetuximab (HR, 0.27; 95% CI, 0.07–0.99; $P = .034$). However, the primary endpoint of complete response rate was not met, and other phase II trials have not shown a clear benefit to the addition



of cetuximab in this setting.^{283,284} Further evaluation of this regimen is warranted.

The randomized, multicenter, phase II SAKK 41/07 trial evaluated the addition of panitumumab to preoperative capecitabine-based chemoRT in patients with locally advanced, *KRAS* wild-type rectal cancer.²⁸⁵ The primary endpoint was pathologic near-complete plus complete tumor response, which occurred in 53% (95% CI, 36%–69%) of patients in the panitumumab arm versus 32% (95% CI, 16%–52%) in the control arm. Patients receiving panitumumab experienced increased rates of grade 3 or greater toxicity.

A phase II study of 57 patients with resectable T3/T4 rectal cancer evaluated preoperative treatment with capecitabine, oxaliplatin, bevacizumab, and RT, followed by surgery 8 weeks later and adjuvant FOLFOX/bevacizumab.²⁸⁶ The 5-year OS rate was 80%, and the 5-year relapse-free survival rate was 81%. However, the primary endpoint of pathologic complete response was not met, significant toxicities were observed, and compliance with adjuvant therapy was low.

Additional phase II trials assessing the effects of adding irinotecan or bevacizumab to neoadjuvant or adjuvant regimens have begun.²⁸⁷⁻²⁸⁹ However, at this time the panel does not endorse the use of bevacizumab, cetuximab, panitumumab, irinotecan, or oxaliplatin with concurrent radiotherapy for rectal cancer.

Induction Chemotherapy

Several small trials have tested the utility of a course of neoadjuvant chemotherapy preceding chemoRT and resection.²⁹⁰⁻²⁹⁵ In the Spanish GCR-3 randomized phase II trial, patients were randomized to receive CapeOx either before chemoRT or after surgery.^{292,296} Similar pathologic complete response rates were seen, and induction chemotherapy

appeared to be less toxic and better tolerated. Another phase II trial randomized patients to chemoRT and surgery with or without FOLFOX induction therapy.²⁹⁴ There were no differences between the clinical outcomes, but the group receiving induction therapy experienced higher toxicity. The phase II AVACROSS study assessed the safety and efficacy of adding bevacizumab to induction therapy with CapeOx prior to capecitabine/bevacizumab-chemoRT and surgery.²⁹⁵ The regimen was well tolerated with a pathologic complete response rate of 36%.

Possible benefits of using chemotherapy first include the early prevention or eradication of micrometastases, higher rates of pathologic complete response, minimizing the time patients need an ileostomy, facilitating resection, and improving the tolerance and completion rates of chemotherapy. This approach was added to the 2015 version of these guidelines as an acceptable option.

Preoperative Chemotherapy Without Chemoradiation

A small single-center phase II pilot trial treated patients with stage II or III rectal cancer with induction FOLFOX/bevacizumab chemotherapy followed by chemoRT only in those with stable or progressive disease and resection in all patients.²⁹⁷ All 32 of the participants had an R0 resection, and the 4-year DFS was 84% (95% CI, 67%–94%). The ongoing N1048/C81001/Z6092 PROSPECT trial by The Alliance for Clinical Trials in Oncology is also asking whether chemotherapy alone is effective in treating stage II or III high rectal cancer in patients with at least 20% tumor regression following neoadjuvant treatment (clinicaltrials.gov NCT01515787).

The phase III FOWARC trial, discussed above, compared neoadjuvant therapy with and without radiation (without additional therapy for those with stable or progressive disease) and found that neoadjuvant FOLFOX without radiation gave lower rates of pathologic complete



response than regimens that included radiation (6.6% vs. 14.0% for 5-FU-RT and 27.5% for FOLFOX-RT).²⁸⁰ Rates of downstaging were similar to the 5-FU-RT (35.5% vs. 37.1% for 5-FU-RT and 56.4% for FOLFOX-RT).

A 2015 systematic review identified 1 randomized phase III trial, 6 single-arm phase II trials, and 1 retrospective case series study that addressed the effectiveness of neoadjuvant chemotherapy (without chemoRT) and surgery in patients with locally advanced rectal cancer.²⁹⁸ The ranges of R0 resection and pathological complete response rates were 90% to 100% and 4% to 33%, respectively.

This approach could spare patients the morbidities associated with radiation, but the panel considered it investigational at this time.

Technical Aspects of Radiation Therapy

With respect to administration of RT, multiple RT fields should include the tumor or tumor bed with a 2- to 5-cm margin, presacral nodes, and the internal iliac nodes. The external iliac nodes should also be included for T4 tumors involving anterior structures; inclusion of the inguinal nodes for tumors invading into the distal anal canal can also be considered. Recommended doses of radiation are typically 45 to 50 Gy in 25 to 28 fractions to the pelvis using 3 or 4 fields. Positioning and other techniques to minimize radiation to the small bowel are encouraged. The Radiation Therapy Oncology Group (RTOG) has established normal pelvic contouring atlases for females and males (available online at <http://www.rtog.org/CoreLab/ContouringAtlases.aspx>).²⁹⁹ Intensity-modulated RT (IMRT) should only be used in the setting of a clinical trial or in unique clinical situations such as re-irradiation of previously treated recurrent disease or unique anatomical situations.

Coordination of preoperative therapy, surgery, and adjuvant chemotherapy is important. Although longer intervals from completion of chemoRT to surgery have been shown to be associated with an increase in pathologic complete response rates,³⁰⁰⁻³⁰⁵ it is unclear whether such longer intervals are associated with clinical benefit. Results of one National Cancer Data Base analysis suggest that an interval of >8 weeks was associated with increased odds of pathologic complete response,³⁰⁶ whereas other similar analyses concluded that an interval >56 or 60 days (8–8.5 weeks) is associated with higher rates of positive margins, lower rates of sphincter preservation, and/or shorter survival.^{307,308}

The GRECCAR6 phase III, multicenter, randomized, open-label, parallel-group controlled trial randomized patients with stage II/III rectal cancer treated with chemoRT to a 7-week or an 11-week interval before surgery.³⁰⁹ The pathologic complete response rate was not different between the groups (15.0% vs. 17.4%; $P = .60$), but the morbidity (44.5% vs. 32%; $P = .04$), medical complications (32.8% vs. 19.2%; $P = .01$), and rate of complete mesorectal resection (78.7% vs. 90%; $P = .02$) were worse in the 11-week group. The rate of anastomotic leaks and the mean length of hospital stay were similar between the groups.

Based on these data, the panel recommends an interval of 5 to 12 weeks following completion of full-dose 5½-week chemoRT prior to surgical resection for patients treated with preoperative chemoRT in order to allow patient recuperation from chemoRT-associated toxicities.

Short-course Radiation

Several European studies have looked at the efficacy of a shorter course of preoperative radiation (25 Gy over 5 days), not combined with chemotherapy, for the treatment of rectal cancer. The results of the Swedish Rectal Cancer Trial evaluating the use of short-course RT



administered preoperatively for resectable rectal cancer showed a survival advantage and a decreased rate of local recurrence with this approach compared with surgery alone.³¹⁰ However, a follow-up study published in 2005 showed that the patients with short-course preoperative RT had increased relative risk for postoperative hospitalization due to bowel obstructions and other gastrointestinal complications.³¹¹ A number of other studies also investigating the effectiveness of preoperative short-course RT in patients with rectal cancer staged as T1-3 have demonstrated that OS was not significantly affected despite improvements in local control of disease.^{166,312,313} A more recent multicenter, randomized study of 1350 patients with rectal cancer compared 1) short-course preoperative RT and no postoperative treatment with 2) no preoperative RT and a postoperative approach that included chemoRT in selected patients (ie, those with a positive CRM following resection) and no RT in patients without evidence of residual disease following surgery.³¹⁴ Results indicated that patients in the preoperative RT arm had significantly lower local recurrence rates and a 6% absolute improvement in 3-year DFS ($P = .03$), although no difference in OS was observed between the arms of the study.^{314,315}

Long-term (12-year) follow-up of one of the short-course radiation trials (the Dutch TME trial³¹²) was reported.³¹⁶ The analysis showed that 10-year survival was significantly improved in patients with stage III disease with a negative circumferential margin in the radiotherapy plus surgery group compared to the group that received surgery alone (50% vs. 40%; $P = .032$).³¹⁶ However, this long follow-up showed that secondary malignancies and other non-rectal cancer causes of death were more frequent in the radiotherapy group than in the control group (14% vs. 9% for secondary malignancies), negating any survival advantage in the node-negative subpopulation.

A few studies have compared short-course RT to long-course chemoRT. One randomized study of 312 patients in Poland directly compared preoperative short-course radiation and more conventional preoperative long-course chemoRT and found no differences in local recurrence or survival.³¹⁷ Similarly, an Australian/New Zealand trial (Trans-Tasman Radiation Oncology Group [TROG] trial 01.04) that randomized 326 patients to short-course radiation or long-course chemoRT found no differences in local recurrence and OS rates.³¹⁸ In addition, rates of late toxicity, distant recurrence, and relapse-free survival were not significantly different between the arms. Patients in the long-course arm were more likely to experience serious adverse events (eg, radiation dermatitis rates, 0% vs. 5.6%; $P = .003$), whereas patients in the short-course arm were more likely to have a permanent stoma (38.0% vs. 29.8%; $P = .13$).³¹⁹ Finally, a trial compared short-course RT with long-course chemoRT with delayed surgery in both groups.³²⁰ Although the long-course arm experienced greater tumor downsizing and downstaging compared with short-course treatment, no differences were seen in the R0 resection rates or postoperative morbidity.

A 2014 systematic review identified 16 studies (randomized controlled trials, phase II trials, and retrospective studies) that addressed the interval between short-course radiation and resection of rectal cancer.³²¹ Lower rates of severe acute post-radiation toxicity but higher rates of minor postoperative complications were seen in the immediate-surgery group (1- to 2-week interval) compared with the delayed surgery group (5- to 13-week interval). The pCR rates were significantly higher in the delayed-surgery group, with no differences in sphincter preservation and R0 resection rates.

Overall, it appears that short-course RT gives effective local control and the same OS as more conventional RT schedules, and therefore is



considered as an appropriate option for patients with T3N0 or T1-3N1-2 rectal cancer. Short-course RT is not recommended for T4 disease. A multidisciplinary evaluation, including a discussion of the need for down-staging and the possibility of long-term toxicity, is recommended when considering short-course RT.

Response to Neoadjuvant Treatment

Fifty percent to 60% of patients are down-staged following neoadjuvant therapy, with about 20% of patients showing a pathologic complete response.³²²⁻³²⁸ Recent studies have suggested that the response to neoadjuvant treatment correlates with long-term outcomes in patients with rectal cancer. In the MERCURY prospective cohort trial, 111 patients were assessed by MRI and pathologic staging.³²⁹ On multivariate analysis, MRI-assessed tumor regression grade was significantly associated with OS and DFS. Patients with poor tumor regression grade had 5-year survival rates of 27% versus 72% for patients with good tumor regression grade ($P = .001$), and DFS rates were 31% versus 64% ($P = .007$). Similarly, in the CAO/ARO/AIO-94 trial, patients with pathologic complete regression had 10-year cumulative incidence of distant metastasis and DFS of 10.5% and 89.5%, respectively, while those with poor regression had corresponding incidences of 39.6% and 63%.³³⁰ A recent retrospective review of 725 patients with rectal cancer found similar results.³²⁶ In this study, pathologically determined response to neoadjuvant treatment correlated with long-term outcomes. Five-year recurrence-free survival rates were 90.5%, 78.7%, and 58.5% for patients with complete, intermediate, and poor responses, respectively ($P < .001$). Distant metastases and local recurrences also correlated with the level of response.

In addition to its prognostic value, there is some initial evidence of predictive value to neoadjuvant treatment response. Subgroup analysis

of the EORTC 22921 trial showed that patients down-staged to ypT0-2 were more likely to benefit from adjuvant chemotherapy than patients with ypT3-4 staging.³²² Similar results were seen from another retrospective review.³³¹ Although no prospective data to predict the benefit of adjuvant therapy in patients with tumor downstaging or a pathologic complete response exist, the panel believes that such patients should be strongly considered for adjuvant chemotherapy.

Wait-and-See Nonoperative Approach for Clinical Complete Responders

As preoperative treatment and imaging modalities have improved, some have suggested that patients with a clinical complete response to chemoRT may be able to be spared the morbidities of surgery. In 2004, Habr-Gama et al³³² retrospectively compared the outcomes of 71 patients who were observed without surgery following complete clinical response (27% of patients) to the outcome of 22 patients (8%) who had incomplete clinical responses but complete pathologic responses post-TME. The OS and DFS rates at 5 years were 100% and 92%, respectively, in the nonoperative group compared to 88% and 83%, respectively, in the resected group. However, other studies did not achieve as impressive results, and many clinicians were skeptical of the approach.³³³

A more recent prospective study included a more thorough assessment of treatment response and used very strict criteria to select 21 of 192 patients (11%) with clinical complete responses who were then observed with careful follow-up and compared to 20 patients with a complete pathologic response after resection.³³⁴ Only 1 patient in the nonoperative group developed a local recurrence after a mean follow-up of 25 months; that patient underwent successful surgery. No statistical differences in long-term outcomes were seen between the groups. The cumulative probabilities for 2-year DFS and OS were 89% (95% CI,



43%–98%) and 100%, respectively, in the wait-and-see group and 93% (95% CI, 59%–99%) and 91% (95% CI, 59%–99%), respectively, in the resected group. Short-term functional outcomes, however, were better in the wait-and-see group, with better bowel function scores, less incontinence, and 10 patients avoiding permanent colostomy.

Other non-randomized, prospective studies have added to the growing evidence that the nonoperative approach may warrant further study.³³⁵⁻

³³⁸ For example, one study showed that 49% of patients experienced a complete clinical response after 5-FU-based chemoRT, and found that strict surveillance in these patients, with resection of recurrences when possible, resulted in a 5-year recurrence-free survival of 69%, which rose to 94% after resections were performed.³³⁶

Despite these impressive results, many still believe that longer follow-up, larger sample sizes, and additional careful observational studies are needed before patients with a clinical complete response are routinely managed by a wait-and-see approach.³³⁹ Furthermore, recent studies have found that neither FDG-PET, nor MRI, nor CT can accurately determine a pathologic complete response, complicating the selection of appropriate patients for a nonoperative approach.^{185-193,340} In addition, lymph node metastases are still seen in a subset of patients with pathologic complete response.³⁴¹ Overall, the panel does not support this approach in the routine management of localized rectal cancer.

Adjuvant Chemotherapy

Adjuvant chemotherapy is recommended for all patients with stage II/III rectal cancer following neoadjuvant chemoRT and surgery if they did not receive neoadjuvant chemotherapy regardless of the surgical pathology results, although few studies have evaluated the effect of adjuvant chemotherapy in patients with rectal cancer, and its role is not well defined.^{342,343} The addition of 5-FU adjuvant chemotherapy to

preoperative chemoRT provided no benefit to the rate of local recurrence in the EORTC Radiotherapy Group Trial 22921.²⁶⁸ However, this study did show an improvement in DFS (HR, 0.87; 95% CI, 0.72–1.04; $P = .13$) of patients receiving adjuvant chemotherapy (+/- RT) following preoperative RT (+/- 5-FU–based chemotherapy).²⁶⁸ Long-term results of the 22921 trial confirmed that adjuvant 5-FU chemotherapy did not improve OS, and the difference in DFS was less pronounced than following the previous analysis (HR, 0.91; 95% CI, 0.77–1.08; $P = .29$).³⁴⁴ Limitations of this trial include the fact that only 43% of participants received the full course of adjuvant chemotherapy. Other trials have failed to show an improvement in OS or DFS with adjuvant therapy with a fluoropyrimidine alone in this setting.^{345,346}

Other trials have investigated the use of more modern agents in the adjuvant setting. The phase III ECOG E3201 trial was designed to investigate the effect of adding either oxaliplatin (FOLFOX) or irinotecan (FOLFIRI) to 5-FU/LV-based adjuvant chemotherapy administered to patients with stage II/III rectal cancer after either preoperative or postoperative chemoRT. This study was replaced with an alternative trial with bevacizumab, but results from an initial 165 patients indicate that adjuvant FOLFOX can be safely used in this patient population.³⁴⁷ The open-label phase II ADORE trial randomized 321 patients with resected rectal cancer and neoadjuvant therapy to adjuvant 5-FU/LV or FOLFOX.³⁴⁸ The FOLFOX arm had higher 3-year DFS, at 71.6% versus 62.9% (HR, 0.66; 95% CI, 0.43–0.99; $P = .047$). The CAO/ARO/AIO-04 trial found an improvement in 3-year DFS when oxaliplatin was added to 5-FU in both neoadjuvant and adjuvant treatment (75.9% vs. 71.2%; $P = .03$).²⁷⁷

A recent study in which patients who received neoadjuvant chemoRT and experienced a pathologic complete response were observed without additional adjuvant chemotherapy found 5-year DFS and OS



rates of 96% and 100%, respectively.³⁴⁹ In addition, a meta-analysis of 4 randomized trials (1196 patients) concluded that adjuvant fluorouracil-based chemotherapy (5-FU/LV, capecitabine, or CapeOx) after preoperative therapy and surgery did not improve OS, DFS, or the rate of distant recurrences in patients with stage II or III rectal cancer.³⁵⁰ However, more recent trials that found a DFS benefit to the addition of adjuvant oxaliplatin-based adjuvant therapy were not included in this study, and other meta-analyses have come to the opposite conclusion.^{351,352} The panel continues to support the use of adjuvant therapy in this setting.

A recent analysis of the NCCN Colorectal Cancer Database found that, of 2073 patients with stage II/III rectal cancer who received neoadjuvant chemoRT treatment, 203 patients (9.8%) did not receive any adjuvant chemotherapy as recommended by these guidelines.³⁵³ Multivariate analysis found that complete pathologic response, infection, no closure of ileostomy/colostomy, age, poor performance status, and being on Medicaid or indigent were associated with not receiving adjuvant chemotherapy. Results from the SEER database indicated that even fewer patients in the general population are receiving adjuvant therapy (61.5%) in this setting.³⁵⁴ Pathologic stage, age, and postoperative readmissions were associated with a decreased likelihood of receiving adjuvant treatment.

Although conclusive data on the use of adjuvant therapy in patients with stage II/III rectal cancer are lacking, the panel recommends use of FOLFOX or CapeOx as preferred options. 5-FU/leucovorin or capecitabine can also be used in this setting and might be especially appropriate in patients who responded to neoadjuvant treatment with 5-FU or capecitabine.

Timing and Duration of Adjuvant Therapy: A 2011 systematic review and meta-analysis of 10 studies involving more than 15,000 patients with colorectal cancer looked at the effect of timing of adjuvant therapy following resection.³⁵⁵ Results of this analysis showed that each 4-week delay in chemotherapy results in a 14% decrease in OS, indicating that adjuvant therapy should be administered as soon as the patient is medically able. These results are consistent with other similar analyses.³⁵⁶

The optimal duration of adjuvant treatment in rectal cancer is still unclear.^{357,358} In the MOSAIC trial, patients with stage II/III colon cancer were treated with 6 months of adjuvant FOLFOX.³⁵⁹ The use of a shorter course of adjuvant FOLFOX in rectal cancer (ie, 4 months) is justified when preoperative chemoRT is administered.

Multigene Assays

Several multigene assays have been developed in hopes of providing prognostic and predictive information to aid in decisions regarding adjuvant therapy in patients with stage II or III colon cancer (see the NCCN Guidelines for Colon Cancer, available at www.NCCN.org, for a full discussion).³⁶⁰

Among the multigene assays used in colon cancer is the Oncotype DX colon cancer assay, which quantifies the expression of 7 recurrence-risk genes and 5 reference genes as a prognostic classifier of low, intermediate, or high likelihood of recurrence.³⁶¹ Clinical validation in patients with stage II and III colon cancer from QUASAR and National Surgical Adjuvant Breast and Bowel Project (NSABP) C-07 trials showed that recurrence scores are prognostic for recurrence, DFS, and OS in stage II and III colon cancer, but are not predictive of benefit to adjuvant therapy.³⁶² For the low, intermediate, and high recurrence risk



groups, recurrence at 3 years was 12%, 18%, and 22%, respectively. Similar results were found in other prospectively designed studies.^{363,364}

A recent prospectively designed validation study assessed this assay for predicting recurrence risk in patients with stage II and III rectal cancer.³⁶⁵ For patients who underwent surgery without neoadjuvant therapy in the Dutch Total Mesorectal Excision (TME) trial, recurrence score was predictive of recurrence, distant recurrence, and rectal-cancer-specific survival. In patients with stage II rectal cancer, recurrence at 5 years was 11%, 27%, and 43% for the low, intermediate, and high recurrence risk groups, respectively.

The panel believes the information from this test can further inform the risk of recurrence over other risk factors, but they question the value added. Furthermore, there is no evidence of predictive value in terms of the potential benefit of chemotherapy in patients with colon or rectal cancer with any of the available multigene assays. The panel believes that there are insufficient data to recommend the use of multigene assays to determine adjuvant therapy for patients with colorectal cancer.

Leucovorin Shortage

A leucovorin shortage recently existed in the United States. No specific data guide management under these circumstances, and all proposed strategies are empiric. The panel recommends several possible options to help alleviate the problems associated with this shortage. One is the use of levo-leucovorin, which is commonly used in Europe. A dose of 200 mg/m² of levo-leucovorin is equivalent to 400 mg/m² of standard leucovorin. Another option is for practices or institutions to use lower doses of leucovorin for all doses in all patients, since the panel feels that lower doses are likely to be as efficacious as higher doses, based on several studies. The QUASAR study found that 175 mg leucovorin

gave similar survival and 3-year recurrence rates as 25 mg leucovorin when given with bolus 5-FU to patients as adjuvant therapy following R0 resections for colorectal cancer.³⁶⁶ Another study showed no difference in response rate or survival in patients with metastatic colorectal cancer receiving bolus 5-FU with either high-dose (500 mg/m²) or low-dose (20 mg/m²) leucovorin.³⁶⁷ Also, the Mayo Clinic and NCCTG determined that there was no therapeutic difference between the use of high- (200 mg/m²) or low- (20 mg/m²) dose leucovorin with bolus 5-FU in the treatment of advanced colorectal cancer, although 5-FU doses were different in the 2 arms.³⁶⁸ Finally, if none of the above options is available, treatment without leucovorin would be reasonable. For patients who tolerate this without grade II or higher toxicity, a modest increase in 5-FU dose (in the range of 10%) may be considered.

Recommendations for Patients with T1 and T2 Lesions

Node-negative T1 lesions are treated with transabdominal resection or transanal excision, as appropriate (see section on *Surgical Approaches*, above). If pathology review after local excision reveals no high-risk features or if a transabdominal resection was performed, then no additional treatment is required. If, however, pathology review after local excision reveals a poorly differentiated histology, positive margins, invasion into the lower third of the submucosa (sm3 level), or LVI or if the tumor is restaged to T2, additional treatment is required. The initial options are transabdominal re-resection or chemoRT. For resected patients with positive nodes or pT3-4 disease, chemotherapy and chemo RT “sandwich regimen” (as described below) should be given. For patients treated with transanal excision and then chemoRT, observation can be considered if a complete response is seen. Alternatively, the patient can undergo a transabdominal resection with optional adjuvant chemotherapy or chemotherapy can be given without resection.

Node-negative T2 lesions are treated with transabdominal resection, since local recurrence rates of 11% to 45% have been observed for T2 lesions following local excision alone.^{168,369,370} Following transabdominal resection, patients with tumors staged as pT1-2, N0, M0 require no further treatment. If pathology review reveals pT3-4, N0, M0 or node-positive disease, a “sandwich regimen,” consisting of: 1) an optional first round of adjuvant chemotherapy with 5-FU with or without LV or FOLFOX or capecitabine with or without oxaliplatin,³⁷¹ followed by 2) concurrent 5-FU/RT (infusional [preferred] or bolus infusion along with LV) or capecitabine/RT (preferred); followed by 3) 5-FU with or without LV or FOLFOX or capecitabine with or without oxaliplatin, is one option. Alternatively, these patients can be observed or can receive chemotherapy without radiation. Patients with pathologic findings of N1 disease should receive a “sandwich regimen” of chemotherapy and chemoRT.

The panel recommends perioperative therapy for a total duration of approximately 6 months.

Recommendations for Patients with T3-4 Lesions, Nodal Involvement, Locally Unresectable Disease, or Who Are Medically Inoperable

Patients clinically staged as having resectable T3-4, N0 T any, N1-2 lesions, and/or who have locally unresectable disease or are medically inoperable have 3 options for the sequence of treatment: 1) chemotherapy with long-course RT, then resection if possible, followed by chemotherapy; 2) short-course RT (not recommended for T4 disease), then resection if possible, followed by chemotherapy; or 3) chemotherapy, then chemoRT, then resection if possible. Infusional 5-FU/RT and capecitabine/RT are the preferred chemoRT options (category 1 for both) regardless of the sequence. An alternative chemoRT regimen is bolus 5-FU/LV/RT. The preferred chemotherapy regimens when given before surgery are FOLFOX or CapeOx, with 5-

FU/leucovorin and capecitabine as additional options. In the postoperative setting, adjuvant FOLFOX (preferred), CapeOx (preferred), 5-FU/leucovorin, or capecitabine should be given.

Resection should be considered following preoperative therapy unless there is a clear contraindication. The panel advises that a poor clinical response does not necessarily imply unresectability, and surgical exploration is usually appropriate. Transabdominal resection should be performed 5 to 12 weeks following completion of neoadjuvant therapy. The panel recommends that the duration of perioperative chemotherapy, including chemotherapy and chemoRT, be approximately 6 months. When resection is contraindicated following primary treatment, patients should be treated with a systemic regimen for advanced disease (see discussion of *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for Colon Cancer, available at www.NCCN.org). FOLFOXIRI is not recommended in this setting.

For unresectable cancers, doses higher than 54 Gy may be required; the dose of radiation to the small bowel should be limited to 45 Gy. For patients with T4 tumors or recurrent cancers or if margins are very close or positive, intraoperative RT (IORT),³⁷²⁻³⁷⁶ which involves direct exposure of tumors to RT during surgery while removing normal structures from the field of treatment, may be considered as an additional boost to facilitate resection. If IORT is not available, 10 to 20 Gy and/or brachytherapy to a limited volume can be considered.

Principles of the Management of Metastatic Disease

Approximately 50% to 60% of patients diagnosed with colorectal cancer will develop colorectal metastases,³⁷⁷⁻³⁷⁹ and 80% to 90% of these patients have unresectable metastatic liver disease.^{378,380-383} Metastatic disease most frequently develops metachronously after treatment for



locoregional colorectal cancer, with the liver as the most common site of involvement.³⁸⁴ However, 20% to 34% of patients with colorectal cancer present with synchronous liver metastases.^{383,385} Some evidence indicates that synchronous metastatic colorectal liver disease is associated with a more disseminated disease state and a worse prognosis than metastatic colorectal liver disease that develops metachronously. In a retrospective study of 155 patients who underwent hepatic resection for colorectal liver metastases, patients with synchronous liver metastases had more sites of liver involvement ($P = .008$) and more bilobar metastases ($P = .016$) than patients diagnosed with metachronous liver metastases.³⁸⁶

It has been estimated that more than half of patients who die of colorectal cancer have liver metastases at autopsy, with metastatic liver disease as the cause of death in most patients.³⁸⁷ Reviews of autopsy reports of patients who died from colorectal cancer showed that the liver was the only site of metastatic disease in one-third of patients.³⁸²

Furthermore, several studies have shown rates of 5-year survival to be low in patients with metastatic liver disease not undergoing surgery.^{378,388} Certain clinicopathologic factors, such as the presence of extrahepatic metastases, the presence of more than 3 tumors, and a disease-free interval of fewer than 12 months, have been associated with a poor prognosis in patients with colorectal cancer.^{385,389-393}

Other groups, including ESMO, have established guidelines for the treatment of metastatic colorectal cancer.³⁹⁴ The NCCN recommendations are discussed below.

Surgical Management of Colorectal Metastases

Studies of selected patients undergoing surgery to remove colorectal liver metastases have shown that cure is possible in this population and should be the goal for a substantial number of these patients.^{378,395}

Reports have shown 5-year DFS rates of approximately 20% in patients who have undergone resection of liver metastases,^{390,393} and a recent meta-analysis reported a median 5-year survival of 38%.³⁹⁶ In addition, retrospective analyses and meta-analyses have shown that patients with solitary liver metastases have a 5-year OS rate as high as 71% following resection.³⁹⁷⁻³⁹⁹ Therefore, decisions relating to patient suitability, or potential suitability, and subsequent selection for metastatic colorectal surgery are critical junctures in the management of metastatic colorectal liver disease⁴⁰⁰ (discussed further in *Determining Resectability*).

Colorectal metastatic disease sometimes occurs in the lung.³⁷⁷ Most of the treatment recommendations discussed for metastatic colorectal liver disease also apply to the treatment of colorectal pulmonary metastases.^{177,401,402} Combined pulmonary and hepatic resections of resectable metastatic disease have been performed in very highly selected cases.⁴⁰²⁻⁴⁰⁷

Evidence supporting resection of extrahepatic metastases in patients with metastatic colorectal cancer is limited. In a recent retrospective analysis of patients undergoing concurrent complete resection of hepatic and extrahepatic disease, the 5-year survival rate was lower than in patients without extrahepatic disease, and virtually all patients who underwent resection of extrahepatic metastases experienced disease recurrence.^{408,409} However, a recent international analysis of 1629 patients with colorectal liver metastases showed that 16% of the 171 patients (10.4%) who underwent concurrent resection of extrahepatic and hepatic disease remained disease-free at a median follow-up of 26 months, suggesting that concurrent resection may be of significant benefit in well-selected patients (ie, those with a smaller total number of metastases).⁴⁰⁷ A recent systematic review concluded



similarly that carefully selected patients might benefit from this approach.⁴¹⁰

Recent data suggest that a surgical approach to the treatment of recurrent hepatic disease isolated to the liver can be safely undertaken. However, in a retrospective analysis, 5-year survival was shown to decrease with each subsequent curative-intent surgery, and the presence of extrahepatic disease at the time of surgery was independently associated with a poor prognosis.⁴¹¹⁻⁴¹⁴ In a more recent retrospective analysis of 43 patients who underwent repeat hepatectomy for recurrent disease, 5-year overall and PFS rates were reported to be 73% and 22%, respectively.⁴¹¹ A recent meta-analysis of 27 studies including >7200 patients found that those with longer disease-free intervals; those whose recurrences were solitary, smaller, or unilobular; and those lacking extrahepatic disease derived more benefit from repeat hepatectomy.⁴¹⁵ Panel consensus is that re-resection of liver or lung metastases can be considered in carefully selected patients.^{402,416,417}

Patients with a resectable primary rectal tumor and resectable synchronous metastases can be treated with a staged or simultaneous resection, as discussed below in *Recommendations for Treatment of Resectable Synchronous Metastases*. For patients presenting with unresectable metastases and an intact primary that is not acutely obstructed, palliative resection of the primary is rarely indicated, and systemic chemotherapy is the preferred initial maneuver (discussed in more detail below in *Recommendations for Treatment of Unresectable Synchronous Metastases*).⁴¹⁸

Local Therapies for Metastases

The standard of care for patients with resectable metastatic disease is surgical resection. If resection is not feasible, image-guided ablation⁴¹⁹⁻

⁴²¹ or stereotactic body radiation therapy (SBRT; also called stereotactic ablative radiotherapy [SABR])^{381,422,423} are reasonable options, as discussed in subsequent paragraphs. Many patients, however, are not surgical candidates and/or have disease that cannot be ablated with clear margins⁴²⁰ or safely treated by SBRT. In select patients with liver-only or liver-dominant metastatic disease that cannot be resected or ablated arterially, other locally directed treatment options may be offered.⁴²⁴⁻⁴²⁶

A meta-analysis of 90 studies concluded that hepatic arterial infusion (HAI), radioembolization, and transcatheter arterial chemoembolization (TACE) have similar efficacy in patients with unresectable colorectal metastases in the liver.⁴²⁷ Local therapies are described in more detail below. The role of non-extirpative local therapies in the treatment of colorectal metastases remains controversial.

Hepatic Arterial Infusion

Placement of a hepatic arterial port or implantable pump during surgical intervention for liver resection with subsequent infusion of chemotherapy directed to the liver metastases through the hepatic artery (ie, HAI) is an option (category 2B). In a randomized study of patients who had undergone hepatic resection, administration of floxuridine with dexamethasone through HAI and intravenous 5-FU with or without leucovorin was shown to be superior to a similar systemic chemotherapy regimen alone with respect to 2-year survival free of hepatic disease.^{382,428} The study was not powered for long-term survival, but a trend (not significant) was seen toward better long-term outcome in the group receiving HAI at later follow-up periods.^{382,429} Several other clinical trials have shown significant improvement in response or time to hepatic disease progression when HAI therapy has been compared with systemic chemotherapy, although most have not shown a survival benefit of HAI therapy.³⁸² Results of some studies also suggest that HAI



may be useful in the conversion of patients from an unresectable to a resectable status.^{430,431}

Some of the uncertainties regarding patient selection for preoperative chemotherapy are also relevant to the application of HAI.³⁹⁵ Limitations on the use of HAI therapy include the potential for biliary toxicity³⁸² and the requirement of specific technical expertise. Panel consensus is that HAI therapy should be considered selectively, and only at institutions with extensive experience in both the surgical and medical oncologic aspects of the procedure.

Arterially Directed Embolic Therapy

TACE involves hepatic artery catheterization to cause vessel occlusion with locally delivered chemotherapy.⁴²⁵ A recent randomized trial using HAI to deliver irinotecan-loaded drug-eluting beads (DEBIRI) reported an OS benefit (22 months vs. 15 months; $P = .031$).⁴³² A 2013 meta-analysis identified 5 observational studies and 1 randomized trial and concluded that, although DEBIRI appears to be safe and effective for patients with unresectable colorectal liver metastases, additional trials are needed.⁴³³ A more recent trial randomized 30 patients with colorectal liver metastases to FOLFOX/bevacizumab and 30 patients to FOLFOX/bevacizumab/DEBIRI.⁴³⁴ DEBIRI resulted in an improvement in the primary outcome measure of response rate (78% vs. 54% at 2 months; $P = .02$).

Doxorubicin-eluting beads have also been studied; the strongest data supporting their effectiveness come from several phase II trials in hepatocellular carcinoma.⁴³⁵⁻⁴⁴⁰ A recent systematic review concluded that data are not strong enough to recommend TACE for the treatment of colorectal liver metastases except as part of a clinical trial.⁴⁴¹

The panel believes that arterially directed catheter therapy and, in particular, yttrium 90 microsphere selective internal radiation (see *Radioembolization*, below) is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases.

Liver- or Lung-Directed Radiation

Local radiation therapies include arterial radioembolization with microspheres⁴⁴²⁻⁴⁵² and conformal (stereotactic) external beam RT (EBRT).⁴⁵³

EBRT to the metastatic site can be considered in highly selected cases in which the patient has a limited number of liver or lung metastases or the patient is symptomatic or in the setting of a clinical trial. It should be delivered in a highly conformal manner and should not be used in place of surgical resection. The possible techniques include three-dimensional conformal radiotherapy, SBRT,^{381,422,423,454,455} and IMRT, which uses computer imaging to focus radiation to the tumor site and potentially decrease toxicity to normal tissue.⁴⁵⁶⁻⁴⁵⁹

Radioembolization

A prospective, randomized, phase III trial of 44 patients showed that radioembolization combined with chemotherapy can lengthen time to progression in patients with liver-limited metastatic colorectal cancer following progression on initial therapy (2.1 vs. 4.5 months; $P = .03$).⁴⁶⁰ The effect on the primary endpoint of time to liver progression was more pronounced (2.1 vs. 5.5 months; $P = .003$). Treatment of liver metastases with yttrium-90 glass radioembolization in a prospective, multicenter, phase II study resulted in a median PFS of 2.9 months for patients with colorectal primaries who were refractory to standard treatment.⁴⁶¹ In the refractory setting, a CEA level ≥ 90 and LVI at the time of primary resection were negative prognostic factors for OS.⁴⁵¹



Several large case series have been reported for yttrium-90 radioembolization in patients with refractory unresectable colorectal liver metastases, and the technique appears to be safe with some clinical benefit.^{444,462,463}

Results from the phase III randomized controlled SIRFLOX trial (yttrium-90 resin microspheres with FOLFOX+/- bevacizumab vs. FOLFOX+/- bevacizumab) have been published.⁴⁶⁴ The trial assessed the safety and efficacy of yttrium-90 radioembolization as first-line therapy in 530 patients with colorectal liver metastases. Although the primary endpoint was not met, with PFS in the FOLFOX +/- bevacizumab arm at 10.2 months versus 10.7 months in the FOLFOX/Y-90 arm (HR, 0.93; 95% CI, 0.77–1.12; $P = .43$), a prolonged liver PFS was demonstrated for the study arm (20.5 months for the FOLFOX/Y90 arm vs. 12.6 months for the chemotherapy only arm; HR, 0.69; 95% CI, 0.55–0.90; $P = .002$).

Whereas very little data show any impact on patient survival and the data supporting its efficacy are limited, toxicity with radioembolization is relatively low.⁴⁶⁴⁻⁴⁶⁷ Consensus amongst panel members is that arterially directed catheter therapy and, in particular, yttrium 90 microsphere selective internal radiation is an option in highly selected patients with chemotherapy-resistant/-refractory disease and with predominant hepatic metastases.

Tumor Ablation

Although resection is the standard approach for the local treatment of resectable metastatic disease, patients with liver or lung oligometastases can be considered for tumor ablation therapy.⁴⁶⁸ Ablative techniques include radiofrequency ablation (RFA),^{420,469} microwave ablation, cryoablation, percutaneous ethanol injection, and electro-coagulation. Evidence on the use of RFA as a reasonable treatment option for non-surgical candidates and those with recurrent

disease after hepatectomy with small liver metastases that can be treated with clear margins is growing.^{420,469-472} Data on ablative techniques other than RFA are extremely limited.⁴⁷³⁻⁴⁷⁹

A small number of retrospective studies have compared RFA and resection in the treatment of liver or lung metastases.^{398,479-482} Most of these studies have shown RFA to be inferior to resection in terms of rates of local recurrence and 5-year OS.^{468,483} Whether the differences in outcome observed for patients with liver metastases treated with RFA versus resection alone are from patient selection bias, technological limitations of RFA, or a combination of these factors is currently unclear.⁴⁸¹ A 2010 ASCO clinical evidence review determined that RFA has not been well-studied in the setting of colorectal cancer liver metastases, with no randomized controlled trials having been reported.⁴⁷⁹ The ASCO panel concluded that a compelling need exists for more research in this area. A 2012 Cochrane Database systematic review came to similar conclusions, as have separate meta-analyses.^{475,478,484}

Recently, a trial was reported in which 119 patients were randomized to receive systemic treatment or systemic treatment plus RFA with or without resection.⁴⁸⁵ No difference in OS was seen, but PFS was improved at 3 years in the RFA group (27.6% vs. 10.6%; HR, 0.63; 95% CI, 0.42–0.95; $P = .025$). Similarly 2 recent studies and a position paper by a panel of experts on ablation indicated that ablation may provide acceptable oncologic outcomes for selected patients with small liver metastases that can be ablated with sufficient margins.⁴¹⁹⁻⁴²¹

Resection or ablation (either alone or in combination with resection) should be reserved for patients with metastatic disease that is completely amenable to local therapy with adequate margins. Use of surgery, ablation, or the combination, with the goal of less-than-

complete resection/ablation of all known sites of disease, is not recommended.

Peritoneal Carcinomatosis

Approximately 17% of patients with metastatic colorectal cancer have peritoneal carcinomatosis, with 2% having the peritoneum as the only site of metastasis.⁹¹ Patients with peritoneal metastases generally have a shorter PFS and OS than those without peritoneal involvement.⁹¹ The goal of treatment for most abdominal/peritoneal metastases is palliative, rather than curative, and primarily consists of systemic therapy (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for Colon Cancer) with palliative surgery or stenting if needed for obstruction or impending obstruction.⁴⁸⁶⁻⁴⁸⁸ If an R0 resection can be achieved, however, surgical resection of isolated peritoneal disease may be considered at experienced centers. The panel cautions that the use of bevacizumab in patients with colon or rectal stents is associated with a possible increased risk of bowel perforation.^{489,490}

Cytoreductive Debulking with Heated Intraperitoneal Chemotherapy

Several surgical series and retrospective analyses have addressed the role of cytoreductive surgery (ie, peritoneal stripping surgery) and perioperative hyperthermic intraperitoneal chemotherapy (HIPEC) for the treatment of peritoneal carcinomatosis without extra-abdominal metastases.⁴⁹¹⁻⁴⁹⁹ In the only randomized controlled trial of this approach, Verwaal et al⁵⁰⁰ randomized 105 patients to receive standard therapy (5-FU/LV with or without palliative surgery) or undergo aggressive cytoreductive surgery and HIPEC with mitomycin C; postoperative 5-FU/LV was given to 33 of 47 patients. OS was 12.6 months in the standard arm and 22.3 months in the HIPEC arm ($P = .032$). However, treatment-related morbidity was high, and the mortality was 8% in the HIPEC group, mostly related to bowel leakage. In addition, long-term survival does not seem to be improved by this

treatment as seen by follow-up results.⁵⁰¹ Importantly, this trial was performed without oxaliplatin, irinotecan, or molecularly targeted agents. Some experts have argued that the OS difference seen might have been much smaller if these agents had been used (ie, the control group would have had better outcomes).⁵⁰²

Other criticisms of the Verwaal trial have been published.⁵⁰² One important point is that the trial included patients with peritoneal carcinomatosis of appendiceal origin, a group that has seen greater benefit with the cytoreductive surgery/HIPEC approach.^{492,503-505} A retrospective, multicenter, cohort study reported overall median survival times of 30 and 77 months for patients with peritoneal carcinomatosis of colorectal origin and appendiceal origin, respectively, treated with HIPEC or with cytoreductive surgery and early postoperative intraperitoneal chemotherapy.⁵⁰⁴ The median OS time for patients with pseudomyxoma peritonei, which arises from mucinous appendiceal carcinomas, was not reached at the time of publication. A recent retrospective international registry study reported 10- and 15-year survival rates of 63% and 59%, respectively, in patients with pseudomyxoma peritonei from mucinous appendiceal carcinomas treated with cytoreductive surgery and HIPEC.⁵⁰⁶ HIPEC was not shown to be associated with improvements in OS in this study, whereas completeness of cytoreduction was. Thus, for patients with pseudomyxoma peritonei, optimal treatment is still unclear.⁵⁰⁷

The individual components of this approach have not been well-studied. In fact, studies in rats have suggested that the hyperthermia component of the treatment is irrelevant.⁵⁰⁸ Results of a retrospective cohort study also suggest that heat may not affect outcomes from the procedure.⁴⁹³ In addition, a randomized trial compared systemic 5-FU/oxaliplatin to cytoreductive surgery and intraperitoneal 5-FU without heat.⁵⁰⁹ Although terminated prematurely because of poor accrual, analysis suggested



that the cytoreductive surgery plus IPEC approach may have been superior to the systemic therapy approach (2-year OS, 54% vs. 38%; $P = .04$) for patients with resectable colorectal peritoneal metastases.

In addition, significant morbidity and mortality are associated with this procedure. A 2006 meta-analysis of 2 randomized controlled trials and 12 other studies reported morbidity rates ranging from 23% to 44% and mortality rates ranging from 0% to 12%.⁴⁹⁹ Furthermore, recurrences after the procedure are very common.⁵¹⁰ Whereas the risks are reportedly decreasing with time (ie, recent studies report 1%–5% mortality rates at centers of excellence^{496,502}), the benefits of the approach have not been definitively shown, and HIPEC remains very controversial.⁵¹¹⁻⁵¹⁴

The panel currently believes that complete cytoreductive surgery and/or intraperitoneal chemotherapy can be considered in experienced centers for selected patients with limited peritoneal metastases for whom R0 resection can be achieved. The panel recognizes the need for randomized clinical trials that will address the risks and benefits associated with each of these modalities.

Determining Resectability

The consensus of the panel is that patients diagnosed with potentially resectable metastatic colorectal cancer should undergo an upfront evaluation by a multidisciplinary team, including surgical consultation (ie, with an experienced hepatic surgeon in cases involving liver metastases) to assess resectability status. The criteria for determining patient suitability for resection of metastatic disease are the likelihood of achieving complete resection of all evident disease with negative surgical margins and maintaining adequate liver reserve.⁵¹⁵⁻⁵¹⁸ When the remnant liver is insufficient in size based on cross-sectional imaging volumetrics, preoperative portal vein embolization of the involved liver

can be done to expand the future liver remnant.⁵¹⁹ It should be noted that size alone is rarely a contraindication to resection of a tumor. Resectability differs fundamentally from endpoints that focus more on palliative measures. Instead, the resectability endpoint is focused on the potential of surgery to cure the disease.⁵²⁰ Resection should not be undertaken unless complete removal of all known tumor is realistically possible (R0 resection), because incomplete resection or debulking (R1/R2 resection) has not been shown to be beneficial.^{379,515}

The role of PET/CT in determining resectability of patients with metastatic colorectal cancer is discussed in *Recommendations for Treatment of Metachronous Metastases*, below.

Conversion to Resectability

The majority of patients diagnosed with metastatic colorectal disease have unresectable disease. However, for those with liver-limited unresectable disease that, because of involvement of critical structures, cannot be resected unless regression is accomplished, preoperative chemotherapy is being increasingly considered in highly selected cases in an attempt to downsize colorectal metastases and convert them to a resectable status. Patients presenting with large numbers of metastatic sites within the liver or lung are unlikely to achieve an R0 resection simply on the basis of a favorable response to chemotherapy, as the probability of complete eradication of a metastatic deposit by chemotherapy alone is low. These patients should be regarded as having unresectable disease not amenable to conversion therapy. In some highly selected cases, however, patients with significant response to conversion chemotherapy can be converted from unresectable to resectable status.⁴⁶⁸

Any active metastatic systemic regimen can be used in an attempt to convert a patient's unresectable status to a resectable status, because

the goal is not specifically to eradicate micrometastatic disease, but rather to obtain the optimal size regression of the visible metastases. An important point to keep in mind is that irinotecan- and oxaliplatin-based chemotherapeutic regimens may cause liver steatohepatitis and sinusoidal liver injury, respectively.⁵²¹⁻⁵²⁵ To limit the development of hepatotoxicity, it is therefore recommended that surgery be performed as soon as possible after the patient becomes resectable. Some of the trials addressing various conversion therapy regimens are discussed below.

In the study of Pozzo et al, it was reported that chemotherapy with irinotecan combined with 5-FU/LV enabled a significant portion (32.5%) of the patients with initially unresectable liver metastases to undergo liver resection.⁵¹⁷ The median time to progression was 14.3 months, with all of these patients alive at a median follow-up of 19 months. In a phase II study conducted by the NCCTG,³⁸⁰ 42 patients with unresectable liver metastases were treated with FOLFOX. Twenty-five patients (60%) had tumor reduction and 17 patients (40%; 68% of the responders) were able to undergo resection after a median period of 6 months of chemotherapy. In another study, 1104 patients with initially unresectable colorectal liver disease were treated with chemotherapy, which included oxaliplatin in the majority of cases, and 138 patients (12.5%) classified as “good responders” underwent secondary hepatic resection.³⁸⁹ The 5-year DFS rate for these 138 patients was 22%. In addition, results from a retrospective analysis of 795 previously untreated patients with metastatic colorectal cancer enrolled in the Intergroup N9741 randomized phase III trial evaluating the efficacy of mostly oxaliplatin-containing chemotherapy regimens indicated that 24 patients (3.3%; 2 of the 24 had lung metastases) were able to undergo curative resection after treatment.⁵²⁶ The median OS time in this group was 42.4 months.

In addition, first-line FOLFOXIRI (infusional 5-FU, LV, oxaliplatin, irinotecan) has been compared with FOLFIRI (infusional 5-FU, LV, irinotecan) in 2 randomized clinical trials in patients with unresectable disease.^{527,528} In both studies, FOLFOXIRI led to an increase in R0 secondary resection rates: 6% versus 15%, $P = .033$ in the Gruppo Oncologico Nord Ovest (GONO) trial⁵²⁷; and 4% versus 10%, $P = .08$ in the Gastrointestinal Committee of the Hellenic Oncology Research Group (HORG) trial.⁵²⁸ In a follow-up study of the GONO trial, the 5-year survival rate was higher in the group receiving FOLFOXIRI (15% vs. 8%), with a median OS of 23.4 vs. 16.7 months ($P = .026$).⁵²⁹

More recent favorable results of randomized clinical trials evaluating FOLFIRI or FOLFOX for the purpose of conversion of unresectable disease to resectable disease in combination with anti-epidermal growth factor receptor (EGFR) inhibitors have been reported.⁵³⁰ For instance, in the CELIM phase II trial, patients were randomized to receive cetuximab with either FOLFOX6 or FOLFIRI.⁵³⁰ Retrospective analysis showed that, in both treatment arms combined, resectability increased from 32% to 60% after chemotherapy in patients with wild-type *KRAS* exon 2 ($P < .0001$) with the addition of cetuximab. Another recent randomized controlled trial compared chemotherapy (mFOLFOX6 or FOLFIRI) plus cetuximab to chemotherapy alone in patients with unresectable colorectal cancer metastatic to the liver.⁵³¹ The primary endpoint was the rate of conversion to resectability based on evaluation by a multidisciplinary team. After evaluation, 20 of 70 patients (29%) in the cetuximab arm and 9 of 68 patients (13%) in the control arm were determined to be eligible for curative-intent hepatic resection. R0 resection rates were 25.7% in the cetuximab arm and 7.4% in the control arm ($P < .01$). In addition, surgery improved the median survival time compared to unresected participants in both arms, with longer survival in patients receiving cetuximab (46.4 vs. 25.7 months; $P = .007$



for the cetuximab arm and 36.0 vs. 19.6 months; $P = .016$ for the control arm). A recent meta-analysis of 4 randomized controlled trials concluded that the addition of cetuximab or panitumumab to chemotherapy significantly increased the response rate, the R0 resection rate (from 11%–18%; RR, 1.59; $P = .04$), and PFS, but not OS in patients with wild-type *KRAS* exon 2-containing tumors.⁵³²

The role of bevacizumab in the patient with unresectable, metastatic colorectal cancer, whose disease is felt to be potentially convertible to resectability with a reduction in tumor size, has also been studied. Data seem to suggest that bevacizumab modestly improves the response rate to irinotecan-based regimens.⁵³³ As such, when an irinotecan-based regimen is selected for an attempt to convert unresectable disease to resectability, the use of bevacizumab would seem to be an appropriate consideration. On the other hand, a 1400-patient, randomized, double-blind, placebo-controlled trial of CapeOx or FOLFOX with or without bevacizumab showed no benefit in terms of response rate or tumor regression for the addition of bevacizumab, as measured by both investigators and an independent radiology review committee.⁵³⁴ Therefore, arguments for use of bevacizumab with oxaliplatin-based therapy in this “convert to resectability” setting are not compelling. However, because it is not known in advance whether resectability will be achieved, the use of bevacizumab with oxaliplatin-based therapy in this setting is acceptable.

When initial chemotherapy is planned for patients with unresectable disease that is felt to be potentially convertible to resectability, the panel recommends that a surgical re-evaluation be planned approximately 2 months after initiation of chemotherapy, and that those patients who continue to receive chemotherapy undergo surgical re-evaluation approximately every 2 months thereafter.^{525,535-537} Reported risks associated with chemotherapy include the potential for development of

liver steatosis or steatohepatitis when oxaliplatin or irinotecan-containing chemotherapeutic regimens are administered.⁵²¹ To limit the development of hepatotoxicity, it is therefore recommended that surgery should be performed as soon as possible after the patient becomes resectable.

Neoadjuvant and Adjuvant Therapy for Resectable Metastatic Disease

The panel recommends consideration of administration of a course of an active systemic therapy regimen for metastatic disease, for a total perioperative treatment time of approximately 6 months, for most patients undergoing liver or lung resection, to increase the likelihood that residual microscopic disease will be eradicated (category 2B for the use of biologic agents in the perioperative metastatic setting). A recent meta-analysis identified 3 randomized clinical trials comparing surgery alone to surgery plus systemic therapy with 642 evaluable patients with colorectal liver metastases.⁵³⁸ The pooled analysis showed a benefit of chemotherapy in PFS (pooled HR, 0.75; CI, 0.62–0.91; $P = .003$) and DFS (pooled HR, 0.71; CI, 0.58–0.88; $P = .001$), but not in OS (pooled HR, 0.74; CI, 0.53–1.05; $P = .088$). Another meta-analysis published in 2015 combined data on 1896 patients from 10 studies and also found that perioperative chemotherapy improved DFS (HR, 0.81; 95% CI, 0.72–0.91; $P = .0007$) but not OS (HR, 0.88; 95% CI, 0.77–1.01; $P = .07$) in patients with resectable colorectal liver metastases.⁵³⁹ Additional recent meta-analyses have also failed to observe an OS benefit with the addition of adjuvant chemotherapy in resectable metastatic colorectal cancer.^{540,541}

The choice of chemotherapy regimen in the preoperative setting is dependent on a number of factors, including the chemotherapy history of the patient, whether disease is synchronous or metachronous, and the response rates and safety/toxicity issues associated with the



regimens, as outlined in the guidelines. Biologics are not recommended in the perioperative metastatic setting, with the exception of initial therapy in unresectable patients who may be converted to a resectable state.

Although the benefits of perioperative chemotherapy for patients with liver metastases have not yet been fully validated in randomized clinical trials, a recent EORTC phase III study (EORTC 40983) evaluating use of perioperative FOLFOX (6 cycles before and 6 cycles after surgery) for patients with initially resectable liver metastases demonstrated absolute improvements in 3-year PFS of 8.1% ($P = .041$) and 9.2% ($P = .025$) for all eligible patients and all resected patients, respectively, when chemotherapy in conjunction with surgery was compared with surgery alone.⁵⁴² The partial response rate after preoperative FOLFOX was 40%, and operative mortality was <1% in both treatment groups. However, no difference in OS was seen between the groups, perhaps because second-line therapy was given to 77% of the patients in the surgery only arm and to 59% of the patients in the chemotherapy arm.⁵⁴³

The optimal sequencing of chemotherapy remains unclear. Patients with initially resectable disease may undergo liver resection first, followed by postoperative adjuvant chemotherapy. Alternatively, perioperative (neoadjuvant plus postoperative) chemotherapy can be used.^{540,544}

Potential advantages of the preoperative chemotherapy approach include earlier treatment of micrometastatic disease; determination of responsiveness to chemotherapy, which can be prognostic and help plan postoperative therapy; and avoidance of local therapy in those who progress early. Potential disadvantages include missing the “window of opportunity” for resection because of the possibility of disease

progression or achievement of a complete response, thereby making it difficult to identify areas for resection.^{382,545,546} Importantly, results from a study of patients with colorectal cancer receiving preoperative chemotherapy indicated that viable cancer was still present in most of the original sites of metastases when these sites were examined pathologically despite achievement of a complete response as evaluated on CT scan.^{546,547} It is therefore essential that during treatment with preoperative chemotherapy, frequent evaluations are undertaken and close communication is maintained between medical oncologists, radiologists, surgeons, and patients so that a treatment strategy can be developed that optimizes exposure to the preoperative regimen and facilitates an appropriately timed surgical intervention.⁵²¹

Other reported risks associated with the preoperative chemotherapy approach include the potential for development of liver steatohepatitis and sinusoidal liver injury when irinotecan- and oxaliplatin-based chemotherapeutic regimens are administered, respectively.⁵²¹⁻⁵²⁵ To reduce the development of hepatotoxicity, the neoadjuvant period is usually limited to 2 to 3 months, and patients should be carefully monitored by a multidisciplinary team.

It is important to note that some of the treatment approaches for patients diagnosed with rectal cancer and resectable synchronous lung or liver metastases differ relative to those for patients diagnosed with similarly staged colon cancer. In particular, initial treatment options for synchronous resectable rectal cancer include preoperative chemoRT directed toward treatment of the primary cancer; a preoperative combination chemotherapy regimen plus a biologic agent to target metastatic disease; and a surgical approach (ie, staged or synchronous resection of metastases and rectal lesion). Advantages of an initial chemoRT approach include a possible decreased risk of pelvic failure following surgery, while a disadvantage is that preoperative pelvic RT



may decrease tolerance to systemic bevacizumab-containing adjuvant regimens, thereby limiting subsequent treatment of systemic disease. Data to guide decisions regarding optimal treatment approaches in this population of patients are very limited.

Based largely on extrapolation from stage III disease and limited randomized data for stage IV disease, the panel recommends the use of postoperative adjuvant chemotherapy in patients who have undergone liver or lung resection and who have received preoperative chemoRT. Postoperative chemoRT is recommended for patients with synchronous metastases who have not received prior chemoRT and who are at higher risk for pelvic recurrence following staged or synchronous resection of metastases and rectal lesion (ie, patients with disease staged as pT3-4, Any N, M1 or Any T, N1-2, M1).

Perioperative Bevacizumab for Resectable Metastatic Disease

The efficacy of bevacizumab in combination with FOLFOX and FOLFIRI in the treatment of unresectable metastatic disease (see *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for Colon Cancer) has led to study of its use in combination with these regimens in the preoperative setting. However, the safety of administering bevacizumab pre- or postoperatively in combination with 5-FU–based regimens has not been adequately evaluated. A retrospective evaluation of data from 2 randomized clinical trials of 1132 patients receiving chemotherapy with or without bevacizumab as initial therapy for metastatic colorectal cancer indicated that the incidence of wound healing complications was increased for the group of patients undergoing a major surgical procedure while receiving a bevacizumab-containing regimen when compared to the group receiving chemotherapy alone while undergoing major surgery (13% vs. 3.4%, respectively; $P = .28$).⁵⁴⁸ However, when chemotherapy plus bevacizumab or chemotherapy alone was administered prior to surgery,

the incidence of wound healing complications in either group of patients was low (1.3% vs. 0.5%; $P = .63$).

A recent meta-analysis of randomized controlled trials demonstrated that the addition of bevacizumab to chemotherapy is associated with a higher incidence of treatment-related mortality than chemotherapy alone (RR, 1.33; 95% CI, 1.02–1.73; $P = .04$); hemorrhage (23.5%), neutropenia (12.2%), and gastrointestinal perforation (7.1%) were the most common causes of fatality.⁵⁴⁹ Venous thromboembolisms, however, were not increased in patients receiving bevacizumab with chemotherapy versus those receiving chemotherapy alone.⁵⁵⁰

The panel recommends against the use of bevacizumab as neoadjuvant treatment of patients with resectable metastatic rectal cancer. For patients who receive bevacizumab for unresectable disease and are converted to a resectable state, the panel recommends at least a 6-week interval (which corresponds to 2 half-lives of the drug⁵⁵¹) between the last dose of bevacizumab and surgery.

Perioperative Cetuximab and Panitumumab for Resectable Metastatic Disease

The New EPOC trial, which was stopped early because it met protocol-defined futility criteria, found a lack of benefit to cetuximab with chemotherapy in the perioperative metastatic setting (>85% received FOLFOX or CapeOx; patients with prior oxaliplatin received FOLFIRI).⁵⁵² In fact, with less than half of expected events observed, PFS was significantly reduced in the cetuximab arm (14.8 vs. 24.2 months; HR 1.50; 95% CI, 1.00–2.25; $P < .048$). The panel thus recommends against panitumumab and cetuximab in the neoadjuvant setting. The panel also points out cetuximab and panitumumab should be used with caution in patients with unresectable disease that could potentially be converted to a resectable status.

Systemic Therapy for Advanced or Metastatic Disease

The current management of disseminated metastatic colorectal cancer involves various active drugs, either in combination or as single agents: 5-FU/LV, capecitabine, irinotecan, oxaliplatin, bevacizumab, cetuximab, panitumumab, ziv-aflibercept, ramucirumab, regorafenib, trifluridine-tipiracil, pembrolizumab, and nivolumab. The choice of therapy is based on consideration of the goals of therapy, the type and timing of prior therapy, the mutational profile of the tumor, and the differing efficacy and toxicity profiles of the constituent drugs. Although the specific chemotherapy regimens listed in the guidelines are designated according to whether they pertain to initial therapy or therapy after first, second, or third progression, it is important to clarify that these recommendations represent a continuum of care and that these lines of treatment are blurred rather than discrete. For example, if oxaliplatin is administered as a part of an initial treatment regimen but is discontinued after 12 weeks or earlier for escalating neurotoxicity, continuation of the remainder of the treatment regimen would still be considered initial therapy.

Principles to consider at the start of therapy include preplanned strategies for altering therapy for patients exhibiting a tumor response or disease characterized as stable or progressive, and plans for adjusting therapy for patients who experience certain toxicities. For example, decisions related to therapeutic choices after first progression of disease should be based partly on the prior therapies received (ie, exposing the patient to a range of cytotoxic agents). Furthermore, an evaluation of the efficacy and safety of these regimens for an individual patient must take into account not only the component drugs, but also the doses, schedules, and methods of administration of these agents, and the potential for surgical cure and the performance status of the patient.

The continuum of care approach to the management of patients with metastatic rectal cancer is the same as described for patients with metastatic colon cancer. Please refer to *Systemic Therapy for Advanced or Metastatic Disease* in the NCCN Guidelines for Colon Cancer for a detailed discussion of the various options for systemic treatment (available at www.NCCN.org). The roles of predictive markers for treatment selection in the advanced and metastatic disease setting are discussed below.

Predictive Markers for Cetuximab and Panitumumab *Primary Tumor Sidedness*

A growing body of data has shown that the location of the primary tumor can be both prognostic and predictive of response to EGFR inhibitors in metastatic colorectal cancer.⁵⁵³⁻⁵⁶⁰ For example, outcomes of 75 patients with metastatic colorectal cancer treated with cetuximab, panitumumab, or cetuximab/irinotecan in first-line or subsequent lines of therapy at 3 Italian centers were analyzed based on sidedness of the primary tumor.⁵⁵⁴ No responses were seen in the patients with right-sided primary tumors compared with a response rate of 41% in those with left-sided primaries ($P = .003$). The median PFS was 2.3 and 6.6 months in patients with right-sided and left-sided tumors, respectively (HR, 3.97; 95% CI, 2.09–7.53; $P < .0001$).

The strongest evidence for the predictive value of primary tumor sidedness and response to EGFR inhibitors is in the first-line treatment of patients in the phase III CALGB/SWOG 80405 trial.^{560,561} The study showed that patients with all *RAS* wild-type, right-sided primary tumors (cecum to hepatic flexure) had longer OS if treated with bevacizumab than if treated with cetuximab in first line (HR, 1.36; 95% CI, 0.93–1.99; $P = .10$), whereas patients with all *RAS* wild-type, left-sided primary tumors (splenic flexure to rectum) had longer OS if treated with cetuximab than if treated with bevacizumab (HR, 0.77; 95% CI, 0.59–



0.99; $P = 0.04$).⁵⁶¹ OS was prolonged with cetuximab versus bevacizumab in the left-sided primary group (39.3 months vs. 32.6 months), but was shortened in the right-sided primary group (13.6 months vs. 29.2 months).

These and other data suggest that cetuximab and panitumumab confer little if any benefit to patients with metastatic colorectal cancer if the primary tumor originated on the right side.^{553,554,556,557} The panel believes that primary tumor sidedness is a surrogate for the non-random distribution of molecular subtypes across the colon and that the ongoing analysis of tumor specimens from the study will enable a better understanding of the biologic explanation of the observed difference in response to EGFR inhibitors. Until that time, only patients whose primary tumors originated on the left side of the colon (splenic flexure to rectum) should be offered cetuximab or panitumumab in the first-line treatment of metastatic disease. Evidence also suggests that sidedness is predictive of response to EGFR inhibitors in subsequent lines of therapy,^{553,554,557} but the panel awaits more definitive studies. Until such data are available, all patients with *RAS* wild-type tumors can be considered for panitumumab or cetuximab in subsequent lines of therapy if neither was previously given.

KRAS, NRAS, and BRAF Status

EGFR has been shown to be overexpressed in 49% to 82% of colorectal tumors.⁵⁶²⁻⁵⁶⁵ EGFR testing of colorectal tumor cells has no proven predictive value in determining likelihood of response to either cetuximab or panitumumab. Data from the BOND-1 study indicated that the intensity of immunohistochemical staining of EGFR in colorectal tumor cells did not correlate with the response rate to cetuximab.⁵⁶⁶ A similar conclusion was drawn with respect to panitumumab.⁵⁶⁷

Therefore, routine EGFR testing is not recommended, and no patient

should be considered for or excluded from cetuximab or panitumumab therapy based on EGFR test results.

Cetuximab and panitumumab are monoclonal antibodies directed against EGFR that inhibit its downstream signaling pathways, but EGFR status as assessed using immunohistochemistry is not predictive of treatment efficacy.^{566,568} Furthermore, cetuximab and panitumumab are only effective in approximately 10% to 20% of patients with colorectal cancer.^{566,568,569} The *RAS/RAF/MAPK* pathway is downstream of EGFR; mutations in components of this pathway are being studied in search of predictive markers for efficacy of these therapies.

A sizable body of literature has shown that these *KRAS* exon 2 mutations are predictive of response to cetuximab or panitumumab therapy.⁵⁷⁰⁻⁵⁷⁹ More recent evidence shows that mutations in *KRAS* outside of exon 2 and mutations in *NRAS* are also predictive for a lack of benefit to cetuximab and panitumumab (see *NRAS and Other KRAS Mutations, below*).^{580,581} The panel therefore strongly recommends *KRAS/NRAS* genotyping of tumor tissue (either primary tumor or metastasis) in all patients with metastatic colorectal cancer. Patients with known *KRAS* or *NRAS* mutations should not be treated with either cetuximab or panitumumab, either alone or in combination with other anticancer agents, because they have virtually no chance of benefit and the exposure to toxicity and expense cannot be justified. It is implied throughout the guidelines that NCCN recommendations involving cetuximab or panitumumab relate only to patients with disease characterized by *KRAS/NRAS* wild-type genes. ASCO released a Provisional Clinical Opinion Update on extended *RAS* testing in patients with metastatic colorectal cancer that is consistent with the NCCN panel's recommendations.⁵⁸²



The panel strongly recommends genotyping of tumor tissue (either primary tumor or metastasis) in all patients with metastatic colorectal cancer for *RAS* (*KRAS* exon 2 and non-exon 2; *NRAS*) and *BRAF* at diagnosis of stage IV disease. The recommendation for *KRAS/NRAS* testing at this point is not meant to indicate a preference regarding regimen selection in the first-line setting. Rather, this early establishment of *KRAS/NRAS* status is appropriate to plan for the treatment continuum so that the information may be obtained in a non-time-sensitive manner, and the patient and provider can discuss the implications of a *KRAS/NRAS* mutation, if present, while other treatment options still exist. Note that because anti-EGFR agents have no role in the management of stage I, II, or III disease, *KRAS/NRAS* genotyping of colorectal cancers at these earlier stages is not recommended.

KRAS mutations are early events in colorectal cancer formation, and therefore a very tight correlation exists between mutation status in the primary tumor and the metastases.^{583,584} For this reason, *KRAS/NRAS* genotyping can be performed on archived specimens of either the primary tumor or a metastasis. Fresh biopsies should not be obtained solely for the purpose of *KRAS/NRAS* genotyping unless an archived specimen from either the primary tumor or a metastasis is unavailable.

The panel recommends that *KRAS*, *NRAS*, and *BRAF* gene testing be performed only in laboratories that are certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA-88) as qualified to perform highly complex molecular pathology testing.⁵⁸⁵ No specific testing methodology is recommended.⁵⁸⁶

***KRAS* Exon 2 Mutations:** Approximately 40% of colorectal cancers are characterized by mutations in codons 12 and 13 in exon 2 of the coding region of the *KRAS* gene.^{570,587} A sizable body of literature has shown that these *KRAS* exon 2 mutations are predictive of lack of response to

cetuximab or panitumumab therapy,⁵⁷⁰⁻⁵⁷⁹ and FDA labels for cetuximab and panitumumab specifically state that these agents are not recommended for the treatment of colorectal cancer characterized by these mutations.^{588,589} Results are mixed as far as the prognostic value of *KRAS* mutations. In the Alliance N0147 trial, patients with *KRAS* exon 2 mutations experienced a shorter DFS than patients without such mutations.⁵⁹⁰ At this time, however, the test is not recommended for prognostic reasons.

A retrospective study from De Roock et al⁵⁹¹ raised the possibility that codon 13 mutations (G13D) in *KRAS* may not be absolutely predictive of non-response to EGFR inhibition. Another retrospective study showed similar results.⁵⁹² Furthermore, a more recent retrospective analysis of 3 randomized controlled phase III trials concluded that patients with *KRAS* G13D mutations were unlikely to respond to panitumumab.⁵⁹³ Results from a prospective phase II single-arm trial assessed the benefit of cetuximab monotherapy in 12 patients with refractory metastatic colorectal cancer whose tumors contained *KRAS* G13D mutations.⁵⁹⁴ The primary endpoint of 4-month progression-free rate was not met (25%), and no responses were seen. Preliminary results of the AGITG phase II ICECREAM trial also failed to see a benefit of cetuximab monotherapy in patients with *KRAS* G13D mutations.⁵⁹⁵ However, partial responses were reported after treatment with irinotecan plus cetuximab in 9% of this irinotecan-refractory population. The panel believes that patients with any known *KRAS* mutation, including G13D, should not be treated with cetuximab or panitumumab.

***NRAS* and Other *KRAS* Mutations:** In the AGITG MAX study, 10% of patients with wild-type *KRAS* exon 2 had mutations in *KRAS* exons 3 or 4 or in *NRAS* exons 2, 3, and 4.⁵⁹⁶ In the PRIME trial, 17% of 641 patients without *KRAS* exon 2 mutations were found to have mutations



in exons 3 and 4 of *KRAS* or mutations in exons 2, 3, and 4 of *NRAS*. A predefined retrospective subset analysis of data from PRIME revealed that PFS (HR, 1.31; 95% CI, 1.07–1.60; $P = .008$) and OS (HR, 1.21; 95% CI, 1.01–1.45; $P = .04$) were decreased in patients with any *KRAS* or *NRAS* mutation who received panitumumab plus FOLFOX compared to those who received FOLFOX alone.⁵⁸⁰ These results show that panitumumab does not benefit patients with *KRAS* or *NRAS* mutations and may even have a detrimental effect in these patients.

Updated analysis of the FIRE-3 trial was recently published.⁵⁹⁷ When all *RAS* (*KRAS*/*NRAS*) mutations were considered, PFS was significantly worse in patients with *RAS*-mutant tumors receiving FOLFIRI plus cetuximab than patients with *RAS*-mutant tumors receiving FOLFIRI plus bevacizumab (6.1 months vs. 12.2 months; $P = .004$). On the other hand, patients with *KRAS*/*NRAS* wild-type tumors showed no difference in PFS between the regimens (10.4 months vs. 10.2 months; $P = .54$). This result indicates that cetuximab likely has a detrimental effect in patients with *KRAS* or *NRAS* mutations.

The FDA indication for panitumumab was updated to state that panitumumab is not indicated for the treatment of patients with *KRAS* or *NRAS* mutation-positive disease in combination with oxaliplatin-based chemotherapy.⁵⁸⁸ The NCCN Colon/Rectal Cancer Panel believes that non-exon 2 *KRAS* mutation status and *NRAS* mutation status should be determined at diagnosis of stage IV disease. Patients with any known *KRAS* mutation (exon 2 or non-exon 2) or *NRAS* mutation should not be treated with either cetuximab or panitumumab.

***BRAF* V600E Mutations:** Although mutations of *KRAS*/*NRAS* indicate a lack of response to EGFR inhibitors, many tumors containing wild-type *KRAS*/*NRAS* still do not respond to these therapies. Therefore, studies have addressed factors downstream of *KRAS*/*NRAS* as

possible additional biomarkers predictive of response to cetuximab or panitumumab. Approximately 5% to 9% of colorectal cancers are characterized by a specific mutation in the *BRAF* gene (V600E).^{598,599} *BRAF* mutations are, for all practical purposes, limited to tumors that do not have *KRAS* exon 2 mutations.^{598,600} Activation of the protein product of the non-mutated *BRAF* gene occurs downstream of the activated *KRAS* protein in the EGFR pathway. The mutated *BRAF* protein product is believed to be constitutively active,⁶⁰¹⁻⁶⁰³ thereby putatively bypassing inhibition of EGFR by cetuximab or panitumumab.

Limited data from unplanned retrospective subset analyses of patients with metastatic colorectal cancer treated in the first-line setting suggest that although a *BRAF* V600E mutation confers a poor prognosis regardless of treatment, patients with disease characterized by this mutation may receive some benefit from the addition of cetuximab to front-line therapy.^{604,605} On the other hand, results from the randomized phase III Medical Research Council (MRC) COIN trial suggest that cetuximab may have no effect or even a detrimental effect in patients with *BRAF*-mutated tumors treated with CapeOx or FOLFOX in the first-line setting.⁶⁰⁰

In subsequent lines of therapy, retrospective evidence suggests that mutated *BRAF* is a marker of resistance to anti-EGFR therapy in the non-first-line setting of metastatic disease.⁶⁰⁶⁻⁶⁰⁸ A retrospective study of 773 primary tumor samples from patients with chemotherapy-refractory disease showed that *BRAF* mutations conferred a significantly lower response rate to cetuximab (2/24; 8.3%) compared with tumors with wild-type *BRAF* (124/326; 38.0%; $P = .0012$).⁶⁰⁹ Furthermore, data from the multicenter randomized controlled PICCOLO trial are consistent with this conclusion, with a suggestion of harm seen for the addition of panitumumab to irinotecan in the non-first-line setting in the small subset of patients with *BRAF* mutations.⁶¹⁰



A meta-analysis published in 2015 identified 9 phase III trials and 1 phase II trial that compared cetuximab or panitumumab with standard therapy or best supportive care including 463 patients with metastatic colorectal tumors with *BRAF* mutations (first-line, second-line, or refractory settings).⁶¹¹ The addition of an EGFR inhibitor did not improve PFS (HR, 0.88; 95% CI, 0.67–1.14; $P = .33$), OS (HR, 0.91; 95% CI, 0.62–1.34; $P = .63$), or overall response rate (ORR) (RR, 1.31; 95% CI, 0.83–2.08, $P = .25$) compared with control arms. Similarly, another meta-analysis identified 7 randomized controlled trials and found that cetuximab and panitumumab did not improve PFS (HR, 0.86; 95% CI, 0.61–1.21) or OS (HR, 0.97; 95% CI, 0.67–1.41) in patients with *BRAF* mutations.⁶¹²

Despite uncertainty over its role as a predictive marker, it is clear that mutations in *BRAF* are a strong prognostic marker.^{587,600,605,613-616} A recent prospective analysis of tissues from patients with stage II and III colon cancer enrolled in the PETACC-3 trial showed that the *BRAF* mutation is prognostic for OS in patients with low levels of MSI (MSI-L) or stable microsatellites (MSS) (HR, 2.2; 95% CI, 1.4–3.4; $P = .0003$).⁵⁸⁷ Moreover, an updated analysis of the CRYSTAL trial showed that patients with metastatic colorectal tumors carrying a *BRAF* mutation have a worse prognosis than those with the wild-type gene.⁶⁰⁵ Additionally, *BRAF* mutation status predicted OS in the AGITG MAX trial, with an HR of 0.49 (CI, 0.33–0.73; $P = .001$).⁶¹⁴ The OS in patients with *BRAF* mutations in the COIN trial was 8.8 months, while those with *KRAS* exon 2 mutations and wild-type *KRAS* exon 2 tumors had OS times of 14.4 months and 20.1 months, respectively.⁶⁰⁰ Results from a recent systematic review and meta-analysis of 21 studies, including 9885 patients, suggest that *BRAF* mutation may accompany specific high-risk clinicopathologic characteristics.⁶¹⁷ In particular, an association was observed between *BRAF* mutation and proximal tumor location

(OR, 5.22; 95% CI, 3.80–7.17; $P < .001$), T4 tumors (OR, 1.76; 95% CI, 1.16–2.66; $P = .007$), and poor differentiation (OR, 3.82, 95% CI, 2.71–5.36; $P < .001$).

Overall, the panel believes that evidence increasingly suggests that *BRAF* V600E mutation makes response to panitumumab or cetuximab, as single agents or in combination with cytotoxic chemotherapy, highly unlikely. The panel recommends *BRAF* genotyping of tumor tissue (either primary tumor or metastasis⁶¹⁸) at diagnosis of stage IV disease. Testing for the *BRAF* V600E mutation can be performed on formalin-fixed paraffin-embedded tissues and is usually performed by PCR amplification and direct DNA sequence analysis. Allele-specific PCR is another acceptable method for detecting this mutation.

HER2 Overexpression

HER2 is a member of the same family of signaling kinase receptors as EGFR and has been successfully targeted in breast cancer in both the advanced and adjuvant settings. *HER2* is rarely overexpressed in colorectal cancer (approximately 3% overall), but the prevalence is higher in *RAS/BRAF*–wild type tumors (reported at 5%–14%).^{619,620} Specific molecular diagnostic methods have been proposed for *HER2* testing in colorectal cancer,⁶²¹ and various therapeutic approaches are being tested in patients with tumors that have *HER2* overexpression (eg, trastuzumab plus lapatinib, trastuzumab plus pertuzumab).^{620,622} These approaches are currently considered investigational, and enrollment in a clinical trial is encouraged.

Evidence does not support a prognostic role of *HER2* overexpression.⁶²³ However, initial results indicate *HER2* overexpression may be predictive of resistance to EGFR-targeting monoclonal antibodies.^{619,624} For example, in a cohort of 97 patients with *RAS/BRAF*–wild type metastatic colorectal cancer, median PFS on first-line therapy without an EGFR

inhibitor was similar regardless of HER2 status.⁶¹⁹ However, in second-line therapy with an EGFR inhibitor, the PFS was significantly shorter in those with HER2 amplification compared with those without HER2 amplification (2.9 months vs. 8.1 months; HR, 5.0; $P < .0001$). Larger confirmatory studies are needed, and the panel does not recommend HER2 testing for prognostication or treatment planning at this time.

Predictive Marker for Pembrolizumab and Nivolumab: MSI Status

The percentage of stage IV colorectal tumors characterized as MSI-H (mismatch repair-deficient; dMMR) ranges from 3.5% to 5.0% in clinical trials and was 6.5% in the Nurses' Health Study and Health Professionals Follow-up Study.⁶²⁵⁻⁶²⁷ dMMR tumors contain thousands of mutations, which can encode mutant proteins with the potential to be recognized and targeted by the immune system. However, programmed death-ligands PD-L1 and PD-L2 on tumor cells can suppress the immune response by binding to programmed death receptor-1 (PD-1) on T-effector cells. This system evolved to protect the host from an unchecked immune response. Many tumors upregulate PD-L1 and thus evade the immune system.⁶²⁸ It has therefore been hypothesized that dMMR tumors may be sensitive to PD-1 inhibitors.

Pembrolizumab is a humanized, IgG4 monoclonal antibody that binds to PD-1 with high affinity, preventing its interaction with PD-L1 and PD-L2 and thus allowing immune recognition and response.⁶²⁹ Pembrolizumab is FDA-approved for the treatment of patients with unresectable or metastatic melanoma following disease progression on other standard therapies and has shown activity against non-small cell lung cancer.^{629,630}

A recent phase II study evaluated the activity of pembrolizumab in 11 patients with dMMR colorectal cancer, 21 patients with MMR-proficient colorectal cancer, and 9 patients with dMMR non-colorectal

carcinomas.⁶³¹ All patients had progressive metastatic disease; the patients in the colorectal arms had progressed through 2 to 4 previous therapies. The primary endpoints were the immune-related objective response rate and the 20-week immune-related PFS rate. The immune-related objective response rates were 40% (95% CI, 12%–74%) in the dMMR colorectal cancer group, 0% (95% CI, 0%–20%) in the MMR-proficient colorectal cancer group, and 71% (95% CI, 29%–96%) in the dMMR non-colorectal group. The 20-week immune-related PFS rates were 78% (95% CI, 40–97), 11% (95% CI, 1–35), and 67% (95% CI, 22–96), respectively. These results indicate that MSI is a predictive marker for the effectiveness of pembrolizumab across tumor types. Furthermore, the median PFS and OS were not reached in the arm with dMMR colorectal cancer and were 2.2 and 5.0 months, respectively, in the MMR-proficient colorectal cancer group (HR for disease progression or death, 0.10; $P < .001$).

Nivolumab is another humanized IgG4 PD-1 blocking antibody, with FDA indications in melanoma and non-small cell lung cancer.⁶³²

Nivolumab was studied with or without ipilimumab in patients with metastatic colorectal cancer in a phase II trial.⁶³³ The median PFS was 5.3 months (95% CI, 1.4–not estimable) in the MMR-deficient patients who received nivolumab monotherapy, not reached in the MMR-deficient patients who received nivolumab plus ipilimumab, and 1.4 months (95% CI, 1.2–1.9) in the pooled MMR-proficient group.

Based on these data, the panel recommends pembrolizumab or nivolumab as treatment options in patients with metastatic MMR-deficient colorectal cancer in second- or third-line therapy. Patients progressing on either of these drugs should not be offered the other. Additional clinical trials are ongoing to confirm the benefit of these drugs in this setting.



Although PD-1 immune checkpoint inhibitors are generally well tolerated, serious adverse reactions—many immune-mediated—occur in as many as 21% to 41% of patients.^{631,633,634} The most common immune-mediated side effects are to the skin, liver, kidneys, gastrointestinal tract, lungs, and endocrine systems.⁶³⁵⁻⁶³⁷ Pneumonitis, occurring in approximately 3% to 7% of patients on pembrolizumab or nivolumab, is one of the most serious side effects of PD-1 inhibitors.^{635,638-640}

Recommendations for Treatment of Resectable Synchronous Metastases

As part of the pre-treatment workup, the panel recommends tumor *KRAS/NRAS* gene status testing for all patients with metastatic colorectal cancer at the time of diagnosis of metastatic disease. If *KRAS/NRAS* are found to be wild-type, *BRAF* testing can be considered (see *KRAS, NRAS, and BRAF Status*, above).

When patients present with colorectal cancer and synchronous liver metastases, resection of the primary tumor and liver can be done in a simultaneous or staged approach following neoadjuvant treatment (options discussed below).⁶⁴¹⁻⁶⁴⁸ Historically, in the staged approach, the primary tumor was usually resected first. However, the approach of liver resection before resection of the primary tumor is now well-accepted. In addition, emerging data suggest that chemotherapy, followed by resection of liver metastases before resection of the primary tumor, might be an effective approach in some patients, although more studies are needed.⁶⁴⁹⁻⁶⁵¹ In addition, neoadjuvant short-course radiation of T1-T3 primary rectal tumors is an option in this setting.⁶⁵² Locally ablative procedures can be considered instead of or in addition to resection in cases of liver or lung oligometastases (see *Local Therapies for Metastases*, above), but resection is preferred.

There are several acceptable sequences of therapy in the setting of resectable synchronous disease as outlined in the guidelines. As in other settings, the total perioperative therapy should not exceed 6 months.

In the 2014 version of these guidelines, the panel removed the option of surgery as the initial treatment because it believes that the majority of patients should receive preoperative therapy. Upfront systemic treatment has the goal of early eradication of micrometastases, while the goal of consolidating chemoRT is local control of disease prior to surgery/local therapy. For patients receiving neoadjuvant therapy, surgery/local therapy should be performed 5 to 12 weeks following completion of treatment. The panel acknowledges that some patients may not be candidates for chemotherapy or radiation; clinical judgment should be used in such cases.

Recommendations for Treatment of Unresectable Synchronous Metastases

Patients with unresectable metastases or who are medically inoperable are treated according to whether they are symptomatic or asymptomatic. Symptomatic patients are treated with chemotherapy alone, combined modality therapy with 5-FU/RT or capecitabine/RT, short-course radiation (not recommended for T4 tumors), resection of the involved rectal segment, diverting colostomy, or stenting. Primary treatment should be followed by an active systemic therapy regimen for advanced or metastatic disease.

For patients with asymptomatic liver or lung disease that is deemed to be unresectable, the panel recommends systemic therapy for advanced or metastatic disease to attempt to render these patients candidates for resection (see *Determining Resectability and Conversion to Resectability*, above). Chemotherapy regimens with high response rates



should be considered for patients with potentially convertible disease.⁶⁵³ These patients should be re-evaluated for resection after 2 months of chemotherapy and every 2 months thereafter while undergoing such therapy.

Results from a recent study suggest that there may be some benefit in both OS and PFS from resection of the primary in the setting of unresectable colorectal metastases.⁶⁵⁴ Other retrospective analyses have also shown a potential benefit.^{655,656} However, the prospective, multicenter, phase II NSABP C-10 trial showed that patients with an asymptomatic primary colon tumor and unresectable metastatic disease who received mFOLFOX6 with bevacizumab experienced an acceptable level of morbidity without upfront resection of the primary tumor.⁶⁵⁷ The median OS was 19.9 months. Notably, symptomatic improvement in the primary is often seen with first-line systemic chemotherapy even within the first 1 to 2 weeks. Furthermore, complications from the primary lesion are uncommon in these circumstances,⁴¹⁸ and its removal delays initiation of systemic chemotherapy. In fact, a recent systematic review concluded that resection of the primary does not reduce complications and does not improve OS.⁶⁵⁸ However, a different systematic review concluded that, while data are not strong, resection of the primary tumor may provide a survival benefit.⁶⁵⁹ Another systematic review and meta-analysis identified 5 studies that compared open to laparoscopic palliative colectomies in this setting.⁶⁶⁰ The laparoscopic approach resulted in shorter lengths of hospital stays ($P < .001$), fewer postoperative complications ($P = 0.01$), and lower estimated blood loss ($P < .01$).

Overall, the panel believes that the risks of surgery outweigh the possible benefits of this approach. Routine palliative resection of a synchronous primary lesion should therefore only be considered if the

patient has an unequivocal imminent risk of obstruction or acute significant bleeding.⁴¹⁸

An intact primary tumor is not a contraindication to bevacizumab use. The risk of gastrointestinal perforation in the setting of bevacizumab is not decreased by removal of the primary tumor, as large bowel perforations, in general, and perforation of the primary lesion, in particular, are rare (see *Systemic Therapy for Advanced or Metastatic Disease* in the Discussion section of the NCCN Guidelines for Colon Cancer, available at www.NCCN.org).

Recommendations for Treatment of Metachronous Metastases

On documentation of metachronous, potentially resectable, metastatic disease with dedicated contrast-enhanced CT or MRI, characterization of the disease extent using PET/CT scan should be considered in select cases if a surgical cure of M1 disease is feasible. PET/CT is used at this juncture to promptly characterize the extent of metastatic disease, and to identify possible sites of extrahepatic disease that could preclude surgery.^{661,662} A recent randomized clinical trial of patients with resectable metachronous metastases also assessed the role of PET/CT in the workup of potential curable disease.⁶⁶³ While there was no impact of PET/CT on survival, surgical management was changed in 8% of patients after PET/CT. For example, resection was not undertaken for 2.7% of patients because additional metastatic disease was identified (bone, peritoneum/omentum, abdominal nodes). In addition, 1.5% of patients had more extensive hepatic resections and 3.4% had additional organ surgery. An additional 8.4% of patients in the PET/CT arm had false-positive results, many of which were investigated with biopsies or additional imaging. A meta-analysis of 18 studies including 1059 patients with hepatic colorectal metastases found that PET or PET/CT results changed management in 24% of patients.⁶⁶⁴

As with other conditions in which stage IV disease is diagnosed, a tumor analysis (metastases or original primary) of *KRAS/NRAS* genotype should be performed to define whether anti-EGFR agents can be considered among the potential options. Although *BRAF* genotyping can be considered for patients with tumors characterized by the wild-type *KRAS/NRAS* genes, this testing is currently optional and is not a necessary part of deciding whether to use anti-EGFR agents (see *Perioperative Cetuximab and Panitumumab for Resectable Metastatic Disease: The Role of KRAS, NRAS, and BRAF Status*, above). Close communication between members of the multidisciplinary treatment team is recommended, including upfront evaluation by a surgeon experienced in the resection of hepatobiliary and lung metastases.

The management of metachronous metastatic disease is distinguished from that of synchronous disease through also including an evaluation of the chemotherapy history of the patient and through the absence of transabdominal resection. Patients with resectable disease are classified according to whether they have undergone previous chemotherapy. For patients who have resectable metastatic disease, treatment is resection with 6 months of perioperative chemotherapy (pre- or postoperative or a combination of both), with choice of regimens based on previous therapy. Locally ablative procedures can be considered instead of or in addition to resection in cases of liver oligometastases (see *Local Therapies for Metastases*, above), but resection is preferred. For patients without a history of chemotherapy use, FOLFOX or CapeOx are preferred, with FLOX, capecitabine, and 5-FU/LV as additional category 2B options. There are also cases when perioperative chemotherapy is not recommended in resectable metachronous disease. In particular, patients with a history of previous chemotherapy and upfront resection can be observed or may be given an active regimen for advanced disease, and the same is true for

patients whose tumors grew on therapy before resection (category 2B for the use of biologic agents in these settings). Observation is preferred if oxaliplatin-based therapy was previously administered. In addition, observation is an appropriate option for patients whose tumors grew through neoadjuvant treatment.

Patients determined to have unresectable disease through cross-sectional imaging scan (including those considered potentially convertible) should receive an active systemic therapy regimen based on prior chemotherapy history (see *Therapy after Progression* in the Discussion section of the NCCN Guidelines for Colon Cancer, available at www.NCCN.org). In the case of liver metastases only, HAI therapy with or without systemic 5-FU/LV (category 2B) is an option at centers with experience in the surgical and medical oncologic aspects of this procedure. Patients receiving palliative chemotherapy should be monitored with CT or MRI scans approximately every 2 to 3 months.

Endpoints for Advanced Colorectal Cancer Clinical Trials

In the past few years, there has been much debate over what endpoints are most appropriate for clinical trials in advanced colorectal cancer.⁶⁶⁵ Quality of life is an outcome that is rarely measured but is of unquestioned clinical relevance.⁶⁶⁶ While OS is also of clear clinical relevance, it is often not used because large numbers of patients and long follow-up periods are required.⁶⁶⁶ PFS is often used as a surrogate, but its correlation with OS is inconsistent at best, especially when subsequent lines of therapy are administered.⁶⁶⁶⁻⁶⁶⁸ GROUP Español Multidisciplinar en Cancer Digestivo (GEMCAD) recently proposed particular aspects of clinical trial design to be incorporated into trials that use PFS as an endpoint.⁶⁶⁹



A recent study, in which individual patient data from 3 randomized controlled trials were pooled, tested endpoints that take into account subsequent lines of therapy: duration of disease control, which is the sum of PFS times of each active treatment; and time to failure of strategy, which includes intervals between treatment courses and ends when the planned lines of treatment end (because of death, progression, or administration of a new agent).⁶⁶⁷ The authors found a better correlation between these endpoints and OS than between PFS and OS. Another alternative endpoint, time to tumor growth, has also been suggested to predict OS.^{670,671} Further evaluation of these and other surrogate endpoints is warranted.

Post-Treatment Surveillance

After curative-intent surgery and adjuvant chemotherapy, if administered, post-treatment surveillance of patients with colorectal cancer is performed to evaluate for possible therapeutic complications, discover a recurrence that is potentially resectable for cure, and identify new metachronous neoplasms at a preinvasive stage. An analysis of data from 20,898 patients enrolled in 18 large, adjuvant colon cancer, randomized trials showed that 80% of recurrences occurred in the first 3 years after surgical resection of the primary tumor,⁶⁷² and a recent study found that 95% of recurrences occurred in the first 5 years.⁶⁷³

The approach to monitoring and surveillance of patients with rectal cancer is similar to that described for colon cancer.

Surveillance for Locoregional Disease

Advantages of more intensive follow-up of patients after treatment of stage II and/or stage III disease have been demonstrated prospectively in several older studies⁶⁷⁴⁻⁶⁷⁶ and in multiple meta-analyses of randomized controlled trials designed to compare low-intensity and

high-intensity programs of surveillance.⁶⁷⁷⁻⁶⁸⁰ In the final analysis of Intergroup trial 0114 comparing bolus 5-FU to bolus 5-FU/LV in patients with surgically resectable rectal cancer, local recurrence rates continued to rise after 5 years.²⁵⁴ Further, a population-based report indicated that long-term survival is possible in patients treated for local recurrence of rectal cancer (overall 5-year relative survival rate of 15.6%), thereby providing support for more intensive post-treatment follow-up in these patients.⁶⁸¹

Results from a recent randomized controlled trial of 1202 patients with resected stage I to III disease showed that intensive surveillance imaging or CEA screening resulted in an increased rate of curative-intent surgical treatment compared with a minimum follow-up group that only received testing if symptoms occurred, but no advantage was seen in the CEA and CT combination arm (2.3% in the minimum follow-up group, 6.7% in the CEA group, 8% in the CT group, and 6.6% in the CEA plus CT group).⁶⁸² In this study, no mortality benefit to regular monitoring with CEA, CT, or both was observed compared with minimum follow-up (death rate, 18.2% vs. 15.9%; difference, 2.3%; 95% CI, -2.6%–7.1%). The authors concluded that any strategy of surveillance is unlikely to provide a large survival advantage over a symptom-based approach.⁶⁸²

The CEAwatch trial compared usual follow-up care to CEA measurements every two months, with imaging performed if CEA increases were seen twice, in 3223 patients at 11 hospitals treated for non-metastatic colorectal cancer in the Netherlands.⁶⁸³ The intensive CEA surveillance protocol resulted in the detection of more total recurrences and recurrences that could be treated with curative intent than usual follow-up, and the time to detection of recurrent disease was shorter. Another randomized trial of 1228 patients found that more intensive surveillance led to earlier detection of recurrences than a less



intensive program (less frequent colonoscopy and liver ultrasound and the absence of an annual chest x-ray) but did not affect OS.⁶⁸⁴

The randomized phase III PRODIGE 13 trial will compare 5-year OS after intensive radiological monitoring (abdominal ultrasound, chest/abdomen/pelvis CT, and CEA) with a lower intensity program (abdominal ultrasound and chest x-ray) in patients with resected stage II or III colon or rectal tumors.⁶⁸⁵

Clearly, controversies remain regarding selection of optimal strategies for following patients after potentially curative colorectal cancer surgery, and the panel's recommendations are based mainly on consensus. The panel endorses surveillance as a means to identify patients who are potentially curable of metastatic disease with surgical resection.

The following panel recommendations for post-treatment surveillance pertain to patients with stage I through stage III disease who have undergone successful treatment (ie, no known residual disease): history and physical examination every 3 to 6 months for 2 years, and then every 6 months for a total of 5 years; and a CEA test (also see *Managing an Increasing CEA Level*, below) is recommended at baseline and every 3 to 6 months for 2 years,⁶⁸⁶ then every 6 months for a total of 5 years for patients with stage III disease and those with stage II disease if the clinician determines that the patient is a potential candidate for aggressive curative surgery.^{677,686,687}

Colonoscopy is recommended at approximately 1 year following resection (or at approximately 3 to 6 months post-resection if not performed preoperatively due to an obstructing lesion). Repeat colonoscopy is typically recommended at 3 years, and then every 5 years thereafter, unless follow-up colonoscopy indicates advanced adenoma (villous polyp, polyp >1 cm, or high-grade dysplasia), in which

case colonoscopy should be repeated in 1 year.⁶⁸⁸ More frequent colonoscopies may be indicated in patients who present with colorectal cancer before age 50.⁶⁸⁸ Surveillance colonoscopies are primarily aimed at identifying and removing metachronous polyps since data show that patients with a history of colorectal cancer have an increased risk of developing second cancers,⁶⁸⁹ particularly in the first 2 years following resection. The use of post-treatment surveillance colonoscopy has not been shown to improve survival through the early detection of recurrence of the original colorectal cancer.⁶⁸⁸

Proctoscopy with EUS or MRI is recommended to evaluate the rectal anastomosis for local recurrence only in patients treated with transanal excision. Proctoscopy is not recommended for other patients, because isolated local recurrences are rarely found in these patients and are rarely curable. In fact, in a single-center study of 112 patients who had TME for rectal cancer, only one local recurrence occurred, and it was not identified by rectal surveillance but by CEA and symptoms.⁶⁹⁰ In these 112 patients, 20 anoscopies, 44 proctoscopies, and 495 flexible sigmoidoscopies were performed.

Chest, abdominal, and pelvic CT scans are recommended every 3 to 6 months for 2 years and then every 6 to 12 months for up to 5 years.^{677,691} CT scan is recommended to monitor for the presence of potentially resectable metastatic lesions, primarily in the lung and the liver. Hence, CT scan is not routinely recommended in patients who are not candidates for potentially curative resection of liver or lung metastases. A recent analysis of patients with resected or ablated colorectal liver metastases found that the frequency of surveillance imaging did not correlate with time to second procedure or median survival duration.⁶⁹² Those scanned once per year survived a median of 54 months versus 43 months for those scanned 3 to 4 times per year (*P*



= .08), suggesting that annual scans may be sufficient in this population.

Routine CEA monitoring and CT scanning are not recommended beyond 5 years. In addition, use of PET/CT to monitor for disease recurrence is not recommended.^{691,693} The CT that accompanies a PET/CT is usually a noncontrast CT, and therefore is not of ideal quality for routine surveillance.

The ASCO Clinical Practice Guidelines Committee recently endorsed the Follow-up Care, Surveillance Protocol, and Secondary Prevention Measures for Survivors of Colorectal Cancer, from Cancer Care Ontario (CCO).^{694,695} These guidelines differ only slightly from the surveillance recommendations in these NCCN Guidelines for Rectal Cancer. While ASCO/CCO recommend abdominal and chest CT annually for 3 years, the NCCN Panel recommends semi-annual to annual scans for 5 years (category 2B for more frequent than annual scanning). The panel bases its recommendation on the fact that approximately 10% of disease recurrences occur after 3 years.^{673,696} The American Society of Colon and Rectal Surgeons also released surveillance guidelines, which are also very similar to NCCN surveillance recommendations.⁶⁹⁷ One exception is the inclusion of intensive surveillance for patients with resected stage I colon or rectal cancer if the provider deems the patient to be at increased risk for recurrence.

Surveillance for Metastatic Disease

Patients who had resection of metastatic colorectal cancer can undergo subsequent curative-intent resection of recurrent disease (*see Surgical Management of Colorectal Metastases*, above). A retrospective analysis of 952 patients who underwent resection at Memorial Sloan Kettering Cancer Center showed that 27% of patients with recurrent disease underwent curative-intent resection and that 25% of those patients (6%

of recurrences; 4% of the initial population) were free of disease for ≥36 months.⁶⁹⁸

Panel recommendations for surveillance of patients with stage IV rectal cancer with NED after curative-intent surgery and perioperative treatment are similar to those listed for patients treated for locoregional rectal cancer. One difference is that the frequency of chest, abdominal, and pelvic CT scan for the first 2 years is every 3 to 6 months (category 2B for frequency <6 months), whereas it is every 6 to 12 months for patients treated for locoregional disease.

Managing an Increasing CEA Level

Managing patients with an elevated CEA level after resection should include colonoscopy; chest, abdominal, and pelvic CT scans; physical examination; and consideration of a PET/CT scan. If imaging study results are normal in the face of a rising CEA, repeat CT scans are recommended every 3 months until either disease is identified or CEA level stabilizes or declines.

In a recent retrospective chart review at Memorial Sloan Kettering Cancer Center, approximately half of elevations in CEA levels after R0 resection of locoregional colorectal cancer were false positives, with most being single high readings or repeat readings in the range of 5 to 15 ng/mL.⁶⁹⁹ In this study, false-positive results >15 ng/mL were rare, and all results >35 ng/mL represented true-positives. Following a systematic review and meta-analysis, the pooled sensitivity and specificity of CEA at a cut-off of 10 ng/mL were calculated at 68% (95% CI, 53%–79%) and 97% (95% CI, 90%–99%), respectively.^{700,701} In the first 2 years post-resection, a CEA cut-off of 10 ng/mL is estimated to detect 20 recurrences, miss 10 recurrences, and result in 29 false positives.



Panel opinion was divided on the usefulness of PET/CT scan in the scenario of an elevated CEA with negative, good-quality CT scans (ie, some panel members favored use of PET/CT in this scenario whereas others noted that the likelihood of PET/CT identifying surgically curable disease in the setting of negative good-quality CT scans is vanishingly small). A recent systematic review and meta-analysis found 11 studies (510 patients) that addressed the use of PET/CT in this setting.⁷⁰² The pooled estimates of sensitivity and specificity for the detection of tumor recurrence were 94.1% (95% CI, 89.4–97.1%) and 77.2% (95% CI, 66.4–85.9), respectively. Use of PET/CT scans in this scenario is permissible within these guidelines. The panel does not recommend a so-called blind or CEA-directed laparotomy or laparoscopy for patients whose workup for an increased CEA level is negative,⁷⁰³ nor do they recommend use of anti-CEA-radiolabeled scintigraphy.

Treatment of Locally Recurrent Disease

Locally recurrent rectal cancer is characterized by isolated pelvic/anastomotic recurrence of disease. In a single-center study, Yu et al reported low rates of 5-year local recurrence (ie, 5-year locoregional control rate of 91%) for patients with rectal cancer treated with surgery and either RT or chemoRT, and 49% of recurrences occurred in the low pelvic and presacral regions with an additional 14% occurring in the mid and high pelvis.⁷⁰⁴ Patients with disease recurrence at the anastomotic site are more likely to be cured following re-resection than those with an isolated pelvic recurrence.^{705,706}

Potentially resectable isolated pelvic/anastomotic recurrence is optimally managed with resection followed by adjuvant chemoRT or with preoperative RT and concurrent chemotherapy. IORT or brachytherapy should be considered with resection if it can be safely delivered.^{374,707-709} In a study of 43 consecutive patients with advanced

pelvic recurrence of colorectal cancer who had not undergone prior RT, treatment with 5 weeks of 5-FU by infusion concurrent with RT enabled the majority of patients (77%) to undergo re-resection with curative intent.⁷⁰⁶ Studies of patients who previously received pelvic radiotherapy show that re-irradiation can be effective, with acceptable rates of toxicity.⁷¹⁰⁻⁷¹² In one such study of 48 patients with recurrent rectal cancer and a history of pelvic radiation, the 3-year rate of grade 3 to 4 late toxicity was 35%, and 36% of treated patients were able to undergo surgery following radiation.⁷¹⁰ IMRT can be used in this setting of re-irradiation.

Patients with unresectable lesions are treated with chemotherapy with or without radiation according to their ability to tolerate therapy. Debulking that results in gross residual cancer is not recommended.

Survivorship

The panel recommends that a prescription for survivorship and transfer of care to the primary care physician be written.⁷¹³ The oncologist and primary care provider should have defined roles in the surveillance period, with roles communicated to the patient. The care plan should include an overall summary of treatments received, including surgeries, radiation treatments, and chemotherapy. The possible expected time to resolution of acute toxicities, long-term effects of treatment, and possible late sequelae of treatment should be described. Finally, surveillance and health behavior recommendations should be part of the care plan.

Disease preventive measures, such as immunizations; early disease detection through periodic screening for second primary cancers (eg, breast, cervical, or prostate cancers); and routine good medical care and monitoring are recommended (see the NCCN Guidelines for Survivorship, available at www.NCCN.org). Additional health monitoring



should be performed as indicated under the care of a primary care physician. Survivors are encouraged to maintain a therapeutic relationship with a primary care physician throughout their lifetime.⁷¹⁴

Other recommendations include monitoring for late sequelae of rectal cancer or of the treatment of rectal cancer, such as chronic diarrhea or incontinence (eg, patients with stoma).⁷¹⁵⁻⁷²⁰ Urogenital dysfunction following resection and/or pelvic irradiation is common.^{715,721-723} Patients should be screened for sexual dysfunction, erectile dysfunction, dyspareunia, vaginal dryness, and urinary incontinence, frequency, and urgency. Referral to a gynecologist or urologist can be considered for persistent symptoms. Other long-term problems common to colorectal cancer survivors include oxaliplatin-induced peripheral neuropathy, fatigue, insomnia, cognitive dysfunction, and emotional or social distress.⁷²⁴⁻⁷²⁹ Specific management interventions to address side effects of colorectal cancer have been described,⁷³⁰ and a survivorship care plan for patients with colorectal cancer has recently been published.⁷³¹

The NCCN Guidelines for Survivorship, available at www.NCCN.org, provide screening, evaluation, and treatment recommendations for common consequences of cancer and cancer treatment to aid health care professionals who work with survivors of adult-onset cancer in the post-treatment period, including those in specialty cancer survivor clinics and primary care practices. The NCCN Guidelines for Survivorship include many topics with potential relevance to survivors of colorectal cancer, including Anxiety, Depression, and Distress; Cognitive Dysfunction; Fatigue; Pain; Sexual Dysfunction; Healthy Lifestyles; and Immunizations. Concerns related to employment, insurance, and disability are also discussed. The ACS has also established guidelines for the care of survivors of colorectal cancer, including surveillance for recurrence, screening for subsequent primary

malignancies, the management of physical and psychosocial effects of cancer and its treatment, and promotion of healthy lifestyles.⁷¹⁴

Healthy Lifestyles for Survivors of Colorectal Cancer

Evidence indicates that certain lifestyle characteristics, such as smoking cessation, maintaining a healthy BMI, engaging in regular exercise, and making certain dietary choices are associated with improved outcomes and quality of life after treatment for colorectal cancer. In a prospective observational study of patients with stage III colon cancer enrolled in the CALGB 89803 adjuvant chemotherapy trial, DFS was found to be directly related to how much exercise these patients received.⁷³² In addition, a recent study of a large cohort of men treated for stage I through III colorectal cancer showed an association between increased physical activity and lower rates of colorectal cancer-specific mortality and overall mortality.⁷³³ More recent data support the conclusion that physical activity improves outcomes. In a cohort of over 2000 survivors of non-metastatic colorectal cancer, those who spent more time in recreational activity had a lower mortality than those who spent more leisure time sitting.⁷³⁴ In addition, recent evidence suggests that both pre- and post-diagnosis physical activity decreases colorectal cancer mortality. Women enrolled in the Women's Health Initiative study who subsequently developed colorectal cancer had lower colorectal cancer-specific mortality (HR, 0.68; 95% CI, 0.41–1.13) and all-cause mortality (HR, 0.63; 95% CI, 0.42–0.96) if they reported high levels of physical activity.⁷³⁵ Similar results were seen in other studies and in recent meta-analyses of prospective studies.⁷³⁶⁻⁷³⁹

A retrospective study of patients with stage II and III colon cancer enrolled in NSABP trials from 1989 to 1994 showed that patients with a BMI of 35 kg/m² or greater had an increased risk of disease recurrence and death.⁷⁴⁰ Recent analyses confirm the increased risk for recurrence

and death in obese patients.⁷⁴¹ Data from the ACCENT database also found that pre-diagnosis BMI has a prognostic impact on outcomes in patients with stage II/III colorectal cancer undergoing adjuvant therapy.⁷⁴² However, a recent analysis of participants in the Cancer Prevention Study-II Nutrition Cohort who subsequently developed non-metastatic colorectal cancer found that pre-diagnosis obesity but not post-diagnosis obesity was associated with higher all-cause and colorectal cancer-specific mortality.⁷⁴³ A meta-analysis of prospective cohort studies found that pre-diagnosis obesity was associated with increased colorectal cancer-specific and all-cause mortality.⁷⁴⁴ Other analyses confirm the increased risk for recurrence and death in obese patients.^{741,745-748}

In contrast, pooled data from first-line clinical trials in the ARCAD database indicate that a low BMI may be associated with an increased risk of progression and death in the metastatic setting, whereas a high BMI may not be.⁷⁴⁹ In addition, results of one retrospective observational study of a cohort of 3408 patients with resected stage I to III colorectal cancer suggest that the relationship between mortality and BMI might be U shaped, with the lowest mortality for those with BMI 28 kg/m².⁷⁵⁰ However, several possible explanations for this so-called “obesity paradox” have been suggested.⁷⁵¹ Overall the panel believes that survivors of colorectal cancer should be encouraged to achieve and maintain a healthy body weight (see the NCCN Guidelines for Survivorship at www.NCCN.org).

A diet consisting of more fruits, vegetables, poultry, and fish, less red meat, higher in whole grains, and lower in refined grains and concentrated sweets was found to be associated with an improved outcome in terms of cancer recurrence or death.⁷⁵² There is also some evidence that higher postdiagnosis intake of total milk and calcium may be associated with a lower risk of death in patients with stage I, II, or III

colorectal cancer.⁷⁶ Recent analysis of the CALGB 89803 trial found that higher dietary glycemic load was also associated with an increased risk of recurrence and mortality in patients with stage III disease.⁷⁵³ Another analysis of the data from CALGB 89803 found an association between high intake of sugar-sweetened beverages and an increased risk of recurrence and death in patients with stage III colon cancer.⁷⁵⁴ The link between red and processed meats and mortality in survivors of non-metastatic colorectal cancer has been further supported by recent data from the Cancer Prevention Study II Nutrition Cohort, in which survivors with consistently high intake had a higher risk of colorectal cancer-specific mortality than those with low intake (RR, 1.79; 95% CI, 1.11–2.89).⁶⁹

A discussion of lifestyle characteristics that may be associated with a decreased risk of colorectal cancer recurrence, such as those recommended by the American Cancer Society (ACS),⁷⁵⁵ also provides “a teachable moment” for the promotion of overall health, and an opportunity to encourage patients to make choices and changes compatible with a healthy lifestyle. In addition, a recent trial showed that telephone-based health behavior coaching had a positive effect on physical activity, diet, and BMI in survivors of colorectal cancer, suggesting that survivors may be open to health behavior change.⁷⁵⁶

Therefore, survivors of colorectal cancer should be encouraged to maintain a healthy body weight throughout life; adopt a physically active lifestyle (at least 30 minutes of moderate intensity activity on most days of the week); consume a healthy diet with emphasis on plant sources; limit alcohol consumption; and quit smoking.⁷⁵⁷ Activity recommendations may require modification based on treatment sequelae (ie, ostomy, neuropathy), and diet recommendations may be modified based on the severity of bowel dysfunction.⁷⁵⁸



Secondary Chemoprevention for Colorectal Cancer Survivors

Limited data suggest a link between post-colorectal-cancer-diagnosis statin use and increased survival.^{89,759,760} A meta-analysis that included 4 studies found that post-diagnosis statin use increased OS (HR, 0.76; 95% CI, 0.68–0.85; $P < .001$) and cancer-specific survival (HR, 0.70; 95% CI, 0.60–0.81; $P < .001$).⁷⁵⁹

Abundant data show that low-dose aspirin therapy after a diagnosis of colorectal cancer decreases the risk of recurrence and death.⁷⁶¹⁻⁷⁶⁷ For example, a population-based, observational, retrospective cohort study of 23,162 patients with colorectal cancer in Norway found that post-diagnosis aspirin use was associated with improved colorectal cancer-specific survival (HR, 0.85; 95% CI, 0.79–0.92) and OS (HR, 0.95; 95% CI, 0.90–1.01).⁷⁶¹ Some evidence suggests that tumor mutations in *PIK3CA* may be predictive for response to aspirin, although the data are somewhat inconsistent and other predictive markers have also been suggested.^{763,768-772}

Based on these data, the panel believes that survivors of colorectal cancer can consider taking low-dose aspirin to reduce their risk of recurrence and death. Importantly, aspirin may increase the risk of gastrointestinal bleeding and hemorrhagic stroke, and these risks should be discussed with colorectal cancer survivors.⁷⁷³

Summary

The NCCN Rectal Cancer Panel believes that a multidisciplinary approach, including representation from gastroenterology, medical oncology, surgical oncology, radiation oncology, and radiology is necessary for treating patients with rectal cancer. Adequate pathologic assessment of the resected lymph nodes is important with a goal of evaluating at least 12 nodes when possible. Patients with very-early-

stage tumors that are node-negative by endorectal ultrasound or endorectal or pelvic MRI and who meet carefully defined criteria can be managed with a transanal excision. A transabdominal resection is appropriate for all other rectal lesions. Perioperative chemoRT and chemotherapy are preferred for the majority of patients with suspected or proven T3-4 disease and/or regional node involvement.

The recommended post-treatment surveillance program for patients following treatment for rectal cancer includes serial CEA determinations, as well as periodic chest, abdominal, and pelvic CT scans, and periodic evaluation by colonoscopy. Patients with recurrent localized disease should be considered for resection with chemotherapy and radiation. If resection is not possible, then chemotherapy is given with or without radiation.

A patient with metastatic disease in the liver or lung should be considered for surgical resection if he or she is a candidate for surgery and if complete resection (R0) can be achieved. Perioperative chemotherapy and chemoRT are used in the synchronous setting, and perioperative chemotherapy is used in the metachronous setting.

Recommendations for patients with disseminated, unresectable metastatic disease represent a continuum of care in which lines of treatment are blurred rather than discrete. Principles to consider at the start of therapy include pre-planned strategies for altering therapy for patients in both the presence and absence of disease progression and plans for adjusting therapy for patients who experience certain toxicities. Recommended initial therapy options for advanced or metastatic disease depend on whether or not the patient is appropriate for intensive therapy. The more intensive initial therapy options include FOLFOX, FOLFIRI, CapeOX, and FOLFOXIRI. Addition of a biologic agent (ie, bevacizumab, cetuximab, panitumumab) is listed as an option



in combination with some of these regimens, depending on available data. Systemic therapy options for patients with progressive disease are dependent on the choice of initial therapy. The panel endorses the concept that treating patients in a clinical trial has priority over standard treatment regimens.



References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA Cancer J Clin* 2016;66:7-30. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26742998>.
2. Cheng L, Eng C, Nieman LZ, et al. Trends in colorectal cancer incidence by anatomic site and disease stage in the United States from 1976 to 2005. *Am J Clin Oncol* 2011;34:573-580. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21217399>.
3. Henley SJ, Singh SD, King J, et al. Invasive cancer incidence and survival--United States, 2011. *MMWR Morb Mortal Wkly Rep* 2015;64:237-242. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25763875>.
4. Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin* 2011;61:212-236. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21685461>.
5. Bailey CE, Hu CY, You YN, et al. Increasing disparities in the age-related incidences of colon and rectal cancers in the United States, 1975-2010. *JAMA Surg* 2014;1-6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25372703>.
6. U.S. National Library of Medicine-Key MEDLINE® Indicators. Available at: http://www.nlm.nih.gov/bsd/bsd_key.html. Accessed March 1, 2016.
7. Ahsan H, Neugut AI, Garbowski GC, et al. Family history of colorectal adenomatous polyps and increased risk for colorectal cancer. *Ann Intern Med* 1998;128:900-905. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9634428>.
8. Bonelli L, Martines H, Conio M, et al. Family history of colorectal cancer as a risk factor for benign and malignant tumours of the large bowel. A case-control study. *Int J Cancer* 1988;41:513-517. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3356486>.
9. Hemminki K, Chen B. Familial risk for colorectal cancers are mainly due to heritable causes. *Cancer Epidemiol Biomarkers Prev* 2004;13:1253-1256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15247139>.
10. Hemminki K, Eng C. Clinical genetic counselling for familial cancers requires reliable data on familial cancer risks and general action plans. *J Med Genet* 2004;41:801-807. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15520403>.
11. Quintero E, Carrillo M, Leoz ML, et al. Risk of advanced neoplasia in first-degree relatives with colorectal cancer: a large multicenter cross-sectional study. *PLoS Med* 2016;13:e1002008. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27138769>.
12. Hampel H, Frankel WL, Martin E, et al. Feasibility of screening for Lynch syndrome among patients with colorectal cancer. *J Clin Oncol* 2008;26:5783-5788. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18809606>.
13. Lynch HT, de la Chapelle A. Hereditary colorectal cancer. *N Engl J Med* 2003;348:919-932. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12621137>.
14. Galiatsatos P, Foulkes WD. Familial adenomatous polyposis. *Am J Gastroenterol* 2006;101:385-398. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16454848>.
15. Hennink SD, van der Meulen-de Jong AE, Wolterbeek R, et al. Randomized comparison of surveillance intervals in familial colorectal cancer. *J Clin Oncol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26527788>.
16. Aaltonen LA, Salovaara R, Kristo P, et al. Incidence of hereditary nonpolyposis colorectal cancer and the feasibility of molecular screening for the disease. *N Engl J Med* 1998;338:1481-1487. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9593786>.



17. Hampel H, Frankel WL, Martin E, et al. Screening for the Lynch syndrome (hereditary nonpolyposis colorectal cancer). *N Engl J Med* 2005;352:1851-1860. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15872200>.

18. Hendriks YM, de Jong AE, Morreau H, et al. Diagnostic approach and management of Lynch syndrome (hereditary nonpolyposis colorectal carcinoma): a guide for clinicians. *CA Cancer J Clin* 2006;56:213-225. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16870997>.

19. Beamer LC, Grant ML, Espenschied CR, et al. Reflex immunohistochemistry and microsatellite instability testing of colorectal tumors for Lynch syndrome among US cancer programs and follow-up of abnormal results. *J Clin Oncol* 2012;30:1058-1063. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22355048>.

20. Burt RW. Who should have genetic testing for the lynch syndrome? *Ann Intern Med* 2011;155:127-128. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21768586>.

21. Matloff J, Lucas A, Polydorides AD, Itzkowitz SH. Molecular tumor testing for Lynch syndrome in patients with colorectal cancer. *J Natl Compr Canc Netw* 2013;11:1380-1385. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24225971>.

22. Ward RL, Hicks S, Hawkins NJ. Population-based molecular screening for Lynch syndrome: implications for personalized medicine. *J Clin Oncol* 2013;31:2554-2562. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23733757>.

23. Recommendations from the EGAPP Working Group: genetic testing strategies in newly diagnosed individuals with colorectal cancer aimed at reducing morbidity and mortality from Lynch syndrome in relatives. *Genet Med* 2009;11:35-41. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19125126>.

24. Ladabaum U, Wang G, Terdiman J, et al. Strategies to identify the Lynch syndrome among patients with colorectal cancer: a cost-effectiveness analysis. *Ann Intern Med* 2011;155:69-79. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21768580>.

25. Palomaki GE, McClain MR, Melillo S, et al. EGAPP supplementary evidence review: DNA testing strategies aimed at reducing morbidity and mortality from Lynch syndrome. *Genet Med* 2009;11:42-65. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19125127>.

26. Giardiello FM, Allen JI, Axilbund JE, et al. Guidelines on genetic evaluation and management of Lynch syndrome: a consensus statement by the US Multi-Society Task Force on Colorectal Cancer. *Am J Gastroenterol* 2014;109:1159-1179. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25070057>.

27. Rubenstein JH, Enns R, Heidelbaugh J, et al. American Gastroenterological Association Institute guideline on the diagnosis and management of Lynch syndrome. *Gastroenterology* 2015;149:777-782. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26226577>.

28. Heald B, Plesec T, Liu X, et al. Implementation of universal microsatellite instability and immunohistochemistry screening for diagnosing lynch syndrome in a large academic medical center. *J Clin Oncol* 2013;31:1336-1340. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23401454>.

29. de Rosa N, Rodriguez-Bigas MA, Chang GJ, et al. DNA mismatch repair deficiency in rectal cancer: benchmarking its impact on prognosis, neoadjuvant response prediction, and clinical cancer genetics. *J Clin Oncol* 2016;34:3039-3046. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27432916>.

30. Guinney J, Dienstmann R, Wang X, et al. The consensus molecular subtypes of colorectal cancer. *Nat Med* 2015;21:1350-1356. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26457759>.



31. Beaugerie L, Svrcek M, Seksik P, et al. Risk of colorectal high-grade dysplasia and cancer in a prospective observational cohort of patients with inflammatory bowel disease. *Gastroenterology* 2013;145:166-175 e168. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23541909>.
32. Johnson CM, Wei C, Ensor JE, et al. Meta-analyses of colorectal cancer risk factors. *Cancer Causes Control* 2013;24:1207-1222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23563998>.
33. Lutgens MW, van Oijen MG, van der Heijden GJ, et al. Declining risk of colorectal cancer in inflammatory bowel disease: an updated meta-analysis of population-based cohort studies. *Inflamm Bowel Dis* 2013;19:789-799. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23448792>.
34. Alexander DD, Weed DL, Cushing CA, Lowe KA. Meta-analysis of prospective studies of red meat consumption and colorectal cancer. *Eur J Cancer Prev* 2011;20:293-307. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21540747>.
35. Cheng J, Chen Y, Wang X, et al. Meta-analysis of prospective cohort studies of cigarette smoking and the incidence of colon and rectal cancers. *Eur J Cancer Prev* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24722538>.
36. De Bruijn KM, Arends LR, Hansen BE, et al. Systematic review and meta-analysis of the association between diabetes mellitus and incidence and mortality in breast and colorectal cancer. *Br J Surg* 2013;100:1421-1429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24037561>.
37. Esposito K, Chiodini P, Capuano A, et al. Colorectal cancer association with metabolic syndrome and its components: a systematic review with meta-analysis. *Endocrine* 2013;44:634-647. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23546613>.
38. Fedirko V, Tramacere I, Bagnardi V, et al. Alcohol drinking and colorectal cancer risk: an overall and dose-response meta-analysis of published studies. *Ann Oncol* 2011;22:1958-1972. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21307158>.
39. Huxley RR, Ansary-Moghaddam A, Clifton P, et al. The impact of dietary and lifestyle risk factors on risk of colorectal cancer: a quantitative overview of the epidemiological evidence. *Int J Cancer* 2009;125:171-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19350627>.
40. Kitahara CM, Berndt SI, de Gonzalez AB, et al. Prospective investigation of body mass index, colorectal adenoma, and colorectal cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *J Clin Oncol* 2013;31:2450-2459. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23715565>.
41. Kyu HH, Bachman VF, Alexander LT, et al. Physical activity and risk of breast cancer, colon cancer, diabetes, ischemic heart disease, and ischemic stroke events: systematic review and dose-response meta-analysis for the Global Burden of Disease Study 2013. *BMJ* 2016;354:i3857. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27510511>.
42. Larsson SC, Orsini N, Wolk A. Diabetes mellitus and risk of colorectal cancer: a meta-analysis. *J Natl Cancer Inst* 2005;97:1679-1687. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16288121>.
43. Lauby-Secretan B, Scoccianti C, Loomis D, et al. Body fatness and cancer--viewpoint of the IARC Working Group. *N Engl J Med* 2016;375:794-798. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27557308>.
44. Levi Z, Kark JD, Barchana M, et al. Measured body mass index in adolescence and the incidence of colorectal cancer in a cohort of 1.1 million males. *Cancer Epidemiol Biomarkers Prev* 2011;20:2524-2531. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22056504>.
45. Luo W, Cao Y, Liao C, Gao F. Diabetes mellitus and the incidence and mortality of colorectal cancer: a meta-analysis of 24 cohort studies.



Colorectal Dis 2012;14:1307-1312. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23046351>.

46. Ma Y, Yang Y, Wang F, et al. Obesity and risk of colorectal cancer: a systematic review of prospective studies. PLoS One 2013;8:e53916. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23349764>.

47. Magalhaes B, Peleteiro B, Lunet N. Dietary patterns and colorectal cancer: systematic review and meta-analysis. Eur J Cancer Prev 2012;21:15-23. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21946864>.

48. Moore SC, Lee IM, Weiderpass E, et al. Association of leisure-time physical activity with risk of 26 types of cancer in 1.44 million adults. JAMA Intern Med 2016;176:816-825. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/27183032>.

49. Parajuli R, Bjerkaas E, Tverdal A, et al. The increased risk of colon cancer due to cigarette smoking may be greater in women than men. Cancer Epidemiol Biomarkers Prev 2013;22:862-871. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23632818>.

50. Schmid D, Leitzmann MF. Television viewing and time spent sedentary in relation to cancer risk: a meta-analysis. J Natl Cancer Inst 2014;106. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24935969>.

51. Yuhara H, Steinmaus C, Cohen SE, et al. Is diabetes mellitus an independent risk factor for colon cancer and rectal cancer? Am J Gastroenterol 2011;106:1911-1921; quiz 1922. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21912438>.

52. Keum N, Greenwood DC, Lee DH, et al. Adult weight gain and adiposity-related cancers: a dose-response meta-analysis of prospective observational studies. J Natl Cancer Inst 2015;107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25618901>.

53. Klatsky AL, Li Y, Nicole Tran H, et al. Alcohol intake, beverage choice, and cancer: a cohort study in a large kaiser permanente

population. Perm J 2015;19:28-34. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25785639>.

54. Shen D, Mao W, Liu T, et al. Sedentary behavior and incident cancer: a meta-analysis of prospective studies. PLoS One 2014;9:e105709. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25153314>.

55. Kohler LN, Garcia DO, Harris RB, et al. Adherence to diet and physical activity cancer prevention guidelines and cancer outcomes: a systematic review. Cancer Epidemiol Biomarkers Prev 2016;25:1018-1028. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27340121>.

56. Song M, Giovannucci E. Preventable incidence and mortality of carcinoma associated with lifestyle factors among white adults in the United States. JAMA Oncol 2016. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/27196525>.

57. Keum N, Aune D, Greenwood DC, et al. Calcium intake and colorectal cancer risk: dose-response meta-analysis of prospective observational studies. Int J Cancer 2014;135:1940-1948. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24623471>.

58. Murphy N, Norat T, Ferrari P, et al. Consumption of dairy products and colorectal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC). PLoS One 2013;8:e72715. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24023767>.

59. Ralston RA, Truby H, Palermo CE, Walker KZ. Colorectal cancer and nonfermented milk, solid cheese, and fermented milk consumption: a systematic review and meta-analysis of prospective studies. Crit Rev Food Sci Nutr 2014;54:1167-1179. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24499149>.

60. Orlich MJ, Singh PN, Sabate J, et al. Vegetarian dietary patterns and the risk of colorectal cancers. JAMA Intern Med 2015;175:767-776. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25751512>.

61. Yu XF, Zou J, Dong J. Fish consumption and risk of gastrointestinal cancers: a meta-analysis of cohort studies. *World J Gastroenterol* 2014;20:15398-15412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25386090>.
62. Zhu B, Sun Y, Qi L, et al. Dietary legume consumption reduces risk of colorectal cancer: evidence from a meta-analysis of cohort studies. *Sci Rep* 2015;5:8797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25739376>.
63. Cao Y, Nishihara R, Wu K, et al. Population-wide impact of long-term use of aspirin and the risk for cancer. *JAMA Oncol* 2016;2:762-769. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26940135>.
64. Chan AT, Giovannucci EL, Meyerhardt JA, et al. Long-term use of aspirin and nonsteroidal anti-inflammatory drugs and risk of colorectal cancer. *JAMA* 2005;294:914-923. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16118381>.
65. Flossmann E, Rothwell PM, British Doctors Aspirin T, the UKTIAAT. Effect of aspirin on long-term risk of colorectal cancer: consistent evidence from randomised and observational studies. *Lancet* 2007;369:1603-1613. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17499602>.
66. Friis S, Poulsen AH, Sorensen HT, et al. Aspirin and other non-steroidal anti-inflammatory drugs and risk of colorectal cancer: a Danish cohort study. *Cancer Causes Control* 2009;20:731-740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19122977>.
67. Friis S, Riis AH, Erichsen R, et al. Low-dose aspirin or nonsteroidal anti-inflammatory drug use and colorectal cancer risk: a population-based, case-control study. *Ann Intern Med* 2015;163:347-355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26302241>.
68. Rothwell PM, Wilson M, Elwin CE, et al. Long-term effect of aspirin on colorectal cancer incidence and mortality: 20-year follow-up of five randomised trials. *Lancet* 2010;376:1741-1750. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20970847>.
69. McCullough ML, Gapstur SM, Shah R, et al. Association between red and processed meat intake and mortality among colorectal cancer survivors. *J Clin Oncol* 2013;31:2773-2782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23816965>.
70. Phipps AI, Shi Q, Newcomb PA, et al. Associations between cigarette smoking status and colon cancer prognosis among participants in North Central Cancer Treatment Group phase III trial N0147. *J Clin Oncol* 2013;31:2016-2023. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23547084>.
71. Walter V, Jansen L, Hoffmeister M, Brenner H. Smoking and survival of colorectal cancer patients: systematic review and meta-analysis. *Ann Oncol* 2014;25:1517-1525. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24692581>.
72. Yang B, Jacobs EJ, Gapstur SM, et al. Active smoking and mortality among colorectal cancer survivors: the Cancer Prevention Study II nutrition cohort. *J Clin Oncol* 2015;33:885-893. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25646196>.
73. Song M, Zhang X, Meyerhardt JA, et al. Marine omega-3 polyunsaturated fatty acid intake and survival after colorectal cancer diagnosis. *Gut* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27436272>.
74. Morris EJ, Penegar S, Whitehouse LE, et al. A retrospective observational study of the relationship between family history and survival from colorectal cancer. *Br J Cancer* 2013;108:1502-1507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23511565>.
75. Dik VK, Murphy N, Siersema PD, et al. Prediagnostic intake of dairy products and dietary calcium and colorectal cancer survival-results from the EPIC Cohort Study. *Cancer Epidemiol Biomarkers Prev*



2014;23:1813-1823. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24917183>.

76. Yang B, McCullough ML, Gapstur SM, et al. Calcium, vitamin D, dairy products, and mortality among colorectal cancer survivors: the Cancer Prevention Study-II nutrition cohort. *J Clin Oncol* 2014;32:2335-2343. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24958826>.

77. Bu WJ, Song L, Zhao DY, et al. Insulin therapy and the risk of colorectal cancer in patients with type 2 diabetes: a meta-analysis of observational studies. *Br J Clin Pharmacol* 2014;78:301-309. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25099257>.

78. Cardel M, Jensen SM, Pottegard A, et al. Long-term use of metformin and colorectal cancer risk in type II diabetics: a population-based case-control study. *Cancer Med* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25091592>.

79. Guraya SY. Association of type 2 diabetes mellitus and the risk of colorectal cancer: A meta-analysis and systematic review. *World J Gastroenterol* 2015;21:6026-6031. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26019469>.

80. Karlstad O, Starup-Linde J, Vestergaard P, et al. Use of insulin and insulin analogs and risk of cancer - systematic review and meta-analysis of observational studies. *Curr Drug Saf* 2013;8:333-348. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24215311>.

81. Rokkas T, Portincasa P. Colon neoplasia in patients with type 2 diabetes on metformin: A meta-analysis. *Eur J Intern Med* 2016;33:60-66. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27318643>.

82. Sehdev A, Shih YC, Vekhter B, et al. Metformin for primary colorectal cancer prevention in patients with diabetes: a case-control study in a US population. *Cancer* 2015;121:1071-1078. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25424411>.

83. Singh S, Singh H, Singh PP, et al. Antidiabetic medications and the risk of colorectal cancer in patients with diabetes mellitus: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2013;22:2258-2268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24042261>.

84. Zhang ZJ, Li S. The prognostic value of metformin for cancer patients with concurrent diabetes: a systematic review and meta-analysis. *Diabetes Obes Metab* 2014;16:707-710. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24460896>.

85. Higurashi T, Hosono K, Takahashi H, et al. Metformin for chemoprevention of metachronous colorectal adenoma or polyps in post-polypectomy patients without diabetes: a multicentre double-blind, placebo-controlled, randomised phase 3 trial. *Lancet Oncol* 2016;17:475-483. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26947328>.

86. Mills KT, Bellows CF, Hoffman AE, et al. Diabetes mellitus and colorectal cancer prognosis: a meta-analysis. *Dis Colon Rectum* 2013;56:1304-1319. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24105007>.

87. Mei ZB, Zhang ZJ, Liu CY, et al. Survival benefits of metformin for colorectal cancer patients with diabetes: a systematic review and meta-analysis. *PLoS One* 2014;9:e91818. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24647047>.

88. Kowall B, Stang A, Rathmann W, Kostev K. No reduced risk of overall, colorectal, lung, breast, and prostate cancer with metformin therapy in diabetic patients: database analyses from Germany and the UK. *Pharmacoepidemiol Drug Saf* 2015;24:865-874. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26132313>.

89. Zanders MM, van Herk-Sukel MP, Vissers PA, et al. Are metformin, statin and aspirin use still associated with overall mortality among colorectal cancer patients with diabetes if adjusted for one another? *Br*

J Cancer 2015;113:403-410. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26180924>.

90. Amin MB, Greene FL, Edge S, et al., eds. AJCC Cancer Staging Manual (ed 8th Edition). New York: Springer; 2016.

91. Franko J, Shi Q, Goldman CD, et al. Treatment of colorectal peritoneal carcinomatosis with systemic chemotherapy: a pooled analysis of north central cancer treatment group phase III trials N9741 and N9841. J Clin Oncol 2012;30:263-267. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22162570>.

92. Compton CC, Fielding LP, Burgart LJ, et al. Prognostic factors in colorectal cancer. College of American Pathologists Consensus Statement 1999. Arch Pathol Lab Med 2000;124:979-994. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10888773>.

93. Compton CC, Greene FL. The staging of colorectal cancer: 2004 and beyond. CA Cancer J Clin 2004;54:295-308. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15537574>.

94. Nagtegaal ID, Marijnen CA, Kranenburg EK, et al. Circumferential margin involvement is still an important predictor of local recurrence in rectal carcinoma: not one millimeter but two millimeters is the limit. Am J Surg Pathol 2002;26:350-357. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11859207>.

95. Nagtegaal ID, Quirke P. What is the role for the circumferential margin in the modern treatment of rectal cancer? J Clin Oncol 2008;26:303-312. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18182672>.

96. Wibe A, Rendedal PR, Svensson E, et al. Prognostic significance of the circumferential resection margin following total mesorectal excision for rectal cancer. Br J Surg 2002;89:327-334. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11872058>.

97. Gavioli M, Luppi G, Losi L, et al. Incidence and clinical impact of sterilized disease and minimal residual disease after preoperative radiochemotherapy for rectal cancer. Dis Colon Rectum 2005;48:1851-1857. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16132481>.

98. Rodel C, Martus P, Papadopoulos T, et al. Prognostic significance of tumor regression after preoperative chemoradiotherapy for rectal cancer. J Clin Oncol 2005;23:8688-8696. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16246976>.

99. Nissan A, Stojadinovic A, Shia J, et al. Predictors of recurrence in patients with T2 and early T3, N0 adenocarcinoma of the rectum treated by surgery alone. J Clin Oncol 2006;24:4078-4084. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16943525>.

100. Fujita S, Shimoda T, Yoshimura K, et al. Prospective evaluation of prognostic factors in patients with colorectal cancer undergoing curative resection. J Surg Oncol 2003;84:127-131. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/14598355>.

101. Liebig C, Ayala G, Wilks J, et al. Perineural invasion is an independent predictor of outcome in colorectal cancer. J Clin Oncol 2009;27:5131-5137. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19738119>.

102. Quah HM, Chou JF, Gonen M, et al. Identification of patients with high-risk stage II colon cancer for adjuvant therapy. Dis Colon Rectum 2008;51:503-507. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18322753>.

103. Lo DS, Pollett A, Siu LL, et al. Prognostic significance of mesenteric tumor nodules in patients with stage III colorectal cancer. Cancer 2008;112:50-54. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/18008365>.

104. Nagtegaal ID, Quirke P. Colorectal tumour deposits in the mesorectum and pericolon; a critical review. Histopathology



2007;51:141-149. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17532768>.

105. Puppa G, Maisonneuve P, Sonzogni A, et al. Pathological assessment of pericolonic tumor deposits in advanced colonic carcinoma: relevance to prognosis and tumor staging. *Mod Pathol* 2007;20:843-855. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17491597>.

106. Ueno H, Mochizuki H. Clinical significance of extrabowel skipped cancer infiltration in rectal cancer. *Surg Today* 1997;27:617-622.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9306563>.

107. Ueno H, Mochizuki H, Hashiguchi Y, et al. Extramural cancer deposits without nodal structure in colorectal cancer: optimal categorization for prognostic staging. *Am J Clin Pathol* 2007;127:287-294. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17210518>.

108. Compton CC. Key issues in reporting common cancer specimens: problems in pathologic staging of colon cancer. *Arch Pathol Lab Med* 2006;130:318-324. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16519558>.

109. Lai LL, Fuller CD, Kachnic LA, Thomas CR, Jr. Can pelvic radiotherapy be omitted in select patients with rectal cancer? *Semin Oncol* 2006;33:S70-74. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17178292>.

110. Glynne-Jones R, Mawdsley S, Novell JR. The clinical significance of the circumferential resection margin following preoperative pelvic chemo-radiotherapy in rectal cancer: why we need a common language. *Colorectal Dis* 2006;8:800-807. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17032329>.

111. Adam IJ, Mohamdee MO, Martin IG, et al. Role of circumferential margin involvement in the local recurrence of rectal cancer. *Lancet* 1994;344:707-711. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/7915774>.

112. Mawdsley S, Glynne-Jones R, Grainger J, et al. Can histopathologic assessment of circumferential margin after preoperative pelvic chemoradiotherapy for T3-T4 rectal cancer predict for 3-year disease-free survival? *Int J Radiat Oncol Biol Phys* 2005;63:745-752.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16199310>.

113. Hwang MR, Park JW, Park S, et al. Prognostic impact of circumferential resection margin in rectal cancer treated with preoperative chemoradiotherapy. *Ann Surg Oncol* 2014;21:1345-1351.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24468928>.

114. Tang L, Berlin J, Branton P, et al. Protocol for the examination of specimens from patients with primary carcinoma of the colon and rectum. College of American Pathologists 2016. Available at:

<http://www.cap.org/ShowProperty?nodePath=/UCMCon/Contribution%20Folders/WebContent/pdf/cp-colon-16protocol-3400.pdf>.

115. Sarli L, Bader G, Iusco D, et al. Number of lymph nodes examined and prognosis of TNM stage II colorectal cancer. *Eur J Cancer* 2005;41:272-279. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15661553>.

116. Wong SL, Ji H, Hollenbeck BK, et al. Hospital lymph node examination rates and survival after resection for colon cancer. *JAMA* 2007;298:2149-2154. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18000198>.

117. Pocard M, Panis Y, Malassagne B, et al. Assessing the effectiveness of mesorectal excision in rectal cancer: prognostic value of the number of lymph nodes found in resected specimens. *Dis Colon Rectum* 1998;41:839-845. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9678368>.

118. Tepper JE, O'Connell MJ, Niedzwiecki D, et al. Impact of number of nodes retrieved on outcome in patients with rectal cancer. *J Clin Oncol* 2001;19:157-163. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11134208>.



119. Kidner TB, Ozao-Choy JJ, Yoon J, Bilchik AJ. Should quality measures for lymph node dissection in colon cancer be extrapolated to rectal cancer? *Am J Surg* 2012;204:843-847; discussion 847-848.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22981183>.

120. Baxter NN, Morris AM, Rothenberger DA, Tepper JE. Impact of preoperative radiation for rectal cancer on subsequent lymph node evaluation: a population-based analysis. *Int J Radiat Oncol Biol Phys* 2005;61:426-431. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15667963>.

121. Han J, Noh GT, Yeo SA, et al. The number of retrieved lymph nodes needed for accurate staging differs based on the presence of preoperative chemoradiation for rectal cancer. *Medicine (Baltimore)* 2016;95:e4891. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27661032>.

122. Wichmann MW, Muller C, Meyer G, et al. Effect of preoperative radiochemotherapy on lymph node retrieval after resection of rectal cancer. *Arch Surg* 2002;137:206-210. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11822961>.

123. de Campos-Lobato LF, Stocchi L, de Sousa JB, et al. Less than 12 nodes in the surgical specimen after total mesorectal excision following neoadjuvant chemoradiation: it means more than you think! *Ann Surg Oncol* 2013;20:3398-3406. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23812804>.

124. Kim HJ, Jo JS, Lee SY, et al. Low lymph node retrieval after preoperative chemoradiation for rectal cancer is associated with improved prognosis in patients with a good tumor response. *Ann Surg Oncol* 2015;22:2075-2081. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25395150>.

125. Turner RR, Nora DT, Trocha SD, Bilchik AJ. Colorectal carcinoma nodal staging. Frequency and nature of cytokeratin-positive cells in sentinel and nonsentinel lymph nodes. *Arch Pathol Lab Med*

2003;127:673-679. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12741889>.

126. Wood TF, Nora DT, Morton DL, et al. One hundred consecutive cases of sentinel lymph node mapping in early colorectal carcinoma: detection of missed micrometastases. *J Gastrointest Surg* 2002;6:322-329; discussion 229-330. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12022982>.

127. Yasuda K, Adachi Y, Shiraishi N, et al. Pattern of lymph node micrometastasis and prognosis of patients with colorectal cancer. *Ann Surg Oncol* 2001;8:300-304. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11352302>.

128. Braat AE, Oosterhuis JW, Moll FC, et al. Sentinel node detection after preoperative short-course radiotherapy in rectal carcinoma is not reliable. *Br J Surg* 2005;92:1533-1538. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16231281>.

129. Wiese D, Sirop S, Yestrepky B, et al. Ultrastaging of sentinel lymph nodes (SLNs) vs. non-SLNs in colorectal cancer--do we need both? *Am J Surg* 2010;199:354-358; discussion 358. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20226909>.

130. Noura S, Yamamoto H, Miyake Y, et al. Immunohistochemical assessment of localization and frequency of micrometastases in lymph nodes of colorectal cancer. *Clin Cancer Res* 2002;8:759-767. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11895906>.

131. Sloothaak DA, Sahami S, van der Zaag-Loonen HJ, et al. The prognostic value of micrometastases and isolated tumour cells in histologically negative lymph nodes of patients with colorectal cancer: a systematic review and meta-analysis. *Eur J Surg Oncol* 2014;40:263-269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24368050>.

132. Mescoli C, Albertoni L, Pucciarelli S, et al. Isolated tumor cells in regional lymph nodes as relapse predictors in stage I and II colorectal

cancer. *J Clin Oncol* 2012;30:965-971. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22355061>.

133. Rahbari NN, Bork U, Mutschall E, et al. Molecular detection of tumor cells in regional lymph nodes is associated with disease recurrence and poor survival in node-negative colorectal cancer: a systematic review and meta-analysis. *J Clin Oncol* 2012;30:60-70. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22124103>.

134. Ryan R, Gibbons D, Hyland JM, et al. Pathological response following long-course neoadjuvant chemoradiotherapy for locally advanced rectal cancer. *Histopathology* 2005;47:141-146. Available at:

135. Al-Sukhni E, Attwood K, Gabriel EM, et al. Lymphovascular and perineural invasion are associated with poor prognostic features and outcomes in colorectal cancer: A retrospective cohort study. *Int J Surg* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27600906>.

136. Knijn N, Mogk SC, Teerenstra S, et al. Perineural invasion is a strong prognostic factor in colorectal cancer: a systematic review. *Am J Surg Pathol* 2016;40:103-112. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26426380>.

137. Mayo E, Llanos AA, Yi X, et al. Prognostic value of tumour deposit and perineural invasion status in colorectal cancer patients: a SEER-based population study. *Histopathology* 2016;69:230-238. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26802566>.

138. Yagi R, Shimada Y, Kameyama H, et al. Clinical significance of extramural tumor deposits in the lateral pelvic lymph node area in low rectal cancer: a retrospective study at two institutions. *Ann Surg Oncol* 2016;23:552-558. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27393567>.

139. Gopal P, Lu P, Ayers GD, et al. Tumor deposits in rectal adenocarcinoma after neoadjuvant chemoradiation are associated with poor prognosis. *Mod Pathol* 2014;27:1281-1287. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24434897>.

140. Zhang LN, Xiao WW, Xi SY, et al. Tumor deposits: markers of poor prognosis in patients with locally advanced rectal cancer following neoadjuvant chemoradiotherapy. *Oncotarget* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26695441>.

141. Autier P, Boniol M, Pizot C, Mullie P. Vitamin D status and ill health: a systematic review. *Lancet Diabetes Endocrinol* 2014;2:76-89. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24622671>.

142. Chung M, Lee J, Terasawa T, et al. Vitamin D with or without calcium supplementation for prevention of cancer and fractures: an updated meta-analysis for the U.S. Preventive Services Task Force. *Ann Intern Med* 2011;155:827-838. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22184690>.

143. Gorham ED, Garland CF, Garland FC, et al. Optimal vitamin D status for colorectal cancer prevention: a quantitative meta analysis. *Am J Prev Med* 2007;32:210-216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17296473>.

144. Lappe JM, Travers-Gustafson D, Davies KM, et al. Vitamin D and calcium supplementation reduces cancer risk: results of a randomized trial. *Am J Clin Nutr* 2007;85:1586-1591. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17556697>.

145. Ma Y, Zhang P, Wang F, et al. Association between vitamin D and risk of colorectal cancer: a systematic review of prospective studies. *J Clin Oncol* 2011;29:3775-3782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21876081>.

146. Fedirko V, Riboli E, Tjonneland A, et al. Prediagnostic 25-hydroxyvitamin D, VDR and CASR polymorphisms, and survival in patients with colorectal cancer in western European populations. *Cancer Epidemiol Biomarkers Prev* 2012;21:582-593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22278364>.

147. Ng K, Meyerhardt JA, Wu K, et al. Circulating 25-hydroxyvitamin D levels and survival in patients with colorectal cancer. *J Clin Oncol*



2008;26:2984-2991. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18565885>.

148. Ng K, Venook AP, Sato K, et al. Vitamin D status and survival of metastatic colorectal cancer patients: results from CALGB/SWOG 80405 (Alliance) [abstract]. ASCO Meeting Abstracts 2015;33:3503.

Available at: <http://meetinglibrary.asco.org/content/139861-158>.

149. Zgaga L, Theodoratou E, Farrington SM, et al. Plasma vitamin D concentration influences survival outcome after a diagnosis of colorectal cancer. *J Clin Oncol* 2014;32:2430-2439. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25002714>.

150. Maalmi H, Ordonez-Mena JM, Schottker B, Brenner H. Serum 25-hydroxyvitamin D levels and survival in colorectal and breast cancer patients: systematic review and meta-analysis of prospective cohort studies. *Eur J Cancer* 2014;50:1510-1521. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24582912>.

151. Ou B, Zhao J, Guan S, Lu A. Plasma 25-hydroxyvitamin D levels and survival of colorectal cancer patients: a meta-analysis. *Eur J Cancer* 2015;51:786-788. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25746389>.

152. Baron JA, Barry EL, Mott LA, et al. A trial of calcium and vitamin D for the prevention of colorectal adenomas. *N Engl J Med* 2015;373:1519-1530. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26465985>.

153. Ross AC, Taylor CL, Yaktine AL, Valle HBD, eds. Dietary Reference Intakes for Calcium and Vitamin D. Washington (DC): National Academies Press (US); 2011.

154. Cooper HS, Deppisch LM, Gourley WK, et al. Endoscopically removed malignant colorectal polyps: clinicopathologic correlations. *Gastroenterology* 1995;108:1657-1665. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/7768369>.

155. Hamilton SR, Bosman FT, Boffetta P, et al. Carcinoma of the colon and rectum. In: Bosman FT, Carneiro F, Hruban RH, Theise ND, eds. *WHO Classification of Tumours of the Digestive System*. Lyon: IARC; 2010.

156. Yoshii S, Nojima M, Noshio K, et al. Factors associated with risk for colorectal cancer recurrence after endoscopic resection of T1 tumors. *Clin Gastroenterol Hepatol* 2014;12:292-302 e293. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23962552>.

157. Seitz U, Bohnacker S, Seewald S, et al. Is endoscopic polypectomy an adequate therapy for malignant colorectal adenomas? Presentation of 114 patients and review of the literature. *Dis Colon Rectum* 2004;47:1789-1796. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15622570>.

158. Ueno H, Mochizuki H, Hashiguchi Y, et al. Risk factors for an adverse outcome in early invasive colorectal carcinoma. *Gastroenterology* 2004;127:385-394. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15300569>.

159. Volk EE, Goldblum JR, Petras RE, et al. Management and outcome of patients with invasive carcinoma arising in colorectal polyps. *Gastroenterology* 1995;109:1801-1807. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/7498644>.

160. Choi DH, Sohn DK, Chang HJ, et al. Indications for subsequent surgery after endoscopic resection of submucosally invasive colorectal carcinomas: a prospective cohort study. *Dis Colon Rectum* 2009;52:438-445. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19333043>.

161. Choi JY, Jung SA, Shim KN, et al. Meta-analysis of predictive clinicopathologic factors for lymph node metastasis in patients with early colorectal carcinoma. *J Korean Med Sci* 2015;30:398-406. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25829807>.



162. Park KJ, Choi HJ, Roh MS, et al. Intensity of tumor budding and its prognostic implications in invasive colon carcinoma. *Dis Colon Rectum* 2005;48:1597-1602. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15937624>.

163. Rogers AC, Winter DC, Heeney A, et al. Systematic review and meta-analysis of the impact of tumour budding in colorectal cancer. *Br J Cancer* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27599041>.

164. Garcia-Aguilar J, Hernandez de Anda E, Rothenberger DA, et al. Endorectal ultrasound in the management of patients with malignant rectal polyps. *Dis Colon Rectum* 2005;48:910-916; discussion 916-917. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15868240>.

165. Nelson H, Petrelli N, Carlin A, et al. Guidelines 2000 for colon and rectal cancer surgery. *J Natl Cancer Inst* 2001;93:583-596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11309435>.

166. Kapiteijn E, Marijnen CA, Nagtegaal ID, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer. *N Engl J Med* 2001;345:638-646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11547717>.

167. Schoellhammer HF, Gregorian AC, Sarkisyan GG, Petrie BA. How important is rigid proctosigmoidoscopy in localizing rectal cancer? *Am J Surg* 2008;196:904-908; discussion 908. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19095107>.

168. Baxter NN, Garcia-Aguilar J. Organ preservation for rectal cancer. *J Clin Oncol* 2007;25:1014-1020. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17350952>.

169. Rajput A, Bullard Dunn K. Surgical management of rectal cancer. *Semin Oncol* 2007;34:241-249. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17560986>.

170. Weiser MR, Landmann RG, Wong WD, et al. Surgical salvage of recurrent rectal cancer after transanal excision. *Dis Colon Rectum* 2005;48:1169-1175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15793645>.

171. Wiig JN, Larsen SG, Giercksky KE. Operative treatment of locally recurrent rectal cancer. *Recent Results Cancer Res* 2005;165:136-147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15865028>.

172. Morino M, Risio M, Bach S, et al. Early rectal cancer: the European Association for Endoscopic Surgery (EAES) clinical consensus conference. *Surg Endosc* 2015;29:755-773. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25609317>.

173. Bartram C, Brown G. Endorectal ultrasound and magnetic resonance imaging in rectal cancer staging. *Gastroenterol Clin North Am* 2002;31:827-839. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12481733>.

174. Choi DJ, Kwak JM, Kim J, et al. Preoperative chest computerized tomography in patients with locally advanced mid or lower rectal cancer: its role in staging and impact on treatment strategy. *J Surg Oncol* 2010;102:588-592. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20607759>.

175. Grossmann I, Avenarius JK, Mastboom WJ, Klaase JM. Preoperative staging with chest CT in patients with colorectal carcinoma: not as a routine procedure. *Ann Surg Oncol* 2010;17:2045-2050. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/20151212>.

176. Qiu M, Hu J, Yang D, et al. Pattern of distant metastases in colorectal cancer: a SEER based study. *Oncotarget* 2015;6:38658-38666. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26484417>.

177. Onaitis MW, Petersen RP, Haney JC, et al. Prognostic factors for recurrence after pulmonary resection of colorectal cancer metastases. *Ann Thorac Surg* 2009;87:1684-1688. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19463577>.

178. Bipat S, Glas AS, Slors FJM, et al. Rectal cancer: local staging and assessment of lymph node involvement with endoluminal US, CT, and MR imaging--a meta-analysis. *Radiology* 2004;232:773-783. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15273331>.
179. Klessen C, Rogalla P, Taupitz M. Local staging of rectal cancer: the current role of MRI. *Eur Radiol* 2007;17:379-389. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17008990>.
180. Lahaye MJ, Engelen SM, Nelemans PJ, et al. Imaging for predicting the risk factors--the circumferential resection margin and nodal disease--of local recurrence in rectal cancer: a meta-analysis. *Semin Ultrasound CT MR* 2005;26:259-268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16152740>.
181. Beets-Tan RG, Beets GL. Rectal cancer: review with emphasis on MR imaging. *Radiology* 2004;232:335-346. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15286305>.
182. Xie H, Zhou X, Zhuo Z, et al. Effectiveness of MRI for the assessment of mesorectal fascia involvement in patients with rectal cancer: a systematic review and meta-analysis. *Dig Surg* 2014;31:123-134. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24942675>.
183. Taylor FG, Quirke P, Heald RJ, et al. Preoperative magnetic resonance imaging assessment of circumferential resection margin predicts disease-free survival and local recurrence: 5-year follow-up results of the MERCURY study. *J Clin Oncol* 2014;32:34-43. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24276776>.
184. Beets-Tan RG, Lambregts DM, Maas M, et al. Magnetic resonance imaging for the clinical management of rectal cancer patients: recommendations from the 2012 European Society of Gastrointestinal and Abdominal Radiology (ESGAR) consensus meeting. *Eur Radiol* 2013;23:2522-2531. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23743687>.
185. de Jong EA, Ten Berge JC, Dwarkasing RS, et al. The accuracy of MRI, endorectal ultrasonography, and computed tomography in predicting the response of locally advanced rectal cancer after preoperative therapy: a metaanalysis. *Surgery* 2016;159:688-699. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26619929>.
186. Dickman R, Kundel Y, Levy-Drummer R, et al. Restaging locally advanced rectal cancer by different imaging modalities after preoperative chemoradiation: a comparative study. *Radiat Oncol* 2013;8:278. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24286200>.
187. Guillem JG, Ruby JA, Leibold T, et al. Neither FDG-PET Nor CT can distinguish between a pathological complete response and an incomplete response after neoadjuvant chemoradiation in locally advanced rectal cancer: a prospective study. *Ann Surg* 2013;258:289-295. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23187748>.
188. Hanly AM, Ryan EM, Rogers AC, et al. Multicenter Evaluation of Rectal cancer ReImaging pOst Neoadjuvant (MERRION) therapy. *Ann Surg* 2014;259:723-727. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23744576>.
189. Kuo LJ, Chiou JF, Tai CJ, et al. Can we predict pathologic complete response before surgery for locally advanced rectal cancer treated with preoperative chemoradiation therapy? *Int J Colorectal Dis* 2012;27:613-621. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22080392>.
190. Memon S, Lynch AC, Bressel M, et al. Systematic review and meta-analysis of the accuracy of MRI and endorectal ultrasound in the restaging and response assessment of rectal cancer following neoadjuvant therapy. *Colorectal Dis* 2015;17:748-761. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25891148>.
191. Ryan JE, Warriar SK, Lynch AC, Heriot AG. Assessing pathological complete response to neoadjuvant chemoradiotherapy in locally advanced rectal cancer: a systematic review. *Colorectal Dis*



2015;17:849-861. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26260213>.

192. van der Paardt MP, Zagers MB, Beets-Tan RG, et al. Patients who undergo preoperative chemoradiotherapy for locally advanced rectal cancer restaged by using diagnostic MR imaging: a systematic review and meta-analysis. *Radiology* 2013;269:101-112. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23801777>.

193. Zhao RS, Wang H, Zhou ZY, et al. Restaging of locally advanced rectal cancer with magnetic resonance imaging and endoluminal ultrasound after preoperative chemoradiotherapy: a systemic review and meta-analysis. *Dis Colon Rectum* 2014;57:388-395. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24509465>.

194. Hotker AM, Garcia-Aguilar J, Gollub MJ. Multiparametric MRI of rectal cancer in the assessment of response to therapy: a systematic review. *Dis Colon Rectum* 2014;57:790-799. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24807605>.

195. Joye I, Deroose CM, Vandecaveye V, Haustermans K. The role of diffusion-weighted MRI and (18)F-FDG PET/CT in the prediction of pathologic complete response after radiochemotherapy for rectal cancer: a systematic review. *Radiother Oncol* 2014;113:158-165. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25483833>.

196. Lambregts DM, Rao SX, Sassen S, et al. MRI and diffusion-weighted MRI volumetry for identification of complete tumor responders after preoperative chemoradiotherapy in patients with rectal cancer: a bi-institutional validation study. *Ann Surg* 2015;262:1034-1039. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25211270>.

197. Memon S, Lynch AC, Akhurst T, et al. Systematic review of FDG-PET prediction of complete pathological response and survival in rectal cancer. *Ann Surg Oncol* 2014;21:3598-3607. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24802909>.

198. Guillem JG, Cohen AM. Current issues in colorectal cancer surgery. *Semin Oncol* 1999;26:505-513. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10528898>.

199. Lindsetmo RO, Joh YG, Delaney CP. Surgical treatment for rectal cancer: an international perspective on what the medical gastroenterologist needs to know. *World J Gastroenterol* 2008;14:3281-3289. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18528924>.

200. Willett CG, Compton CC, Shellito PC, Efird JT. Selection factors for local excision or abdominoperineal resection of early stage rectal cancer. *Cancer* 1994;73:2716-2720. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8194011>.

201. Clancy C, Burke JP, Albert MR, et al. Transanal endoscopic microsurgery versus standard transanal excision for the removal of rectal neoplasms: a systematic review and meta-analysis. *Dis Colon Rectum* 2015;58:254-261. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25585086>.

202. Nascimbeni R, Burgart LJ, Nivatvongs S, Larson DR. Risk of lymph node metastasis in T1 carcinoma of the colon and rectum. *Dis Colon Rectum* 2002;45:200-206. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11852333>.

203. Yamamoto S, Watanabe M, Hasegawa H, et al. The risk of lymph node metastasis in T1 colorectal carcinoma. *Hepatogastroenterology* 2004;51:998-1000. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15239233>.

204. You YN, Baxter NN, Stewart A, Nelson H. Is the increasing rate of local excision for stage I rectal cancer in the United States justified?: a nationwide cohort study from the National Cancer Database. *Ann Surg* 2007;245:726-733. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17457165>.

205. Garcia-Aguilar J, Renfro LA, Chow OS, et al. Organ preservation for clinical T2N0 distal rectal cancer using neoadjuvant



chemoradiotherapy and local excision (ACOSOG Z6041): results of an open-label, single-arm, multi-institutional, phase 2 trial. *Lancet Oncol* 2015;16:1537-1546. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26474521>.

206. Shaikh I, Askari A, Ouru S, et al. Oncological outcomes of local excision compared with radical surgery after neoadjuvant chemoradiotherapy for rectal cancer: a systematic review and meta-analysis. *Int J Colorectal Dis* 2015;30:19-29. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25367179>.

207. Landmann RG, Wong WD, Hoepfl J, et al. Limitations of early rectal cancer nodal staging may explain failure after local excision. *Dis Colon Rectum* 2007;50:1520-1525. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17674104>.

208. Kidane B, Chadi SA, Kanters S, et al. Local resection compared with radical resection in the treatment of T1N0M0 rectal adenocarcinoma: a systematic review and meta-analysis. *Dis Colon Rectum* 2015;58:122-140. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25489704>.

209. Nash GM, Weiser MR, Guillem JG, et al. Long-term survival after transanal excision of T1 rectal cancer. *Dis Colon Rectum* 2009;52:577-582. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19404055>.

210. Stitzenberg KB, Sanoff HK, Penn DC, et al. Practice patterns and long-term survival for early-stage rectal cancer. *J Clin Oncol* 2013;31:4276-4282. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24166526>.

211. Sajid MS, Farag S, Leung P, et al. Systematic review and meta-analysis of published trials comparing the effectiveness of transanal endoscopic microsurgery and radical resection in the management of early rectal cancer. *Colorectal Dis* 2014;16:2-14. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24330432>.

212. Lu JY, Lin GL, Qiu HZ, et al. Comparison of transanal endoscopic microsurgery and total mesorectal excision in the treatment of T1 rectal cancer: a meta-analysis. *PLoS One* 2015;10:e0141427. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26505895>.

213. Heald RJ, Husband EM, Ryall RD. The mesorectum in rectal cancer surgery--the clue to pelvic recurrence? *Br J Surg* 1982;69:613-616. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6751457>.

214. Steup WH, Moriya Y, van de Velde CJH. Patterns of lymphatic spread in rectal cancer. A topographical analysis on lymph node metastases. *Eur J Cancer* 2002;38:911-918. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11978516>.

215. Schlag PM. Surgical sphincter preservation in rectal cancer. *Oncologist* 1996;1:288-292. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10388006>.

216. Marr R, Birbeck K, Garvican J, et al. The modern abdominoperineal excision: the next challenge after total mesorectal excision. *Ann Surg* 2005;242:74-82. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15973104>.

217. Russell MM, Ganz PA, Lopa S, et al. Comparative effectiveness of sphincter-sparing surgery versus abdominoperineal resection in rectal cancer: patient-reported outcomes in National Surgical Adjuvant Breast and Bowel Project randomized trial R-04. *Ann Surg* 2015;261:144-148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24670844>.

218. Huang A, Zhao H, Ling T, et al. Oncological superiority of extralevator abdominoperineal resection over conventional abdominoperineal resection: a meta-analysis. *Int J Colorectal Dis* 2014;29:321-327. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24385025>.

219. Nagtegaal ID, van de Velde CJ, van der Worp E, et al. Macroscopic evaluation of rectal cancer resection specimen: clinical significance of the pathologist in quality control. *J Clin Oncol*



2002;20:1729-1734. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11919228>.

220. Parfitt JR, Driman DK. The total mesorectal excision specimen for rectal cancer: a review of its pathological assessment. *J Clin Pathol* 2007;60:849-855. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17046842>.

221. den Dulk M, Putter H, Collette L, et al. The abdominoperineal resection itself is associated with an adverse outcome: the European experience based on a pooled analysis of five European randomised clinical trials on rectal cancer. *Eur J Cancer* 2009;45:1175-1183.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19128956>.

222. Pahlman L, Bohe M, Cedermark B, et al. The Swedish rectal cancer registry. *Br J Surg* 2007;94:1285-1292. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17661309>.

223. Digennaro R, Tondo M, Cuccia F, et al. Coloanal anastomosis or abdominoperineal resection for very low rectal cancer: what will benefit, the surgeon's pride or the patient's quality of life? *Int J Colorectal Dis* 2013;28:949-957. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23274737>.

224. Pachler J, Wille-Jorgensen P. Quality of life after rectal resection for cancer, with or without permanent colostomy. *Cochrane Database Syst Rev* 2012;12:CD004323. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23235607>.

225. Bonjer HJ, Deijen CL, Abis GA, et al. A randomized trial of laparoscopic versus open surgery for rectal cancer. *N Engl J Med* 2015;372:1324-1332. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25830422>.

226. Jeong SY, Park JW, Nam BH, et al. Open versus laparoscopic surgery for mid-rectal or low-rectal cancer after neoadjuvant chemoradiotherapy (COREAN trial): survival outcomes of an open-label, non-inferiority, randomised controlled trial. *Lancet Oncol*

2014;15:767-774. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24837215>.

227. Fleshman J, Branda M, Sargent DJ, et al. Effect of laparoscopic-assisted resection vs open resection of stage II or III rectal cancer on pathologic outcomes: the ACOSOG Z6051 randomized clinical trial. *JAMA* 2015;314:1346-1355. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26441179>.

228. Stevenson AR, Solomon MJ, Lumley JW, et al. Effect of laparoscopic-assisted resection vs open resection on pathological outcomes in rectal cancer: the ALaCaRT randomized clinical trial. *JAMA* 2015;314:1356-1363. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26441180>.

229. Lujan J, Valero G, Biondo S, et al. Laparoscopic versus open surgery for rectal cancer: results of a prospective multicentre analysis of 4,970 patients. *Surg Endosc* 2013;27:295-302. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22736289>.

230. van der Pas MH, Haglund E, Cuesta MA, et al. Laparoscopic versus open surgery for rectal cancer (COLOR II): short-term outcomes of a randomised, phase 3 trial. *Lancet Oncol* 2013;14:210-218.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23395398>.

231. Jayne DG, Guillou PJ, Thorpe H, et al. Randomized trial of laparoscopic-assisted resection of colorectal carcinoma: 3-year results of the UK MRC CLASICC Trial Group. *J Clin Oncol* 2007;25:3061-3068.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17634484>.

232. Jayne DG, Thorpe HC, Copeland J, et al. Five-year follow-up of the Medical Research Council CLASICC trial of laparoscopically assisted versus open surgery for colorectal cancer. *Br J Surg* 2010;97:1638-1645. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20629110>.

233. Kang SB, Park JW, Jeong SY, et al. Open versus laparoscopic surgery for mid or low rectal cancer after neoadjuvant



chemoradiotherapy (COREAN trial): short-term outcomes of an open-label randomised controlled trial. *Lancet Oncol* 2010;11:637-645. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20610322>.

234. Wagman LD. Laparoscopic and open surgery for colorectal cancer: reaching equipoise? *J Clin Oncol* 2007;25:2996-2998. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17634477>.

235. Ahmad NZ, Racheva G, Elmusharaf H. A systematic review and meta-analysis of randomized and non-randomized studies comparing laparoscopic and open abdominoperineal resection for rectal cancer. *Colorectal Dis* 2013;15:269-277. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22958456>.

236. Araujo SE, da Silva eSousa AH, Jr., de Campos FG, et al. Conventional approach x laparoscopic abdominoperineal resection for rectal cancer treatment after neoadjuvant chemoradiation: results of a prospective randomized trial. *Rev Hosp Clin Fac Med Sao Paulo* 2003;58:133-140. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12894309>.

237. Arezzo A, Passera R, Scozzari G, et al. Laparoscopy for rectal cancer reduces short-term mortality and morbidity: results of a systematic review and meta-analysis. *Surg Endosc* 2013;27:1485-1502. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23183871>.

238. Gopall J, Shen XF, Cheng Y. Current status of laparoscopic total mesorectal excision. *Am J Surg* 2012;203:230-241. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22269656>.

239. Kuhry E, Schwenk WF, Gaupset R, et al. Long-term results of laparoscopic colorectal cancer resection. *Cochrane Database Syst Rev* 2008:CD003432. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18425886>.

240. Lee JK, Delaney CP, Lipman JM. Current state of the art in laparoscopic colorectal surgery for cancer: Update on the multi-centric

international trials. *Ann Surg Innov Res* 2012;6:5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22846394>.

241. Jiang JB, Jiang K, Dai Y, et al. Laparoscopic versus open surgery for mid-low rectal cancer: a systematic review and meta-analysis on short- and long-term outcomes. *J Gastrointest Surg* 2015;19:1497-1512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26040854>.

242. Morneau M, Boulanger J, Charlebois P, et al. Laparoscopic versus open surgery for the treatment of colorectal cancer: a literature review and recommendations from the Comité de l'évolution des pratiques en oncologie. *Can J Surg* 2013;56:297-310. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24067514>.

243. Ng SS, Lee JF, Yiu RY, et al. Long-term oncologic outcomes of laparoscopic versus open surgery for rectal cancer: a pooled analysis of 3 randomized controlled trials. *Ann Surg* 2014;259:139-147. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23598381>.

244. Trastulli S, Ciocchi R, Listorti C, et al. Laparoscopic vs open resection for rectal cancer: a meta-analysis of randomized clinical trials. *Colorectal Dis* 2012;14:e277-296. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22330061>.

245. Vennix S, Pelzers L, Bouvy N, et al. Laparoscopic versus open total mesorectal excision for rectal cancer. *Cochrane Database Syst Rev* 2014;4:CD005200. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24737031>.

246. Xiong B, Ma L, Zhang C. Laparoscopic versus open total mesorectal excision for middle and low rectal cancer: a meta-analysis of results of randomized controlled trials. *J Laparoendosc Adv Surg Tech A* 2012;22:674-684. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22881123>.

247. Zhang FW, Zhou ZY, Wang HL, et al. Laparoscopic versus open surgery for rectal cancer: a systematic review and meta-analysis of



randomized controlled trials. Asian Pac J Cancer Prev 2014;15:9985-9996. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25520140>.

248. Zhao D, Li Y, Wang S, Huang Z. Laparoscopic versus open surgery for rectal cancer: a meta-analysis of 3-year follow-up outcomes. Int J Colorectal Dis 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26847617>.

249. Nussbaum DP, Speicher PJ, Ganapathi AM, et al. Laparoscopic versus open low anterior resection for rectal cancer: results from the National Cancer Data Base. J Gastrointest Surg 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25091847>.

250. Miskovic D, Foster J, Agha A, et al. Standardization of laparoscopic total mesorectal excision for rectal cancer: a structured international expert consensus. Ann Surg 2015;261:716-722. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25072446>.

251. Peeters KC, van de Velde CJ, Leer JW, et al. Late side effects of short-course preoperative radiotherapy combined with total mesorectal excision for rectal cancer: increased bowel dysfunction in irradiated patients--a Dutch colorectal cancer group study. J Clin Oncol 2005;23:6199-6206. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16135487>.

252. Rahbari NN, Elbers H, Askoxylakis V, et al. Neoadjuvant radiotherapy for rectal cancer: meta-analysis of randomized controlled trials. Ann Surg Oncol 2013;20:4169-4182. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24002536>.

253. Gunderson LL, Sargent DJ, Tepper JE, et al. Impact of T and N stage and treatment on survival and relapse in adjuvant rectal cancer: a pooled analysis. J Clin Oncol 2004;22:1785-1796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15067027>.

254. Tepper JE, O'Connell M, Niedzwiecki D, et al. Adjuvant therapy in rectal cancer: analysis of stage, sex, and local control--final report of

intergroup 0114. J Clin Oncol 2002;20:1744-1750. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11919230>.

255. Guillem JG, Diaz-Gonzalez JA, Minsky BD, et al. cT3N0 rectal cancer: potential overtreatment with preoperative chemoradiotherapy is warranted. J Clin Oncol 2008;26:368-373. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18202411>.

256. Sauer R, Becker H, Hohenberger W, et al. Preoperative versus postoperative chemoradiotherapy for rectal cancer. N Engl J Med 2004;351:1731-1740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15496622>.

257. Wagman R, Minsky BD, Cohen AM, et al. Sphincter preservation in rectal cancer with preoperative radiation therapy and coloanal anastomosis: long term follow-up. Int J Radiat Oncol Biol Phys 1998;42:51-57. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9747819>.

258. Sauer R, Liersch T, Merkel S, et al. Preoperative versus postoperative chemoradiotherapy for locally advanced rectal cancer: results of the German CAO/ARO/AIO-94 randomized phase III trial after a median follow-up of 11 years. J Clin Oncol 2012;30:1926-1933. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22529255>.

259. Peng LC, Milsom J, Garrett K, et al. Surveillance, epidemiology, and end results-based analysis of the impact of preoperative or postoperative radiotherapy on survival outcomes for T3N0 rectal cancer. Cancer Epidemiol 2014;38:73-78. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24491755>.

260. Kachnic LA. Should preoperative or postoperative therapy be administered in the management of rectal cancer? Semin Oncol 2006;33:S64-69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17178291>.

261. Bujko K, Kepka L, Michalski W, Nowacki MP. Does rectal cancer shrinkage induced by preoperative radio(chemo)therapy increase the



likelihood of anterior resection? A systematic review of randomised trials. *Radiother Oncol* 2006;80:4-12. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16730086>.

262. Wong RK, Tandan V, De Silva S, Figueredo A. Pre-operative radiotherapy and curative surgery for the management of localized rectal carcinoma. *Cochrane Database Syst Rev* 2007;CD002102. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17443515>.

263. Madoff RD. Chemoradiotherapy for rectal cancer--when, why, and how? *N Engl J Med* 2004;351:1790-1792. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15496630>.

264. O'Connell MJ, Martenson JA, Wieand HS, et al. Improving adjuvant therapy for rectal cancer by combining protracted-infusion fluorouracil with radiation therapy after curative surgery. *N Engl J Med* 1994;331:502-507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8041415>.

265. Smalley SR, Benedetti JK, Williamson SK, et al. Phase III trial of fluorouracil-based chemotherapy regimens plus radiotherapy in postoperative adjuvant rectal cancer: GI INT 0144. *J Clin Oncol* 2006;24:3542-3547. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16877719>.

266. Gerard JP, Conroy T, Bonnetain F, et al. Preoperative radiotherapy with or without concurrent fluorouracil and leucovorin in T3-4 rectal cancers: results of FFCD 9203. *J Clin Oncol* 2006;24:4620-4625. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17008704>.

267. Bosset JF, Calais G, Mineur L, et al. Enhanced tumorocidal effect of chemotherapy with preoperative radiotherapy for rectal cancer: preliminary results--EORTC 22921. *J Clin Oncol* 2005;23:5620-5627. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16009958>.

268. Bosset JF, Collette L, Calais G, et al. Chemotherapy with preoperative radiotherapy in rectal cancer. *N Engl J Med*

2006;355:1114-1123. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16971718>.

269. Ceelen WP, Van Nieuwenhove Y, Fierens K. Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer. *Cochrane Database Syst Rev* 2009;CD006041. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19160264>.

270. McCarthy K, Pearson K, Fulton R, Hewitt J. Pre-operative chemoradiation for non-metastatic locally advanced rectal cancer. *Cochrane Database Syst Rev* 2012;12:CD008368. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23235660>.

271. De Caluwe L, Van Nieuwenhove Y, Ceelen WP. Preoperative chemoradiation versus radiation alone for stage II and III resectable rectal cancer. *Cochrane Database Syst Rev* 2013;2:CD006041. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23450565>.

272. Hofheinz RD, Wenz F, Post S, et al. Chemoradiotherapy with capecitabine versus fluorouracil for locally advanced rectal cancer: a randomised, multicentre, non-inferiority, phase 3 trial. *Lancet Oncol* 2012;13:579-588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22503032>.

273. O'Connell MJ, Colangelo LH, Beart RW, et al. Capecitabine and oxaliplatin in the preoperative multimodality treatment of rectal cancer: surgical end points from National Surgical Adjuvant Breast and Bowel Project trial R-04. *J Clin Oncol* 2014;32:1927-1934. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24799484>.

274. Allegra CJ, Yothers G, O'Connell MJ, et al. Neoadjuvant 5-FU or capecitabine plus radiation with or without oxaliplatin in rectal cancer patients: a phase III randomized clinical trial. *J Natl Cancer Inst* 2015;107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26374429>.

275. Aschele C, Cionini L, Lonardi S, et al. Primary tumor response to preoperative chemoradiation with or without oxaliplatin in locally advanced rectal cancer: pathologic results of the STAR-01 randomized



phase III trial. *J Clin Oncol* 2011;29:2773-2780. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21606427>.

276. Gerard JP, Azria D, Gourgou-Bourgade S, et al. Clinical outcome of the ACCORD 12/0405 PRODIGE 2 randomized trial in rectal cancer. *J Clin Oncol* 2012;30:4558-4565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23109696>.

277. Rodel C, Graeven U, Fietkau R, et al. Oxaliplatin added to fluorouracil-based preoperative chemoradiotherapy and postoperative chemotherapy of locally advanced rectal cancer (the German CAO/ARO/AIO-04 study): final results of the multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol* 2015;16:979-989. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26189067>.

278. Rodel C, Liersch T, Becker H, et al. Preoperative chemoradiotherapy and postoperative chemotherapy with fluorouracil and oxaliplatin versus fluorouracil alone in locally advanced rectal cancer: initial results of the German CAO/ARO/AIO-04 randomised phase 3 trial. *Lancet Oncol* 2012;13:679-687. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22627104>.

279. Glynne-Jones R. Rectal cancer--the times they are a-changing. *Lancet Oncol* 2012;13:651-653. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22627103>.

280. Deng Y, Chi P, Lan P, et al. Modified FOLFOX6 with or without radiation versus fluorouracil and leucovorin with radiation in neoadjuvant treatment of locally advanced rectal cancer: initial results of the Chinese FOWARC multicenter, open-label, randomized three-arm phase III trial. *J Clin Oncol* 2016;34:3300-3307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27480145>.

281. Feng YR, Zhu Y, Liu LY, et al. Interim analysis of postoperative chemoradiotherapy with capecitabine and oxaliplatin versus capecitabine alone for pathological stage II and III rectal cancer: a randomized multicenter phase III trial. *Oncotarget* 2016;7:25576-25584. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27014909>.

282. Dewdney A, Cunningham D, Taberero J, et al. Multicenter randomized phase II clinical trial comparing neoadjuvant oxaliplatin, capecitabine, and preoperative radiotherapy with or without cetuximab followed by total mesorectal excision in patients with high-risk rectal cancer (EXPERT-C). *J Clin Oncol* 2012;30:1620-1627. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22473163>.

283. Eisterer W, De Vries A, Ofner D, et al. Preoperative treatment with capecitabine, cetuximab and radiotherapy for primary locally advanced rectal cancer--a phase II clinical trial. *Anticancer Res* 2014;34:6767-6773. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25368289>.

284. Kripp M, Horisberger K, Mai S, et al. Does the addition of cetuximab to radiochemotherapy improve outcome of patients with locally advanced rectal cancer? Long-term results from phase II trials. *Gastroenterol Res Pract* 2015;2015:273489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25861256>.

285. Helbling D, Bodoky G, Gautschi O, et al. Neoadjuvant chemoradiotherapy with or without panitumumab in patients with wild-type KRAS, locally advanced rectal cancer (LARC): a randomized, multicenter, phase II trial SAKK 41/07. *Ann Oncol* 2013;24:718-725. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23139259>.

286. Landry JC, Feng Y, Prabhu RS, et al. Phase II trial of preoperative radiation with concurrent capecitabine, oxaliplatin, and bevacizumab followed by surgery and postoperative 5-fluorouracil, leucovorin, oxaliplatin (FOLFOX), and bevacizumab in patients with locally advanced rectal cancer: 5-year clinical outcomes ECOG-ACRIN Cancer Research Group E3204. *Oncologist* 2015;20:615-616. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25926352>.

287. Chiorean EG, Sanghani S, Schiel MA, et al. Phase II and gene expression analysis trial of neoadjuvant capecitabine plus irinotecan followed by capecitabine-based chemoradiotherapy for locally advanced rectal cancer: Hoosier Oncology Group GI03-53. *Cancer Chemother Pharmacol* 2012;70:25-32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22610353>.



288. Kim SY, Hong YS, Kim DY, et al. Preoperative chemoradiation with cetuximab, irinotecan, and capecitabine in patients with locally advanced resectable rectal cancer: a multicenter Phase II study. *Int J Radiat Oncol Biol Phys* 2011;81:677-683. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20888703>.

289. Spigel DR, Bendell JC, McCleod M, et al. Phase II study of bevacizumab and chemoradiation in the preoperative or adjuvant treatment of patients with stage II/III rectal cancer. *Clin Colorectal Cancer* 2012;11:45-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21840771>.

290. Cercek A, Goodman KA, Hajj C, et al. Neoadjuvant chemotherapy first, followed by chemoradiation and then surgery, in the management of locally advanced rectal cancer. *J Natl Compr Canc Netw* 2014;12:513-519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24717570>.

291. Chau I, Brown G, Cunningham D, et al. Neoadjuvant capecitabine and oxaliplatin followed by synchronous chemoradiation and total mesorectal excision in magnetic resonance imaging-defined poor-risk rectal cancer. *J Clin Oncol* 2006;24:668-674. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16446339>.

292. Fernandez-Martos C, Pericay C, Aparicio J, et al. Phase II, randomized study of concomitant chemoradiotherapy followed by surgery and adjuvant capecitabine plus oxaliplatin (CAPOX) compared with induction CAPOX followed by concomitant chemoradiotherapy and surgery in magnetic resonance imaging-defined, locally advanced rectal cancer: Grupo cancer de recto 3 study. *J Clin Oncol* 2010;28:859-865. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20065174>.

293. Perez K, Safran H, Sikov W, et al. Complete neoadjuvant treatment for rectal cancer: The Brown University Oncology Group CONTRE study. *Am J Clin Oncol* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25374145>.

294. Marechal R, Vos B, Polus M, et al. Short course chemotherapy followed by concomitant chemoradiotherapy and surgery in locally advanced rectal cancer: a randomized multicentric phase II study. *Ann Oncol* 2012;23:1525-1530. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22039087>.

295. Nogue M, Salud A, Vicente P, et al. Addition of bevacizumab to XELOX induction therapy plus concomitant capecitabine-based chemoradiotherapy in magnetic resonance imaging-defined poor-prognosis locally advanced rectal cancer: the AVACROSS study. *Oncologist* 2011;16:614-620. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21467148>.

296. Fernandez-Martos C, Garcia-Albeniz X, Pericay C, et al. Chemoradiation, surgery and adjuvant chemotherapy versus induction chemotherapy followed by chemoradiation and surgery: long-term results of the Spanish GCR-3 phase II randomized trial dagger. *Ann Oncol* 2015;26:1722-1728. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25957330>.

297. Schrag D, Weiser MR, Goodman KA, et al. Neoadjuvant chemotherapy without routine use of radiation therapy for patients with locally advanced rectal cancer: a pilot trial. *J Clin Oncol* 2014;32:513-518. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24419115>.

298. Jalil O, Claydon L, Arulampalam T. Review of neoadjuvant chemotherapy alone in locally advanced rectal cancer. *J Gastrointest Cancer* 2015;46:219-236. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26133151>.

299. Gay HA, Barthold HJ, O'Meara E, et al. Pelvic normal tissue contouring guidelines for radiation therapy: a radiation therapy oncology group consensus panel atlas. *Int J Radiat Oncol Biol Phys* 2012;83:e353-362. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22483697>.

300. Francois Y, Nemoz CJ, Baulieux J, et al. Influence of the interval between preoperative radiation therapy and surgery on downstaging

and on the rate of sphincter-sparing surgery for rectal cancer: the Lyon R90-01 randomized trial. *J Clin Oncol* 1999;17:2396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10561302>.

301. Habr-Gama A, Perez RO, Proscurshim I, et al. Interval between surgery and neoadjuvant chemoradiation therapy for distal rectal cancer: does delayed surgery have an impact on outcome? *Int J Radiat Oncol Biol Phys* 2008;71:1181-1188. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18234443>.

302. Moore HG, Gittleman AE, Minsky BD, et al. Rate of pathologic complete response with increased interval between preoperative combined modality therapy and rectal cancer resection. *Dis Colon Rectum* 2004;47:279-286. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14991488>.

303. Petrelli F, Sgroi G, Sarti E, Barni S. Increasing the interval between neoadjuvant chemoradiotherapy and surgery in rectal cancer: a meta-analysis of published studies. *Ann Surg* 2016;263:458-464. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24263329>.

304. Sloothaak DA, Geijssen DE, van Leersum NJ, et al. Optimal time interval between neoadjuvant chemoradiotherapy and surgery for rectal cancer. *Br J Surg* 2013;100:933-939. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23536485>.

305. Tulchinsky H, Shmueli E, Figer A, et al. An interval >7 weeks between neoadjuvant therapy and surgery improves pathologic complete response and disease-free survival in patients with locally advanced rectal cancer. *Ann Surg Oncol* 2008;15:2661-2667. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18389322>.

306. Probst CP, Becerra AZ, Aquina CT, et al. Extended intervals after neoadjuvant therapy in locally advanced rectal cancer: the key to improved tumor response and potential organ preservation. *J Am Coll Surg* 2015;221:430-440. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26206642>.

307. Huntington CR, Boselli D, Symanowski J, et al. Optimal timing of surgical resection after radiation in locally advanced rectal adenocarcinoma: an analysis of the National Cancer Database. *Ann Surg Oncol* 2016;23:877-887. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26514119>.

308. Sun Z, Adam MA, Kim J, et al. Optimal timing to surgery after neoadjuvant chemoradiotherapy for locally advanced rectal cancer. *J Am Coll Surg* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26897480>.

309. Lefevre JH, Mineur L, Kotti S, et al. Effect of interval (7 or 11 weeks) between neoadjuvant radiochemotherapy and surgery on complete pathologic response in rectal cancer: a multicenter, randomized, controlled trial (GRECCAR-6). *J Clin Oncol* 2016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27432930>.

310. Improved survival with preoperative radiotherapy in resectable rectal cancer. Swedish Rectal Cancer Trial. *N Engl J Med* 1997;336:980-987. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9091798>.

311. Birgisson H, Pahlman L, Gunnarsson U, Glimelius B. Adverse effects of preoperative radiation therapy for rectal cancer: long-term follow-up of the Swedish Rectal Cancer Trial. *J Clin Oncol* 2005;23:8697-8705. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16314629>.

312. Peeters KCMJ, Marijnen CAM, Nagtegaal ID, et al. The TME trial after a median follow-up of 6 years: increased local control but no survival benefit in irradiated patients with resectable rectal carcinoma. *Ann Surg* 2007;246:693-701. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17968156>.

313. Siegel R, Burock S, Wernecke KD, et al. Preoperative short-course radiotherapy versus combined radiochemotherapy in locally advanced rectal cancer: a multi-centre prospectively randomised study of the

Berlin Cancer Society. BMC Cancer 2009;9:50. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19200365>.

314. Sebag-Montefiore D, Stephens RJ, Steele R, et al. Preoperative radiotherapy versus selective postoperative chemoradiotherapy in patients with rectal cancer (MRC CR07 and NCIC-CTG C016): a multicentre, randomised trial. Lancet 2009;373:811-820. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19269519>.

315. Stephens RJ, Thompson LC, Quirke P, et al. Impact of short-course preoperative radiotherapy for rectal cancer on patients' quality of life: data from the Medical Research Council CR07/National Cancer Institute of Canada Clinical Trials Group C016 randomized clinical trial. J Clin Oncol 2010;28:4233-4239. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/20585099>.

316. van Gijn W, Marijnen CA, Nagtegaal ID, et al. Preoperative radiotherapy combined with total mesorectal excision for resectable rectal cancer: 12-year follow-up of the multicentre, randomised controlled TME trial. Lancet Oncol 2011;12:575-582. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21596621>.

317. Bujko K, Nowacki MP, Nasierowska-Guttmejer A, et al. Long-term results of a randomized trial comparing preoperative short-course radiotherapy with preoperative conventionally fractionated chemoradiation for rectal cancer. Br J Surg 2006;93:1215-1223. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16983741>.

318. Ngan SY, Burmeister B, Fisher RJ, et al. Randomized trial of short-course radiotherapy versus long-course chemoradiation comparing rates of local recurrence in patients with T3 rectal cancer: Trans-Tasman Radiation Oncology Group trial 01.04. J Clin Oncol 2012;30:3827-3833. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23008301>.

319. Ansari N, Solomon MJ, Fisher RJ, et al. Acute adverse events and postoperative complications in a randomized trial of preoperative short-course radiotherapy versus long-course chemoradiotherapy for T3

adenocarcinoma of the rectum: Trans-Tasman Radiation Oncology Group trial (TROG 01.04). Ann Surg 2016. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27631775>.

320. Latkauskas T, Pauzas H, Gineikiene I, et al. Initial results of a randomized controlled trial comparing clinical and pathological downstaging of rectal cancer after preoperative short-course radiotherapy or long-term chemoradiotherapy, both with delayed surgery. Colorectal Dis 2012;14:294-298. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/21899712>.

321. Bujko K, Partycki M, Pietrzak L. Neoadjuvant radiotherapy (5 x 5 Gy): immediate versus delayed surgery. Recent Results Cancer Res 2014;203:171-187. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25103005>.

322. Collette L, Bosset J-F, den Dulk M, et al. Patients with curative resection of cT3-4 rectal cancer after preoperative radiotherapy or radiochemotherapy: does anybody benefit from adjuvant fluorouracil-based chemotherapy? A trial of the European Organisation for Research and Treatment of Cancer Radiation Oncology Group. J Clin Oncol 2007;25:4379-4386. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17906203>.

323. Das P, Skibber JM, Rodriguez-Bigas MA, et al. Clinical and pathologic predictors of locoregional recurrence, distant metastasis, and overall survival in patients treated with chemoradiation and mesorectal excision for rectal cancer. Am J Clin Oncol 2006;29:219-224. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16755173>.

324. Das P, Skibber JM, Rodriguez-Bigas MA, et al. Predictors of tumor response and downstaging in patients who receive preoperative chemoradiation for rectal cancer. Cancer 2007;109:1750-1755. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17387743>.

325. Fietkau R, Barten M, Klautke G, et al. Postoperative chemotherapy may not be necessary for patients with ypN0-category after neoadjuvant



chemoradiotherapy of rectal cancer. *Dis Colon Rectum* 2006;49:1284-1292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16758130>.

326. Park IJ, You YN, Agarwal A, et al. Neoadjuvant treatment response as an early response indicator for patients with rectal cancer. *J Clin Oncol* 2012;30:1770-1776. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22493423>.

327. Silberfein EJ, Kattepogu KM, Hu CY, et al. Long-term survival and recurrence outcomes following surgery for distal rectal cancer. *Ann Surg Oncol* 2010;17:2863-2869. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20552409>.

328. Smith KD, Tan D, Das P, et al. Clinical significance of acellular mucin in rectal adenocarcinoma patients with a pathologic complete response to preoperative chemoradiation. *Ann Surg* 2010;251:261-264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19864936>.

329. Patel UB, Taylor F, Blomqvist L, et al. Magnetic resonance imaging-detected tumor response for locally advanced rectal cancer predicts survival outcomes: MERCURY experience. *J Clin Oncol* 2011;29:3753-3760. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21876084>.

330. Fokas E, Liersch T, Fietkau R, et al. Tumor regression grading after preoperative chemoradiotherapy for locally advanced rectal carcinoma revisited: updated results of the CAO/ARO/AIO-94 trial. *J Clin Oncol* 2014;32:1554-1562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24752056>.

331. Janjan NA, Crane C, Feig BW, et al. Improved overall survival among responders to preoperative chemoradiation for locally advanced rectal cancer. *Am J Clin Oncol* 2001;24:107-112. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11319280>.

332. Habr-Gama A, Perez RO, Nadalin W, et al. Operative versus nonoperative treatment for stage 0 distal rectal cancer following chemoradiation therapy: long-term results. *Ann Surg* 2004;240:711-717;

discussion 717-718. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15383798>.

333. Glynne-Jones R, Wallace M, Livingstone JI, Meyrick-Thomas J. Complete clinical response after preoperative chemoradiation in rectal cancer: is a "wait and see" policy justified? *Dis Colon Rectum* 2008;51:10-19; discussion 19-20. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18043968>.

334. Maas M, Beets-Tan RG, Lambregts DM, et al. Wait-and-see policy for clinical complete responders after chemoradiation for rectal cancer. *J Clin Oncol* 2011;29:4633-4640. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22067400>.

335. Appelt AL, Ploen J, Harling H, et al. High-dose chemoradiotherapy and watchful waiting for distal rectal cancer: a prospective observational study. *Lancet Oncol* 2015;16:919-927. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26156652>.

336. Habr-Gama A, Gama-Rodrigues J, Sao Juliao GP, et al. Local recurrence after complete clinical response and watch and wait in rectal cancer after neoadjuvant chemoradiation: impact of salvage therapy on local disease control. *Int J Radiat Oncol Biol Phys* 2014;88:822-828. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24495589>.

337. Li J, Liu H, Yin J, et al. Wait-and-see or radical surgery for rectal cancer patients with a clinical complete response after neoadjuvant chemoradiotherapy: a cohort study. *Oncotarget* 2015;6:42354-42361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26472284>.

338. Renehan AG, Malcomson L, Emsley R, et al. Watch-and-wait approach versus surgical resection after chemoradiotherapy for patients with rectal cancer (the OnCoRe project): a propensity-score matched cohort analysis. *Lancet Oncol* 2016;17:174-183. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26705854>.

339. Glynne-Jones R, Hughes R. Critical appraisal of the 'wait and see' approach in rectal cancer for clinical complete responders after

chemoradiation. Br J Surg 2012;99:897-909. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22539154>.

340. Nahas SC, Rizkallah Nahas CS, Sparapan Marques CF, et al. Pathologic complete response in rectal cancer: can we detect it? Lessons learned from a proposed randomized trial of watch-and-wait treatment of rectal cancer. Dis Colon Rectum 2016;59:255-263. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26953983>.

341. Tranchart H, Lefevre JH, Svrcek M, et al. What is the incidence of metastatic lymph node involvement after significant pathologic response of primary tumor following neoadjuvant treatment for locally advanced rectal cancer? Ann Surg Oncol 2013;20:1551-1559. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23188545>.

342. Bujko K, Glimelius B, Valentini V, et al. Postoperative chemotherapy in patients with rectal cancer receiving preoperative radio(chemo)therapy: A meta-analysis of randomized trials comparing surgery +/- a fluoropyrimidine and surgery + a fluoropyrimidine +/- oxaliplatin. Eur J Surg Oncol 2015;41:713-723. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25911110>.

343. Wolmark N, Wieand HS, Hyams DM, et al. Randomized trial of postoperative adjuvant chemotherapy with or without radiotherapy for carcinoma of the rectum: National Surgical Adjuvant Breast and Bowel Project Protocol R-02. J Natl Cancer Inst 2000;92:388-396. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10699069>.

344. Bosset JF, Calais G, Mineur L, et al. Fluorouracil-based adjuvant chemotherapy after preoperative chemoradiotherapy in rectal cancer: long-term results of the EORTC 22921 randomised study. Lancet Oncol 2014;15:184-190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24440473>.

345. Sainato A, Cernusco Luna Nunzia V, Valentini V, et al. No benefit of adjuvant fluorouracil leucovorin chemotherapy after neoadjuvant chemoradiotherapy in locally advanced cancer of the rectum (LARC): long term results of a randomized trial (I-CNR-RT). Radiother Oncol

2014;113:223-229. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25454175>.

346. Breugom AJ, van Gijn W, Muller EW, et al. Adjuvant chemotherapy for rectal cancer patients treated with preoperative (chemo)radiotherapy and total mesorectal excision: a Dutch Colorectal Cancer Group (DCCG) randomized phase III trial. Ann Oncol 2015;26:696-701. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25480874>.

347. Benson AB, Catalan P, Meropol NJ, et al. ECOG E3201: Intergroup randomized phase III study of postoperative irinotecan, 5-fluorouracil (FU), leucovorin (LV) (FOLFIRI) vs oxaliplatin, FU/LV (FOLFOX) vs FU/LV for patients (pts) with stage II/ III rectal cancer receiving either pre or postoperative radiation (RT)/ FU [abstract]. J Clin Oncol 2006;24 (June 20 suppl):3526. Available at:

348. Hong YS, Nam BH, Kim KP, et al. Oxaliplatin, fluorouracil, and leucovorin versus fluorouracil and leucovorin as adjuvant chemotherapy for locally advanced rectal cancer after preoperative chemoradiotherapy (ADORE): an open-label, multicentre, phase 2, randomised controlled trial. Lancet Oncol 2014;15:1245-1253. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25201358>.

349. Garcia-Albeniz X, Gallego R, Hofheinz RD, et al. Adjuvant therapy sparing in rectal cancer achieving complete response after chemoradiation. World J Gastroenterol 2014;20:15820-15829. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25400468>.

350. Breugom AJ, Swets M, Bosset JF, et al. Adjuvant chemotherapy after preoperative (chemo)radiotherapy and surgery for patients with rectal cancer: a systematic review and meta-analysis of individual patient data. Lancet Oncol 2015;16:200-207. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25589192>.

351. Petersen SH, Harling H, Kirkeby LT, et al. Postoperative adjuvant chemotherapy in rectal cancer operated for cure. Cochrane Database Syst Rev 2012;3:CD004078. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22419291>.



352. Petrelli F, Coinu A, Lonati V, Barni S. A systematic review and meta-analysis of adjuvant chemotherapy after neoadjuvant treatment and surgery for rectal cancer. *Int J Colorectal Dis* 2015;30:447-457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25433820>.

353. Khrizman P, Niland JC, ter Veer A, et al. Postoperative adjuvant chemotherapy use in patients with stage II/III rectal cancer treated with neoadjuvant therapy: a national comprehensive cancer network analysis. *J Clin Oncol* 2013;31:30-38. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23169502>.

354. Haynes AB, You YN, Hu CY, et al. Postoperative chemotherapy use after neoadjuvant chemoradiotherapy for rectal cancer: Analysis of Surveillance, Epidemiology, and End Results-Medicare data, 1998-2007. *Cancer* 2014;120:1162-1170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24474245>.

355. Biagi JJ, Raphael MJ, Mackillop WJ, et al. Association between time to initiation of adjuvant chemotherapy and survival in colorectal cancer: a systematic review and meta-analysis. *JAMA* 2011;305:2335-2342. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21642686>.

356. Des Guetz G, Nicolas P, Perret GY, et al. Does delaying adjuvant chemotherapy after curative surgery for colorectal cancer impair survival? A meta-analysis. *Eur J Cancer* 2010;46:1049-1055. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20138505>.

357. Fakih M. Treating rectal cancer: key issues reconsidered. *Oncology (Williston Park)* 2008;22:1444-1446. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19322952>.

358. Minsky BD, Guillem JG. Multidisciplinary management of resectable rectal cancer. New developments and controversies. *Oncology (Williston Park)* 2008;22:1430-1437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19086601>.

359. Andre T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med*

2004;350:2343-2351. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15175436>.

360. Benson AB, 3rd, Hamilton SR. Path toward prognostication and prediction: an evolving matrix. *J Clin Oncol* 2011;29:4599-4601. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22067398>.

361. O'Connell MJ, Lavery I, Yothers G, et al. Relationship between tumor gene expression and recurrence in four independent studies of patients with stage II/III colon cancer treated with surgery alone or surgery plus adjuvant fluorouracil plus leucovorin. *J Clin Oncol* 2010;28:3937-3944. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20679606>.

362. Gray RG, Quirke P, Handley K, et al. Validation study of a quantitative multigene reverse transcriptase-polymerase chain reaction assay for assessment of recurrence risk in patients with stage II colon cancer. *J Clin Oncol* 2011;29:4611-4619. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22067390>.

363. Venook AP, Niedzwiecki D, Lopatin M, et al. Biologic determinants of tumor recurrence in stage II colon cancer: validation study of the 12-gene recurrence score in cancer and leukemia group B (CALGB) 9581. *J Clin Oncol* 2013;31:1775-1781. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23530100>.

364. Yothers G, O'Connell MJ, Lee M, et al. Validation of the 12-gene colon cancer recurrence score in NSABP C-07 as a predictor of recurrence in patients with stage II and III colon cancer treated with fluorouracil and leucovorin (FU/LV) and FU/LV plus oxaliplatin. *J Clin Oncol* 2013;31:4512-4519. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24220557>.

365. Reimers MS, Kuppen PJ, Lee M, et al. Validation of the 12-gene colon cancer recurrence score as a predictor of recurrence risk in stage II and III rectal cancer patients. *J Natl Cancer Inst* 2014;106. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25261968>.



366. Comparison of fluorouracil with additional levamisole, higher-dose folinic acid, or both, as adjuvant chemotherapy for colorectal cancer: a randomised trial. QUASAR Collaborative Group. *Lancet* 2000;355:1588-1596. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10821362>.

367. Jager E, Heike M, Bernhard H, et al. Weekly high-dose leucovorin versus low-dose leucovorin combined with fluorouracil in advanced colorectal cancer: results of a randomized multicenter trial. Study Group for Palliative Treatment of Metastatic Colorectal Cancer Study Protocol 1. *J Clin Oncol* 1996;14:2274-2279. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8708717>.

368. O'Connell MJ. A phase III trial of 5-fluorouracil and leucovorin in the treatment of advanced colorectal cancer. A Mayo Clinic/North Central Cancer Treatment Group study. *Cancer* 1989;63:1026-1030. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2465076>.

369. Garcia-Aguilar J, Mellgren A, Sirivongs P, et al. Local excision of rectal cancer without adjuvant therapy: a word of caution. *Ann Surg* 2000;231:345-351. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10714627>.

370. Sengupta S, Tjandra JJ. Local excision of rectal cancer: what is the evidence? *Dis Colon Rectum* 2001;44:1345-1361. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11584215>.

371. Schmoll HJ, Cartwright T, Tabernero J, et al. Phase III trial of capecitabine plus oxaliplatin as adjuvant therapy for stage III colon cancer: a planned safety analysis in 1,864 patients. *J Clin Oncol* 2007;25:102-109. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17194911>.

372. Alberda WJ, Verhoef C, Nuyttens JJ, et al. Intraoperative radiation therapy reduces local recurrence rates in patients with microscopically involved circumferential resection margins after resection of locally advanced rectal cancer. *Int J Radiat Oncol Biol Phys* 2014;88:1032-1040. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24661656>.

373. Hahnloser D, Haddock MG, Nelson H. Intraoperative radiotherapy in the multimodality approach to colorectal cancer. *Surg Oncol Clin N Am* 2003;12:993-1013, ix. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14989129>.

374. Hyngstrom JR, Tzeng CW, Beddar S, et al. Intraoperative radiation therapy for locally advanced primary and recurrent colorectal cancer: ten-year institutional experience. *J Surg Oncol* 2014;109:652-658. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24510523>.

375. Valentini V, Balducci M, Tortoreto F, et al. Intraoperative radiotherapy: current thinking. *Eur J Surg Oncol* 2002;28:180-185. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11884054>.

376. Willett CG, Czito BG, Tyler DS. Intraoperative radiation therapy. *J Clin Oncol* 2007;25:971-977. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17350946>.

377. Lee WS, Yun SH, Chun HK, et al. Pulmonary resection for metastases from colorectal cancer: prognostic factors and survival. *Int J Colorectal Dis* 2007;22:699-704. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17109105>.

378. Van Cutsem E, Nordlinger B, Adam R, et al. Towards a pan-European consensus on the treatment of patients with colorectal liver metastases. *Eur J Cancer* 2006;42:2212-2221. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16904315>.

379. Yoo PS, Lopez-Soler RI, Longo WE, Cha CH. Liver resection for metastatic colorectal cancer in the age of neoadjuvant chemotherapy and bevacizumab. *Clin Colorectal Cancer* 2006;6:202-207. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17026789>.

380. Alberts SR, Horvath WL, Sternfeld WC, et al. Oxaliplatin, fluorouracil, and leucovorin for patients with unresectable liver-only metastases from colorectal cancer: a North Central Cancer Treatment Group phase II study. *J Clin Oncol* 2005;23:9243-9249. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16230673>.



381. Dawood O, Mahadevan A, Goodman KA. Stereotactic body radiation therapy for liver metastases. *Eur J Cancer* 2009;45:2947-2959. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19773153>.

382. Kemeny N. Management of liver metastases from colorectal cancer. *Oncology (Williston Park)* 2006;20:1161-1176, 1179; discussion 1179-1180, 1185-1166. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17024869>.

383. Muratore A, Zorzi D, Bouzari H, et al. Asymptomatic colorectal cancer with un-resectable liver metastases: immediate colorectal resection or up-front systemic chemotherapy? *Ann Surg Oncol* 2007;14:766-770. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17103261>.

384. Fong Y, Cohen AM, Fortner JG, et al. Liver resection for colorectal metastases. *J Clin Oncol* 1997;15:938-946. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9060531>.

385. Hayashi M, Inoue Y, Komeda K, et al. Clinicopathological analysis of recurrence patterns and prognostic factors for survival after hepatectomy for colorectal liver metastasis. *BMC Surg* 2010;10:27. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20875094>.

386. Tsai M-S, Su Y-H, Ho M-C, et al. Clinicopathological features and prognosis in resectable synchronous and metachronous colorectal liver metastasis. *Ann Surg Oncol* 2007;14:786-794. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17103254>.

387. Foster JH. Treatment of metastatic disease of the liver: a skeptic's view. *Semin Liver Dis* 1984;4:170-179. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6205450>.

388. Stangl R, Altendorf-Hofmann A, Charnley RM, Scheele J. Factors influencing the natural history of colorectal liver metastases. *Lancet* 1994;343:1405-1410. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7515134>.

389. Adam R, Delvart V, Pascal G, et al. Rescue surgery for unresectable colorectal liver metastases downstaged by chemotherapy: a model to predict long-term survival. *Ann Surg* 2004;240:644-657; discussion 657-648. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15383792>.

390. Choti MA, Sitzmann JV, Tiburi MF, et al. Trends in long-term survival following liver resection for hepatic colorectal metastases. *Ann Surg* 2002;235:759-766. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12035031>.

391. Elias D, Liberale G, Vernerey D, et al. Hepatic and extrahepatic colorectal metastases: when resectable, their localization does not matter, but their total number has a prognostic effect. *Ann Surg Oncol* 2005;12:900-909. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16184442>.

392. Fong Y, Salo J. Surgical therapy of hepatic colorectal metastasis. *Semin Oncol* 1999;26:514-523. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10528899>.

393. Pawlik TM, Scoggins CR, Zorzi D, et al. Effect of surgical margin status on survival and site of recurrence after hepatic resection for colorectal metastases. *Ann Surg* 2005;241:715-722, discussion 722-714. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15849507>.

394. Van Cutsem E, Cervantes A, Adam R, et al. ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. *Ann Oncol* 2016;27:1386-1422. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27380959>.

395. Venook AP. The Kemeny article reviewed: management of liver metastases from colorectal cancer: review 2. *Oncology* 2006;20. Available at: <http://www.cancernetwork.com/display/article/10165/108033>.

396. Kanas GP, Taylor A, Primrose JN, et al. Survival after liver resection in metastatic colorectal cancer: review and meta-analysis of



prognostic factors. Clin Epidemiol 2012;4:283-301. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23152705>.

397. Aloia TA, Vauthey JN, Loyer EM, et al. Solitary colorectal liver metastasis: resection determines outcome. Arch Surg 2006;141:460-466; discussion 466-467. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16702517>.

398. Hur H, Ko YT, Min BS, et al. Comparative study of resection and radiofrequency ablation in the treatment of solitary colorectal liver metastases. Am J Surg 2009;197:728-736. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18789428>.

399. Lee WS, Yun SH, Chun HK, et al. Clinical outcomes of hepatic resection and radiofrequency ablation in patients with solitary colorectal liver metastasis. J Clin Gastroenterol 2008;42:945-949. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18438208>.

400. Charnsangavej C, Clary B, Fong Y, et al. Selection of patients for resection of hepatic colorectal metastases: expert consensus statement. Ann Surg Oncol 2006;13:1261-1268. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16947009>.

401. Gonzalez M, Poncet A, Combescure C, et al. Risk factors for survival after lung metastasectomy in colorectal cancer patients: a systematic review and meta-analysis. Ann Surg Oncol 2013;20:572-579. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23104709>.

402. Gonzalez M, Gervaz P. Risk factors for survival after lung metastasectomy in colorectal cancer patients: systematic review and meta-analysis. Future Oncol 2015;11:31-33. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25662325>.

403. Brouquet A, Vauthey JN, Contreras CM, et al. Improved survival after resection of liver and lung colorectal metastases compared with liver-only metastases: a study of 112 patients with limited lung metastatic disease. J Am Coll Surg 2011;213:62-69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21700179>.

404. Hadden WJ, de Reuver PR, Brown K, et al. Resection of colorectal liver metastases and extra-hepatic disease: a systematic review and proportional meta-analysis of survival outcomes. HPB (Oxford) 2016;18:209-220. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27017160>.

405. Headrick JR, Miller DL, Nagorney DM, et al. Surgical treatment of hepatic and pulmonary metastases from colon cancer. Ann Thorac Surg 2001;71:975-979; discussion 979-980. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11269484>.

406. Marin C, Robles R, Lopez Conesa A, et al. Outcome of strict patient selection for surgical treatment of hepatic and pulmonary metastases from colorectal cancer. Dis Colon Rectum 2013;56:43-50. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23222279>.

407. Pulitano C, Bodingbauer M, Aldrighetti L, et al. Liver resection for colorectal metastases in presence of extrahepatic disease: results from an international multi-institutional analysis. Ann Surg Oncol 2011;18:1380-1388. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21136180>.

408. Carpizo DR, Are C, Jarnagin W, et al. Liver resection for metastatic colorectal cancer in patients with concurrent extrahepatic disease: results in 127 patients treated at a single center. Ann Surg Oncol 2009;16:2138-2146. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19495884>.

409. Carpizo DR, D'Angelica M. Liver resection for metastatic colorectal cancer in the presence of extrahepatic disease. Ann Surg Oncol 2009;16:2411-2421. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19554376>.

410. Chua TC, Saxena A, Liauw W, et al. Hepatectomy and resection of concomitant extrahepatic disease for colorectal liver metastases--a systematic review. Eur J Cancer 2012;48:1757-1765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22153217>.



411. Andreou A, Brouquet A, Abdalla EK, et al. Repeat hepatectomy for recurrent colorectal liver metastases is associated with a high survival rate. *HPB (Oxford)* 2011;13:774-782. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21999590>.

412. de Jong MC, Mayo SC, Pulitano C, et al. Repeat curative intent liver surgery is safe and effective for recurrent colorectal liver metastasis: results from an international multi-institutional analysis. *J Gastrointest Surg* 2009;13:2141-2151. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19795176>.

413. Homayounfar K, Bleckmann A, Conradi LC, et al. Metastatic recurrence after complete resection of colorectal liver metastases: impact of surgery and chemotherapy on survival. *Int J Colorectal Dis* 2013;28:1009-1017. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23371333>.

414. Neeff HP, Drogwitz O, Holzner P, et al. Outcome after repeat resection of liver metastases from colorectal cancer. *Int J Colorectal Dis* 2013;28:1135-1141. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23468250>.

415. Luo LX, Yu ZY, Huang JW, Wu H. Selecting patients for a second hepatectomy for colorectal metastases: An systemic review and meta-analysis. *Eur J Surg Oncol* 2014;40:1036-1048. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24915859>.

416. Adam R, Bismuth H, Castaing D, et al. Repeat hepatectomy for colorectal liver metastases. *Ann Surg* 1997;225:51-60; discussion 60-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8998120>.

417. Salah S, Watanabe K, Park JS, et al. Repeated resection of colorectal cancer pulmonary oligometastases: pooled analysis and prognostic assessment. *Ann Surg Oncol* 2013;20:1955-1961. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23334254>.

418. Poultides GA, Servais EL, Saltz LB, et al. Outcome of primary tumor in patients with synchronous stage IV colorectal cancer receiving

combination chemotherapy without surgery as initial treatment. *J Clin Oncol* 2009;27:3379-3384. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19487380>.

419. Gillams A, Goldberg N, Ahmed M, et al. Thermal ablation of colorectal liver metastases: a position paper by an international panel of ablation experts, the interventional oncology sans frontieres meeting 2013. *Eur Radiol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25994193>.

420. Shady W, Petre EN, Gonen M, et al. Percutaneous radiofrequency ablation of colorectal cancer liver metastases: factors affecting outcomes—a 10-year experience at a single center. *Radiology* 2015;142489. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26267832>.

421. Solbiati L, Ahmed M, Cova L, et al. Small liver colorectal metastases treated with percutaneous radiofrequency ablation: local response rate and long-term survival with up to 10-year follow-up. *Radiology* 2012;265:958-968. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23091175>.

422. Lee MT, Kim JJ, Dinniwell R, et al. Phase I study of individualized stereotactic body radiotherapy of liver metastases. *J Clin Oncol* 2009;27:1585-1591. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19255313>.

423. Rusthoven KE, Kavanagh BD, Cardenas H, et al. Multi-institutional phase I/II trial of stereotactic body radiation therapy for liver metastases. *J Clin Oncol* 2009;27:1572-1578. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19255321>.

424. Alsina J, Choti MA. Liver-directed therapies in colorectal cancer. *Semin Oncol* 2011;38:561-567. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21810515>.



425. Johnston FM, Mavros MN, Herman JM, Pawlik TM. Local therapies for hepatic metastases. *J Natl Compr Canc Netw* 2013;11:153-160. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23411382>.

426. Park J, Chen YJ, Lu WP, Fong Y. The evolution of liver-directed treatments for hepatic colorectal metastases. *Oncology (Williston Park)* 2014;28:991-1003. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25403632>.

427. Zacharias AJ, Jayakrishnan TT, Rajeev R, et al. Comparative effectiveness of hepatic artery based therapies for unresectable colorectal liver metastases: a meta-analysis. *PLoS One* 2015;10:e0139940. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26448327>.

428. Kemeny N, Huang Y, Cohen AM, et al. Hepatic arterial infusion of chemotherapy after resection of hepatic metastases from colorectal cancer. *N Engl J Med* 1999;341:2039-2048. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10615075>.

429. Kemeny NE, Gonen M. Hepatic arterial infusion after liver resection. *N Engl J Med* 2005;352:734-735. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15716576>.

430. Chan DL, Alzahrani NA, Morris DL, Chua TC. Systematic review and meta-analysis of hepatic arterial infusion chemotherapy as bridging therapy for colorectal liver metastases. *Surg Oncol* 2015;24:162-171. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26133575>.

431. Levi FA, Boige V, Hebbar M, et al. Conversion to resection of liver metastases from colorectal cancer with hepatic artery infusion of combined chemotherapy and systemic cetuximab in multicenter trial OPTILIV. *Ann Oncol* 2016;27:267-274. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26578731>.

432. Fiorentini G, Aliberti C, Tilli M, et al. Intra-arterial infusion of irinotecan-loaded drug-eluting beads (DEBIRI) versus intravenous therapy (FOLFIRI) for hepatic metastases from colorectal cancer: final

results of a phase III study. *Anticancer Res* 2012;32:1387-1395. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22493375>.

433. Richardson AJ, Laurence JM, Lam VW. Transarterial chemoembolization with irinotecan beads in the treatment of colorectal liver metastases: systematic review. *J Vasc Interv Radiol* 2013;24:1209-1217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23885916>.

434. Martin RC, 2nd, Scoggins CR, Schreeder M, et al. Randomized controlled trial of irinotecan drug-eluting beads with simultaneous FOLFOX and bevacizumab for patients with unresectable colorectal liver-limited metastasis. *Cancer* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26149602>.

435. Lammer J, Malagari K, Vogl T, et al. Prospective randomized study of doxorubicin-eluting-bead embolization in the treatment of hepatocellular carcinoma: results of the PRECISION V study. *Cardiovasc Intervent Radiol* 2010;33:41-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19908093>.

436. Martin RC, Howard J, Tomalty D, et al. Toxicity of irinotecan-eluting beads in the treatment of hepatic malignancies: results of a multi-institutional registry. *Cardiovasc Intervent Radiol* 2010;33:960-966. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20661569>.

437. Pawlik TM, Reyes DK, Cosgrove D, et al. Phase II trial of sorafenib combined with concurrent transarterial chemoembolization with drug-eluting beads for hepatocellular carcinoma. *J Clin Oncol* 2011;29:3960-3967. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21911714>.

438. Reyes DK, Vossen JA, Kamel IR, et al. Single-center phase II trial of transarterial chemoembolization with drug-eluting beads for patients with unresectable hepatocellular carcinoma: initial experience in the United States. *Cancer J* 2009;15:526-532. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20010173>.

439. van Malenstein H, Maleux G, Vandecaveye V, et al. A randomized phase II study of drug-eluting beads versus transarterial



chemoembolization for unresectable hepatocellular carcinoma.

Onkologie 2011;34:368-376. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21734423>.

440. Vogl TJ, Lammer J, Lencioni R, et al. Liver, gastrointestinal, and cardiac toxicity in intermediate hepatocellular carcinoma treated with PRECISION TACE with drug-eluting beads: results from the PRECISION V randomized trial. AJR Am J Roentgenol 2011;197:W562-570. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21940527>.

441. Riemsma RP, Bala MM, Wolff R, Kleijnen J. Transarterial (chemo)embolisation versus no intervention or placebo intervention for liver metastases. Cochrane Database Syst Rev 2013;4:CD009498. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23633373>.

442. Cosimelli M, Golfieri R, Cagol PP, et al. Multi-centre phase II clinical trial of yttrium-90 resin microspheres alone in unresectable, chemotherapy refractory colorectal liver metastases. Br J Cancer 2010;103:324-331. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20628388>.

443. Gray B, Van Hazel G, Hope M, et al. Randomised trial of SIR-Spheres plus chemotherapy vs. chemotherapy alone for treating patients with liver metastases from primary large bowel cancer. Ann Oncol 2001;12:1711-1720. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11843249>.

444. Hickey R, Lewandowski RJ, Prudhomme T, et al. 90Y radioembolization of colorectal hepatic metastases using glass microspheres: safety and survival outcomes from a 531-patient multicenter study. J Nucl Med 2016;57:665-671. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26635340>.

445. Hong K, McBride JD, Georgiades CS, et al. Salvage therapy for liver-dominant colorectal metastatic adenocarcinoma: comparison between transcatheter arterial chemoembolization versus yttrium-90 radioembolization. J Vasc Interv Radiol 2009;20:360-367. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19167245>.

446. Lewandowski RJ, Memon K, Mulcahy MF, et al. Twelve-year experience of radioembolization for colorectal hepatic metastases in 214 patients: survival by era and chemotherapy. Eur J Nucl Med Mol Imaging 2014;41:1861-1869. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24906565>.

447. Lim L, Gibbs P, Yip D, et al. A prospective evaluation of treatment with Selective Internal Radiation Therapy (SIR-spheres) in patients with unresectable liver metastases from colorectal cancer previously treated with 5-FU based chemotherapy. BMC Cancer 2005;5:132. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16225697>.

448. Mulcahy MF, Lewandowski RJ, Ibrahim SM, et al. Radioembolization of colorectal hepatic metastases using yttrium-90 microspheres. Cancer 2009;115:1849-1858. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19267416>.

449. Seidensticker R, Denecke T, Kraus P, et al. Matched-pair comparison of radioembolization plus best supportive care versus best supportive care alone for chemotherapy refractory liver-dominant colorectal metastases. Cardiovasc Intervent Radiol 2012;35:1066-1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21800231>.

450. Sofocleous CT, Garcia AR, Pandit-Taskar N, et al. Phase I trial of selective internal radiation therapy for chemorefractory colorectal cancer liver metastases progressing after hepatic arterial pump and systemic chemotherapy. Clin Colorectal Cancer 2014;13:27-36. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24370352>.

451. Sofocleous CT, Violari EG, Sotirchos VS, et al. Radioembolization as a salvage therapy for heavily pretreated patients with colorectal cancer liver metastases: factors that affect outcomes. Clin Colorectal Cancer 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26277696>.

452. van Hazel GA, Pavlakis N, Goldstein D, et al. Treatment of fluorouracil-refractory patients with liver metastases from colorectal cancer by using yttrium-90 resin microspheres plus concomitant



systemic irinotecan chemotherapy. *J Clin Oncol* 2009;27:4089-4095. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19652069>.

453. Katz AW, Carey-Sampson M, Muhs AG, et al. Hypofractionated stereotactic body radiation therapy (SBRT) for limited hepatic metastases. *Int J Radiat Oncol Biol Phys* 2007;67:793-798. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17197128>.

454. Agolli L, Bracci S, Nicosia L, et al. Lung metastases treated with stereotactic ablative radiation therapy in oligometastatic colorectal cancer patients: outcomes and prognostic factors after long-term follow-up. *Clin Colorectal Cancer* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27522627>.

455. Chang DT, Swaminath A, Kozak M, et al. Stereotactic body radiotherapy for colorectal liver metastases: a pooled analysis. *Cancer* 2011;117:4060-4069. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21432842>.

456. ACR –ASTRO Practice Parameter for Intensity Modulated Radiation Therapy (IMRT). The American College of Radiology; 2014. Available at: <http://www.acr.org/~media/ACR/Documents/PGTS/guidelines/IMRT.pdf>. Accessed March 1, 2016.

457. Hong TS, Ritter MA, Tome WA, Harari PM. Intensity-modulated radiation therapy: emerging cancer treatment technology. *Br J Cancer* 2005;92:1819-1824. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15856036>.

458. Meyer J, Czito B, Yin F-F, Willett C. Advanced radiation therapy technologies in the treatment of rectal and anal cancer: intensity-modulated photon therapy and proton therapy. *Clin Colorectal Cancer* 2007;6:348-356. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17311699>.

459. Topkan E, Onal HC, Yavuz MN. Managing liver metastases with conformal radiation therapy. *J Support Oncol* 2008;6:9-13, 15. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18257395>.

460. Hendlitz A, Van den Eynde M, Peeters M, et al. Phase III trial comparing protracted intravenous fluorouracil infusion alone or with yttrium-90 resin microspheres radioembolization for liver-limited metastatic colorectal cancer refractory to standard chemotherapy. *J Clin Oncol* 2010;28:3687-3694. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20567019>.

461. Benson AB, 3rd, Geschwind JF, Mulcahy MF, et al. Radioembolisation for liver metastases: results from a prospective 151 patient multi-institutional phase II study. *Eur J Cancer* 2013;49:3122-3130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23777743>.

462. Kennedy AS, Ball D, Cohen SJ, et al. Multicenter evaluation of the safety and efficacy of radioembolization in patients with unresectable colorectal liver metastases selected as candidates for (90)Y resin microspheres. *J Gastrointest Oncol* 2015;6:134-142. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25830033>.

463. Saxena A, Meteling B, Kapoor J, et al. Is yttrium-90 radioembolization a viable treatment option for unresectable, chemorefractory colorectal cancer liver metastases? A large single-center experience of 302 patients. *Ann Surg Oncol* 2015;22:794-802. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25323474>.

464. van Hazel GA, Heinemann V, Sharma NK, et al. SIRFLOX: randomized phase III trial comparing first-line mFOLFOX6 (plus or minus bevacizumab) versus mFOLFOX6 (plus or minus bevacizumab) plus selective internal radiation therapy in patients with metastatic colorectal cancer. *J Clin Oncol* 2016;34:1723-1731. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26903575>.

465. Rosenbaum CE, Verkooijen HM, Lam MG, et al. Radioembolization for treatment of salvage patients with colorectal cancer liver metastases: a systematic review. *J Nucl Med*



2013;54:1890-1895. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24071510>.

466. Saxena A, Bester L, Shan L, et al. A systematic review on the safety and efficacy of yttrium-90 radioembolization for unresectable, chemorefractory colorectal cancer liver metastases. *J Cancer Res Clin Oncol* 2014;140:537-547. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24318568>.

467. Townsend A, Price T, Karapetis C. Selective internal radiation therapy for liver metastases from colorectal cancer. *Cochrane Database Syst Rev* 2009;CD007045. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19821394>.

468. Abdalla EK. Commentary: Radiofrequency ablation for colorectal liver metastases: do not blame the biology when it is the technology. *Am J Surg* 2009;197:737-739. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18789420>.

469. Wang X, Sofocleous CT, Erinjeri JP, et al. Margin size is an independent predictor of local tumor progression after ablation of colon cancer liver metastases. *Cardiovasc Intervent Radiol* 2013;36:166-175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22535243>.

470. Elias D, De Baere T, Smayra T, et al. Percutaneous radiofrequency thermoablation as an alternative to surgery for treatment of liver tumour recurrence after hepatectomy. *Br J Surg* 2002;89:752-756. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12027986>.

471. Sofocleous CT, Petre EN, Gonen M, et al. CT-guided radiofrequency ablation as a salvage treatment of colorectal cancer hepatic metastases developing after hepatectomy. *J Vasc Interv Radiol* 2011;22:755-761. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21514841>.

472. Sucandy I, Cheek S, Golas BJ, et al. Longterm survival outcomes of patients undergoing treatment with radiofrequency ablation for hepatocellular carcinoma and metastatic colorectal cancer liver tumors.

HPB (Oxford) 2016;18:756-763. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/27593593>.

473. Bala MM, Riemsma RP, Wolff R, Kleijnen J. Microwave coagulation for liver metastases. *Cochrane Database Syst Rev* 2013;10:CD010163. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24122576>.

474. Bala MM, Riemsma RP, Wolff R, Kleijnen J. Cryotherapy for liver metastases. *Cochrane Database Syst Rev* 2013;6:CD009058. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23740609>.

475. Cirocchi R, Trastulli S, Boselli C, et al. Radiofrequency ablation in the treatment of liver metastases from colorectal cancer. *Cochrane Database Syst Rev* 2012;6:CD006317. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22696357>.

476. Riemsma RP, Bala MM, Wolff R, Kleijnen J. Percutaneous ethanol injection for liver metastases. *Cochrane Database Syst Rev* 2013;5:CD008717. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23728679>.

477. Riemsma RP, Bala MM, Wolff R, Kleijnen J. Electro-coagulation for liver metastases. *Cochrane Database Syst Rev* 2013;5:CD009497. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23728692>.

478. Weng M, Zhang Y, Zhou D, et al. Radiofrequency ablation versus resection for colorectal cancer liver metastases: a meta-analysis. *PLoS One* 2012;7:e45493. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23029051>.

479. Wong SL, Mangu PB, Choti MA, et al. American Society of Clinical Oncology 2009 clinical evidence review on radiofrequency ablation of hepatic metastases from colorectal cancer. *J Clin Oncol* 2010;28:493-508. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19841322>.

480. Gillams A, Khan Z, Osborn P, Lees W. Survival after radiofrequency ablation in 122 patients with inoperable colorectal lung



metastases. *Cardiovasc Intervent Radiol* 2013;36:724-730. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23070108>.

481. Gleisner AL, Choti MA, Assumpcao L, et al. Colorectal liver metastases: recurrence and survival following hepatic resection, radiofrequency ablation, and combined resection-radiofrequency ablation. *Arch Surg* 2008;143:1204-1212. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19075173>.

482. Reuter NP, Woodall CE, Scoggins CR, et al. Radiofrequency ablation vs. resection for hepatic colorectal metastasis: therapeutically equivalent? *J Gastrointest Surg* 2009;13:486-491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18972167>.

483. de Jong MC, Pulitano C, Ribero D, et al. Rates and patterns of recurrence following curative intent surgery for colorectal liver metastasis: an international multi-institutional analysis of 1669 patients. *Ann Surg* 2009;250:440-448. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19730175>.

484. Bai H, Huangz X, Jing L, et al. The effect of radiofrequency ablation vs. liver resection on survival outcome of colorectal liver metastases (CRLM): a meta-analysis. *Hepatogastroenterology* 2015;62:373-377. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25916066>.

485. Ruers T, Punt C, Van Coevorden F, et al. Radiofrequency ablation combined with systemic treatment versus systemic treatment alone in patients with non-resectable colorectal liver metastases: a randomized EORTC Intergroup phase II study (EORTC 40004). *Ann Oncol* 2012;23:2619-2626. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22431703>.

486. Klaver YL, Leenders BJ, Creemers GJ, et al. Addition of biological therapies to palliative chemotherapy prolongs survival in patients with peritoneal carcinomatosis of colorectal origin. *Am J Clin Oncol* 2013;36:157-161. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22314003>.

487. Takahashi H, Okabayashi K, Tsuruta M, et al. Self-expanding metallic stents versus surgical intervention as palliative therapy for obstructive colorectal cancer: a meta-analysis. *World J Surg* 2015;39:2037-2044. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25894403>.

488. van Hooft JE, van Halsema EE, Vanbiervliet G, et al. Self-expandable metal stents for obstructing colonic and extracolonic cancer: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline. *Gastrointest Endosc* 2014;80:747-761 e741-775. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25436393>.

489. Cennamo V, Fuccio L, Mutri V, et al. Does stent placement for advanced colon cancer increase the risk of perforation during bevacizumab-based therapy? *Clin Gastroenterol Hepatol* 2009;7:1174-1176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19631290>.

490. Small AJ, Coelho-Prabhu N, Baron TH. Endoscopic placement of self-expandable metal stents for malignant colonic obstruction: long-term outcomes and complication factors. *Gastrointest Endosc* 2010;71:560-572. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20189515>.

491. Baratti D, Kusamura S, Pietrantonio F, et al. Progress in treatments for colorectal cancer peritoneal metastases during the years 2010-2015. A systematic review. *Crit Rev Oncol Hematol* 2016;100:209-222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26867984>.

492. Chua TC, Pelz JO, Kerscher A, et al. Critical analysis of 33 patients with peritoneal carcinomatosis secondary to colorectal and appendiceal signet ring cell carcinoma. *Ann Surg Oncol* 2009;16:2765-2770. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19641972>.

493. Elias D, Gilly F, Boutitie F, et al. Peritoneal colorectal carcinomatosis treated with surgery and perioperative intraperitoneal chemotherapy: retrospective analysis of 523 patients from a multicentric



French study. J Clin Oncol 2010;28:63-68. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19917863>.

494. Esquivel J, Sticca R, Sugarbaker P, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy in the management of peritoneal surface malignancies of colonic origin: a consensus statement. Society of Surgical Oncology. Ann Surg Oncol 2007;14:128-133. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17072675>.

495. Goere D, Malka D, Tzani D, et al. Is there a possibility of a cure in patients with colorectal peritoneal carcinomatosis amenable to complete cytoreductive surgery and intraperitoneal chemotherapy? Ann Surg 2013;257:1065-1071. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23299520>.

496. Haslinger M, Francescutti V, Attwood K, et al. A contemporary analysis of morbidity and outcomes in cytoreduction/hyperthermic intraperitoneal chemoperfusion. Cancer Med 2013;2:334-342. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23930210>.

497. Glehen O, Kwiatkowski F, Sugarbaker PH, et al. Cytoreductive surgery combined with perioperative intraperitoneal chemotherapy for the management of peritoneal carcinomatosis from colorectal cancer: a multi-institutional study. J Clin Oncol 2004;22:3284-3292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15310771>.

498. Tabrizian P, Shrager B, Jibara G, et al. Cytoreductive surgery and hyperthermic intraperitoneal chemotherapy for peritoneal carcinomatosis: outcomes from a single tertiary institution. J Gastrointest Surg 2014;18:1024-1031. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24577736>.

499. Yan TD, Black D, Savady R, Sugarbaker PH. Systematic review on the efficacy of cytoreductive surgery combined with perioperative intraperitoneal chemotherapy for peritoneal carcinomatosis from colorectal carcinoma. J Clin Oncol 2006;24:4011-4019. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16921055>.

500. Verwaal VJ, van Ruth S, de Bree E, et al. Randomized trial of cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy and palliative surgery in patients with peritoneal carcinomatosis of colorectal cancer. J Clin Oncol 2003;21:3737-3743. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14551293>.

501. Verwaal VJ, Bruin S, Boot H, et al. 8-year follow-up of randomized trial: cytoreduction and hyperthermic intraperitoneal chemotherapy versus systemic chemotherapy in patients with peritoneal carcinomatosis of colorectal cancer. Ann Surg Oncol 2008;15:2426-2432. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18521686>.

502. Sugarbaker PH, Ryan DP. Cytoreductive surgery plus hyperthermic perioperative chemotherapy to treat peritoneal metastases from colorectal cancer: standard of care or an experimental approach? Lancet Oncol 2012;13:e362-369. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22846841>.

503. El Halabi H, Gushchin V, Francis J, et al. The role of cytoreductive surgery and heated intraperitoneal chemotherapy (CRS/HIPEC) in patients with high-grade appendiceal carcinoma and extensive peritoneal carcinomatosis. Ann Surg Oncol 2012;19:110-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21701929>.

504. Glehen O, Gilly FN, Boutitie F, et al. Toward curative treatment of peritoneal carcinomatosis from nonovarian origin by cytoreductive surgery combined with perioperative intraperitoneal chemotherapy: a multi-institutional study of 1,290 patients. Cancer 2010;116:5608-5618. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20737573>.

505. Shaib WL, Martin LK, Choi M, et al. Hyperthermic intraperitoneal chemotherapy following cytoreductive surgery improves outcome in patients with primary appendiceal mucinous adenocarcinoma: a pooled analysis from three tertiary care centers. Oncologist 2015;20:907-914. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26070916>.

506. Chua TC, Moran BJ, Sugarbaker PH, et al. Early- and long-term outcome data of patients with pseudomyxoma peritonei from



appendiceal origin treated by a strategy of cytoreductive surgery and hyperthermic intraperitoneal chemotherapy. *J Clin Oncol* 2012;30:2449-2456. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22614976>.

507. Faris JE, Ryan DP. Controversy and consensus on the management of patients with pseudomyxoma peritonei. *Curr Treat Options Oncol* 2013;14:365-373. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23934509>.

508. Klaver YL, Hendriks T, Lomme RM, et al. Hyperthermia and intraperitoneal chemotherapy for the treatment of peritoneal carcinomatosis: an experimental study. *Ann Surg* 2011;254:125-130. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21502859>.

509. Cashin PH, Mahteme H, Spang N, et al. Cytoreductive surgery and intraperitoneal chemotherapy versus systemic chemotherapy for colorectal peritoneal metastases: A randomised trial. *Eur J Cancer* 2016;53:155-162. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26751236>.

510. van Oudheusden TR, Nienhuijs SW, Luyer MD, et al. Incidence and treatment of recurrent disease after cytoreductive surgery and intraperitoneal chemotherapy for peritoneally metastasized colorectal cancer: A systematic review. *Eur J Surg Oncol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26175345>.

511. Esquivel J. Colorectal cancer with peritoneal metastases: a plea for cooperation between medical and surgical oncologists. *Oncology (Williston Park)* 2015;29:521-522. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26178340>.

512. Loggie BW, Thomas P. Gastrointestinal cancers with peritoneal carcinomatosis: surgery and hyperthermic intraperitoneal chemotherapy. *Oncology (Williston Park)* 2015;29:515-521. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26178339>.

513. McRee AJ, O'Neil BH. The role of HIPEC in gastrointestinal malignancies: controversies and conclusions. *Oncology (Williston Park)*

2015;29:523-524, C523. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26178341>.

514. O'Dwyer S, Verwaal VJ, Sugarbaker PH. Evolution of treatments for peritoneal metastases from colorectal cancer. *J Clin Oncol* 2015;33:2122-2123. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25897165>.

515. Altendorf-Hofmann A, Scheele J. A critical review of the major indicators of prognosis after resection of hepatic metastases from colorectal carcinoma. *Surg Oncol Clin N Am* 2003;12:165-192. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12735137>.

516. Pawlik TM, Schulick RD, Choti MA. Expanding criteria for resectability of colorectal liver metastases. *Oncologist* 2008;13:51-64. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18245012>.

517. Pozzo C, Basso M, Cassano A, et al. Neoadjuvant treatment of unresectable liver disease with irinotecan and 5-fluorouracil plus folinic acid in colorectal cancer patients. *Ann Oncol* 2004;15:933-939. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15151951>.

518. Vauthey J-N, Zorzi D, Pawlik TM. Making unresectable hepatic colorectal metastases resectable--does it work? *Semin Oncol* 2005;32:118-122. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16399448>.

519. Covey AM, Brown KT, Jarnagin WR, et al. Combined portal vein embolization and neoadjuvant chemotherapy as a treatment strategy for resectable hepatic colorectal metastases. *Ann Surg* 2008;247:451-455. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18376189>.

520. Folprecht G, Grothey A, Alberts S, et al. Neoadjuvant treatment of unresectable colorectal liver metastases: correlation between tumour response and resection rates. *Ann Oncol* 2005;16:1311-1319. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15870084>.



521. Bilchik AJ, Poston G, Curley SA, et al. Neoadjuvant chemotherapy for metastatic colon cancer: a cautionary note. *J Clin Oncol* 2005;23:9073-9078. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16361615>.

522. Choti MA. Chemotherapy-associated hepatotoxicity: do we need to be concerned? *Ann Surg Oncol* 2009;16:2391-2394. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19554374>.

523. Kishi Y, Zorzi D, Contreras CM, et al. Extended preoperative chemotherapy does not improve pathologic response and increases postoperative liver insufficiency after hepatic resection for colorectal liver metastases. *Ann Surg Oncol* 2010;17:2870-2876. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20567921>.

524. Rubbia-Brandt L, Audard V, Sartoretti P, et al. Severe hepatic sinusoidal obstruction associated with oxaliplatin-based chemotherapy in patients with metastatic colorectal cancer. *Ann Oncol* 2004;15:460-466. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14998849>.

525. Vauthey J-N, Pawlik TM, Ribero D, et al. Chemotherapy regimen predicts steatohepatitis and an increase in 90-day mortality after surgery for hepatic colorectal metastases. *J Clin Oncol* 2006;24:2065-2072. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16648507>.

526. Delaunoy T, Alberts SR, Sargent DJ, et al. Chemotherapy permits resection of metastatic colorectal cancer: experience from Intergroup N9741. *Ann Oncol* 2005;16:425-429. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15677624>.

527. Falcone A, Ricci S, Brunetti I, et al. Phase III trial of infusional fluorouracil, leucovorin, oxaliplatin, and irinotecan (FOLFOXIRI) compared with infusional fluorouracil, leucovorin, and irinotecan (FOLFIRI) as first-line treatment for metastatic colorectal cancer: the Gruppo Oncologico Nord Ovest. *J Clin Oncol* 2007;25:1670-1676. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17470860>.

528. Souglakos J, Androulakis N, Syrigos K, et al. FOLFOXIRI (folinic acid, 5-fluorouracil, oxaliplatin and irinotecan) vs FOLFIRI (folinic acid, 5-fluorouracil and irinotecan) as first-line treatment in metastatic colorectal cancer (MCC): a multicentre randomised phase III trial from the Hellenic Oncology Research Group (HORG). *Br J Cancer* 2006;94:798-805. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16508637>.

529. Masi G, Vasile E, Loupakis F, et al. Randomized trial of two induction chemotherapy regimens in metastatic colorectal cancer: an updated analysis. *J Natl Cancer Inst* 2011;103:21-30. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21123833>.

530. Folprecht G, Gruenberger T, Bechstein WO, et al. Tumour response and secondary resectability of colorectal liver metastases following neoadjuvant chemotherapy with cetuximab: the CELIM randomised phase 2 trial. *Lancet Oncol* 2010;11:38-47. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19942479>.

531. Ye LC, Liu TS, Ren L, et al. Randomized controlled trial of cetuximab plus chemotherapy for patients with KRAS wild-type unresectable colorectal liver-limited metastases. *J Clin Oncol* 2013;31:1931-1938. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23569301>.

532. Petrelli F, Barni S. Resectability and outcome with anti-EGFR agents in patients with KRAS wild-type colorectal liver-limited metastases: a meta-analysis. *Int J Colorectal Dis* 2012;27:997-1004. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22358385>.

533. Fuchs CS, Marshall J, Mitchell E, et al. Randomized, controlled trial of irinotecan plus infusional, bolus, or oral fluoropyrimidines in first-line treatment of metastatic colorectal cancer: results from the BICC-C Study. *J Clin Oncol* 2007;25:4779-4786. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17947725>.

534. Saltz LB, Clarke S, Diaz-Rubio E, et al. Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in



metastatic colorectal cancer: a randomized phase III study. *J Clin Oncol* 2008;26:2013-2019. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18421054>.

535. Adam R, Avisar E, Ariche A, et al. Five-year survival following hepatic resection after neoadjuvant therapy for nonresectable colorectal. *Ann Surg Oncol* 2001;8:347-353. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11352309>.

536. Pawlik TM, Olin K, Gleisner AL, et al. Preoperative chemotherapy for colorectal liver metastases: impact on hepatic histology and postoperative outcome. *J Gastrointest Surg* 2007;11:860-868. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17492335>.

537. Rivoire M, De Cian F, Meeus P, et al. Combination of neoadjuvant chemotherapy with cryotherapy and surgical resection for the treatment of unresectable liver metastases from colorectal carcinoma. *Cancer* 2002;95:2283-2292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12436433>.

538. Ciliberto D, Prati U, Roveda L, et al. Role of systemic chemotherapy in the management of resected or resectable colorectal liver metastases: a systematic review and meta-analysis of randomized controlled trials. *Oncol Rep* 2012;27:1849-1856. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22446591>.

539. Wang ZM, Chen YY, Chen FF, et al. Peri-operative chemotherapy for patients with resectable colorectal hepatic metastasis: A meta-analysis. *Eur J Surg Oncol* 2015;41:1197-1203. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26094113>.

540. Araujo R, Gonen M, Allen P, et al. Comparison between perioperative and postoperative chemotherapy after potentially curative hepatic resection for metastatic colorectal cancer. *Ann Surg Oncol* 2013;20:4312-4321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23897009>.

541. Khoo E, O'Neill S, Brown E, et al. Systematic review of systemic adjuvant, neoadjuvant and perioperative chemotherapy for resectable colorectal-liver metastases. *HPB (Oxford)* 2016;18:485-493. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27317952>.

542. Nordlinger B, Sorbye H, Glimelius B, et al. Perioperative chemotherapy with FOLFOX4 and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC Intergroup trial 40983): a randomised controlled trial. *Lancet* 2008;371:1007-1016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18358928>.

543. Nordlinger B, Sorbye H, Glimelius B, et al. Perioperative FOLFOX4 chemotherapy and surgery versus surgery alone for resectable liver metastases from colorectal cancer (EORTC 40983): long-term results of a randomised, controlled, phase 3 trial. *Lancet Oncol* 2013;14:1208-1215. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24120480>.

544. Bilchik AJ, Poston G, Adam R, Choti MA. Prognostic variables for resection of colorectal cancer hepatic metastases: an evolving paradigm. *J Clin Oncol* 2008;26:5320-5321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18936470>.

545. Leonard GD, Brenner B, Kemeny NE. Neoadjuvant chemotherapy before liver resection for patients with unresectable liver metastases from colorectal carcinoma. *J Clin Oncol* 2005;23:2038-2048. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15774795>.

546. van Vledder MG, de Jong MC, Pawlik TM, et al. Disappearing colorectal liver metastases after chemotherapy: should we be concerned? *J Gastrointest Surg* 2010;14:1691-1700. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20839072>.

547. Benoist S, Brouquet A, Penna C, et al. Complete response of colorectal liver metastases after chemotherapy: does it mean cure? *J Clin Oncol* 2006;24:3939-3945. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16921046>.



548. Scappaticci FA, Fehrenbacher L, Cartwright T, et al. Surgical wound healing complications in metastatic colorectal cancer patients treated with bevacizumab. *J Surg Oncol* 2005;91:173-180. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16118771>.

549. Ranpura V, Hapani S, Wu S. Treatment-related mortality with bevacizumab in cancer patients: a meta-analysis. *JAMA* 2011;305:487-494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21285426>.

550. Hurwitz HI, Saltz LB, Van Cutsem E, et al. Venous thromboembolic events with chemotherapy plus bevacizumab: a pooled analysis of patients in randomized phase II and III studies. *J Clin Oncol* 2011;29:1757-1764. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21422411>.

551. Package Insert. AVASTIN® (bevacizumab). Genentech, Inc.; 2015. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/125085s312bl.pdf. Accessed March 1, 2016.

552. Primrose J, Falk S, Finch-Jones M, et al. Systemic chemotherapy with or without cetuximab in patients with resectable colorectal liver metastasis: the New EPOC randomised controlled trial. *Lancet Oncol* 2014;15:601-611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24717919>.

553. Brule SY, Jonker DJ, Karapetis CS, et al. Location of colon cancer (right-sided versus left-sided) as a prognostic factor and a predictor of benefit from cetuximab in NCIC CO.17. *Eur J Cancer* 2015;51:1405-1414. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25979833>.

554. Moretto R, Cremolini C, Rossini D, et al. Location of Primary Tumor and Benefit From Anti-Epidermal Growth Factor Receptor Monoclonal Antibodies in Patients With RAS and BRAF Wild-Type Metastatic Colorectal Cancer. *Oncologist* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27382031>.

555. Loupakis F, Yang D, Yau L, et al. Primary tumor location as a prognostic factor in metastatic colorectal cancer. *J Natl Cancer Inst* 2015;107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25713148>.

556. Lee MS, Advani SM, Morris J, et al. Association of primary (1{degrees}) site and molecular features with progression-free survival (PFS) and overall survival (OS) of metastatic colorectal cancer (mCRC) after anti-epidermal growth factor receptor ({alpha}EGFR) therapy [abstract]. *ASCO Meeting Abstracts* 2016;34:3506. Available at: <http://meetinglibrary.asco.org/content/171167-176>.

557. Chen KH, Shao YY, Chen HM, et al. Primary tumor site is a useful predictor of cetuximab efficacy in the third-line or salvage treatment of KRAS wild-type (exon 2 non-mutant) metastatic colorectal cancer: a nationwide cohort study. *BMC Cancer* 2016;16:327. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27221731>.

558. Warschkow R, Sulz MC, Marti L, et al. Better survival in right-sided versus left-sided stage I - III colon cancer patients. *BMC Cancer* 2016;16:554. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27464835>.

559. Schrag D, Weng S, Brooks G, et al. The relationship between primary tumor sidedness and prognosis in colorectal cancer [abstract]. *ASCO Meeting Abstracts* 2016;34:3505. Available at: <http://meetinglibrary.asco.org/content/167366-176>.

560. Venook AP, Niedzwiecki D, Innocenti F, et al. Impact of primary (1{o}) tumor location on overall survival (OS) and progression-free survival (PFS) in patients (pts) with metastatic colorectal cancer (mCRC): Analysis of CALGB/SWOG 80405 (Alliance) [abstract]. *ASCO Meeting Abstracts* 2016;34:3504. Available at: <http://meetinglibrary.asco.org/content/161936-176>.

561. Venook AP, Niedzwiecki D, Innocenti F, et al. Impact of primary (1°) tumor location on Overall Survival (OS) and Progression Free Survival (PFS) in patients (pts) with metastatic colorectal cancer



(mCRC): Analysis of All RAS wt patients on CALGB / SWOG 80405 (Alliance) [abstract]. ESMO Congress 2016. Available at:

562. Antonacopoulou AG, Tsamandas AC, Petsas T, et al. EGFR, HER-2 and COX-2 levels in colorectal cancer. *Histopathology* 2008;53:698-706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19102009>.

563. McKay JA, Murray LJ, Curran S, et al. Evaluation of the epidermal growth factor receptor (EGFR) in colorectal tumours and lymph node metastases. *Eur J Cancer* 2002;38:2258-2264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12441262>.

564. Spano JP, Lagorce C, Atlan D, et al. Impact of EGFR expression on colorectal cancer patient prognosis and survival. *Ann Oncol* 2005;16:102-108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15598946>.

565. Yen LC, Uen YH, Wu DC, et al. Activating KRAS mutations and overexpression of epidermal growth factor receptor as independent predictors in metastatic colorectal cancer patients treated with cetuximab. *Ann Surg* 2010;251:254-260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20010090>.

566. Cunningham D, Humblet Y, Siena S, et al. Cetuximab monotherapy and cetuximab plus irinotecan in irinotecan-refractory metastatic colorectal cancer. *N Engl J Med* 2004;351:337-345. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15269313>.

567. Hecht JR, Mitchell E, Neubauer MA, et al. Lack of correlation between epidermal growth factor receptor status and response to Panitumumab monotherapy in metastatic colorectal cancer. *Clin Cancer Res* 2010;16:2205-2213. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20332321>.

568. Saltz LB, Meropol NJ, Loehrer PJ, et al. Phase II trial of cetuximab in patients with refractory colorectal cancer that expresses the epidermal growth factor receptor. *J Clin Oncol* 2004;22:1201-1208. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14993230>.

569. Van Cutsem E, Peeters M, Siena S, et al. Open-label phase III trial of panitumumab plus best supportive care compared with best supportive care alone in patients with chemotherapy-refractory metastatic colorectal cancer. *J Clin Oncol* 2007;25:1658-1664. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17470858>.

570. Amado RG, Wolf M, Peeters M, et al. Wild-type KRAS is required for panitumumab efficacy in patients with metastatic colorectal cancer. *J Clin Oncol* 2008;26:1626-1634. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18316791>.

571. Baselga J, Rosen N. Determinants of RASistance to anti-epidermal growth factor receptor agents. *J Clin Oncol* 2008;26:1582-1584. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18316790>.

572. Bokemeyer C, Bondarenko I, Makhson A, et al. Fluorouracil, leucovorin, and oxaliplatin with and without cetuximab in the first-line treatment of metastatic colorectal cancer. *J Clin Oncol* 2009;27:663-671. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19114683>.

573. Dahabreh IJ, Terasawa T, Castaldi PJ, Trikalinos TA. Systematic review: Anti-epidermal growth factor receptor treatment effect modification by KRAS mutations in advanced colorectal cancer. *Ann Intern Med* 2011;154:37-49. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21200037>.

574. De Roock W, Piessevaux H, De Schutter J, et al. KRAS wild-type state predicts survival and is associated to early radiological response in metastatic colorectal cancer treated with cetuximab. *Ann Oncol* 2008;19:508-515. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17998284>.

575. Karapetis CS, Khambata-Ford S, Jonker DJ, et al. K-ras mutations and benefit from cetuximab in advanced colorectal cancer. *N Engl J Med* 2008;359:1757-1765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18946061>.

576. Khambata-Ford S, Garrett CR, Meropol NJ, et al. Expression of epiregulin and amphiregulin and K-ras mutation status predict disease control in metastatic colorectal cancer patients treated with cetuximab. *J Clin Oncol* 2007;25:3230-3237. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17664471>.

577. Lievre A, Bachet J-B, Boige V, et al. KRAS mutations as an independent prognostic factor in patients with advanced colorectal cancer treated with cetuximab. *J Clin Oncol* 2008;26:374-379. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18202412>.

578. Van Cutsem E, Tejpar S, Vanbeckevoort D, et al. Inpatient Cetuximab Dose Escalation in Metastatic Colorectal Cancer According to the Grade of Early Skin Reactions: The Randomized EVEREST Study. *J Clin Oncol* 2012;30:2861-2868. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22753904>.

579. Van Cutsem E, Kohne CH, Hitre E, et al. Cetuximab and chemotherapy as initial treatment for metastatic colorectal cancer. *N Engl J Med* 2009;360:1408-1417. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19339720>.

580. Douillard JY, Oliner KS, Siena S, et al. Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer. *N Engl J Med* 2013;369:1023-1034. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24024839>.

581. Sorich MJ, Wiese MD, Rowland A, et al. Extended RAS mutations and anti-EGFR monoclonal antibody survival benefit in metastatic colorectal cancer: a meta-analysis of randomized, controlled trials. *Ann Oncol* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25115304>.

582. Allegra CJ, Rumble RB, Hamilton SR, et al. Extended RAS gene mutation testing in metastatic colorectal carcinoma to predict response to anti-epidermal growth factor receptor monoclonal antibody therapy: American Society of Clinical Oncology provisional clinical opinion

update 2015. *J Clin Oncol* 2016;34:179-185. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26438111>.

583. Artale S, Sartore-Bianchi A, Veronese SM, et al. Mutations of KRAS and BRAF in primary and matched metastatic sites of colorectal cancer. *J Clin Oncol* 2008;26:4217-4219. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18757341>.

584. Etienne-Grimaldi M-C, Formento J-L, Francoual M, et al. K-Ras mutations and treatment outcome in colorectal cancer patients receiving exclusive fluoropyrimidine therapy. *Clin Cancer Res* 2008;14:4830-4835. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18676755>.

585. Wang HL, Lopategui J, Amin MB, Patterson SD. KRAS mutation testing in human cancers: The pathologist's role in the era of personalized medicine. *Adv Anat Pathol* 2010;17:23-32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20032635>.

586. Monzon FA, Ogino S, Hammond MEH, et al. The role of KRAS mutation testing in the management of patients with metastatic colorectal cancer. *Arch Pathol Lab Med* 2009;133:1600-1606. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19792050>.

587. Roth AD, Tejpar S, Delorenzi M, et al. Prognostic role of KRAS and BRAF in stage II and III resected colon cancer: results of the translational study on the PETACC-3, EORTC 40993, SAKK 60-00 trial. *J Clin Oncol* 2010;28:466-474. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20008640>.

588. Package Insert. Vectibix® (Panitumumab). Thousand Oaks, CA: Amgen Inc.; 2015. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/125147s20_0lbl.pdf. Accessed October 17, 2015.

589. Package Insert. Cetuximab (Erbix®). Branchburg, NJ: ImClone Systems Incorporated; 2015. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/125084s26_2lbl.pdf. Accessed October 7, 2015.



590. Yoon HH, Tougeron D, Shi Q, et al. KRAS codon 12 and 13 mutations in relation to disease-free survival in BRAF-wild-type stage III colon cancers from an adjuvant chemotherapy trial (N0147 alliance). *Clin Cancer Res* 2014;20:3033-3043. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24687927>.

591. De Roock W, Jonker DJ, Di Nicolantonio F, et al. Association of KRAS p.G13D mutation with outcome in patients with chemotherapy-refractory metastatic colorectal cancer treated with cetuximab. *JAMA* 2010;304:1812-1820. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20978259>.

592. Tejpar S, Celik I, Schlichting M, et al. Association of KRAS G13D tumor mutations with outcome in patients with metastatic colorectal cancer treated with first-line chemotherapy with or without cetuximab. *J Clin Oncol* 2012;30:3570-3577. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22734028>.

593. Peeters M, Douillard JY, Van Cutsem E, et al. Mutant KRAS codon 12 and 13 alleles in patients with metastatic colorectal cancer: assessment as prognostic and predictive biomarkers of response to panitumumab. *J Clin Oncol* 2013;31:759-765. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23182985>.

594. Schirripa M, Loupakis F, Lonardi S, et al. Phase II study of single-agent cetuximab in KRAS G13D mutant metastatic colorectal cancer. *Ann Oncol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26371285>.

595. Segelov E, Thavaneswaran S, Waring PM, et al. Response to cetuximab with or without irinotecan in patients with refractory metastatic colorectal cancer harboring the KRAS G13D mutation: Australasian Gastro-Intestinal Trials Group ICECREAM study. *J Clin Oncol* 2016;34:2258-2264. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27114605>.

596. Price TJ, Bruhn MA, Lee CK, et al. Correlation of extended RAS and PIK3CA gene mutation status with outcomes from the phase III

AGITG MAX STUDY involving capecitabine alone or in combination with bevacizumab plus or minus mitomycin C in advanced colorectal cancer. *Br J Cancer* 2015;112:963-970. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25742472>.

597. Heinemann V, von Weikersthal LF, Decker T, et al. FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2014;15:1065-1075. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25088940>.

598. Tol J, Nagtegaal ID, Punt CJA. BRAF mutation in metastatic colorectal cancer. *N Engl J Med* 2009;361:98-99. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19571295>.

599. Van Cutsem E, Lang I, Folprecht G, et al. Cetuximab plus FOLFIRI: final data from the CRYSTAL study on the association of KRAS and BRAF biomarker status with treatment outcome [abstract]. *J Clin Oncol* 2010;28 (May 20 suppl):3570. Available at: http://meeting.ascopubs.org/cgi/content/abstract/28/15_suppl/3570.

600. Maughan TS, Adams RA, Smith CG, et al. Addition of cetuximab to oxaliplatin-based first-line combination chemotherapy for treatment of advanced colorectal cancer: results of the randomised phase 3 MRC COIN trial. *Lancet* 2011;377:2103-2114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21641636>.

601. Davies H, Bignell GR, Cox C, et al. Mutations of the BRAF gene in human cancer. *Nature* 2002;417:949-954. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12068308>.

602. Ikenoue T, Hikiba Y, Kanai F, et al. Functional analysis of mutations within the kinase activation segment of B-Raf in human colorectal tumors. *Cancer Res* 2003;63:8132-8137. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14678966>.

603. Wan PT, Garnett MJ, Roe SM, et al. Mechanism of activation of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. *Cell*



2004;116:855-867. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15035987>.

604. Bokemeyer C, Cutsem EV, Rougier P, et al. Addition of cetuximab to chemotherapy as first-line treatment for KRAS wild-type metastatic colorectal cancer: pooled analysis of the CRYSTAL and OPUS randomised clinical trials. *Eur J Cancer* 2012;48:1466-1475. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22446022>.

605. Van Cutsem E, Kohne CH, Lang I, et al. Cetuximab plus irinotecan, fluorouracil, and leucovorin as first-line treatment for metastatic colorectal cancer: updated analysis of overall survival according to tumor KRAS and BRAF mutation status. *J Clin Oncol* 2011;29:2011-2019. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21502544>.

606. Di Nicolantonio F, Martini M, Molinari F, et al. Wild-type BRAF is required for response to panitumumab or cetuximab in metastatic colorectal cancer. *J Clin Oncol* 2008;26:5705-5712. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19001320>.

607. Laurent-Puig P, Cayre A, Manceau G, et al. Analysis of PTEN, BRAF, and EGFR status in determining benefit from cetuximab therapy in wild-type KRAS metastatic colon cancer. *J Clin Oncol* 2009;27:5924-5930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19884556>.

608. Loupakis F, Ruzzo A, Cremolini C, et al. KRAS codon 61, 146 and BRAF mutations predict resistance to cetuximab plus irinotecan in KRAS codon 12 and 13 wild-type metastatic colorectal cancer. *Br J Cancer* 2009;101:715-721. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19603018>.

609. De Roock W, Claes B, Bernasconi D, et al. Effects of KRAS, BRAF, NRAS, and PIK3CA mutations on the efficacy of cetuximab plus chemotherapy in chemotherapy-refractory metastatic colorectal cancer: a retrospective consortium analysis. *Lancet Oncol* 2010;11:753-762. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20619739>.

610. Seymour MT, Brown SR, Middleton G, et al. Panitumumab and irinotecan versus irinotecan alone for patients with KRAS wild-type, fluorouracil-resistant advanced colorectal cancer (PICCOLO): a prospectively stratified randomised trial. *Lancet Oncol* 2013;14:749-759. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23725851>.

611. Pietrantonio F, Petrelli F, Coinu A, et al. Predictive role of BRAF mutations in patients with advanced colorectal cancer receiving cetuximab and panitumumab: a meta-analysis. *Eur J Cancer* 2015;51:587-594. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25673558>.

612. Rowland A, Dias MM, Wiese MD, et al. Meta-analysis of BRAF mutation as a predictive biomarker of benefit from anti-EGFR monoclonal antibody therapy for RAS wild-type metastatic colorectal cancer. *Br J Cancer* 2015;112:1888-1894. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25989278>.

613. Chen D, Huang JF, Liu K, et al. BRAFV600E mutation and its association with clinicopathological features of colorectal cancer: a systematic review and meta-analysis. *PLoS One* 2014;9:e90607. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24594804>.

614. Price TJ, Hardingham JE, Lee CK, et al. Impact of KRAS and BRAF gene mutation status on outcomes from the phase III AGITG MAX trial of capecitabine alone or in combination with bevacizumab and mitomycin in advanced colorectal cancer. *J Clin Oncol* 2011;29:2675-2682. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21646616>.

615. Saridaki Z, Papadatos-Pastos D, Tzardi M, et al. BRAF mutations, microsatellite instability status and cyclin D1 expression predict metastatic colorectal patients' outcome. *Br J Cancer* 2010;102:1762-1768. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20485284>.

616. Samowitz WS, Sweeney C, Herrick J, et al. Poor survival associated with the BRAF V600E mutation in microsatellite-stable colon cancers. *Cancer Res* 2005;65:6063-6069. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16024606>.



617. Clancy C, Burke JP, Kalady MF, Coffey JC. BRAF mutation is associated with distinct clinicopathological characteristics in colorectal cancer: a systematic review and meta-analysis. *Colorectal Dis* 2013;15:e711-718. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24112392>.

618. Santini D, Spoto C, Loupakis F, et al. High concordance of BRAF status between primary colorectal tumours and related metastatic sites: implications for clinical practice. *Ann Oncol* 2010;21:1565. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20573852>.

619. Raghav KPS, Overman MJ, Yu R, et al. HER2 amplification as a negative predictive biomarker for anti-epidermal growth factor receptor antibody therapy in metastatic colorectal cancer [abstract]. *ASCO Meeting Abstracts* 2016;34:3517. Available at: <http://meetinglibrary.asco.org/content/168395-176>.

620. Sartore-Bianchi A, Trusolino L, Martino C, et al. Dual-targeted therapy with trastuzumab and lapatinib in treatment-refractory, KRAS codon 12/13 wild-type, HER2-positive metastatic colorectal cancer (HERACLES): a proof-of-concept, multicentre, open-label, phase 2 trial. *Lancet Oncol* 2016;17:738-746. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27108243>.

621. Valtorta E, Martino C, Sartore-Bianchi A, et al. Assessment of a HER2 scoring system for colorectal cancer: results from a validation study. *Mod Pathol* 2015;28:1481-1491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26449765>.

622. Hurwitz H, Hainsworth JD, Swanton C, et al. Targeted therapy for gastrointestinal (GI) tumors based on molecular profiles: Early results from MyPathway, an open-label phase IIa basket study in patients with advanced solid tumors [abstract]. *ASCO Meeting Abstracts* 2016;34:653. Available at: <http://meetinglibrary.asco.org/content/159504-173>.

623. Wu SW, Ma CC, Li WH. Does overexpression of HER-2 correlate with clinicopathological characteristics and prognosis in colorectal

cancer? Evidence from a meta-analysis. *Diagn Pathol* 2015;10:144. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26276145>.

624. Martin V, Landi L, Molinari F, et al. HER2 gene copy number status may influence clinical efficacy to anti-EGFR monoclonal antibodies in metastatic colorectal cancer patients. *Br J Cancer* 2013;108:668-675. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23348520>.

625. Koopman M, Kortman GAM, Mekenkamp L, et al. Deficient mismatch repair system in patients with sporadic advanced colorectal cancer. *Br J Cancer* 2009;100:266-273. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19165197>.

626. Lochhead P, Kuchiba A, Imamura Y, et al. Microsatellite instability and BRAF mutation testing in colorectal cancer prognostication. *J Natl Cancer Inst* 2013;105:1151-1156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23878352>.

627. Venderbosch S, Nagtegaal ID, Maughan TS, et al. Mismatch repair status and BRAF mutation status in metastatic colorectal cancer patients: a pooled analysis of the CAIRO, CAIRO2, COIN, and FOCUS studies. *Clin Cancer Res* 2014;20:5322-5330. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25139339>.

628. Topalian SL, Hodi FS, Brahmer JR, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *N Engl J Med* 2012;366:2443-2454. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22658127>.

629. Package Insert. KEYTRUDA® (pembrolizumab). Whitehouse Station, NJ: Merck & Co, Inc.; 2016. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2016/125514s009lbl.pdf. Accessed August 15, 2016.

630. Garon EB, Rizvi NA, Hui R, et al. Pembrolizumab for the treatment of non-small-cell lung cancer. *N Engl J Med* 2015;372:2018-2028. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25891174>.



631. Le DT, Uram JN, Wang H, et al. PD-1 blockade in tumors with mismatch-repair deficiency. *N Engl J Med* 2015;372:2509-2520. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26028255>.

632. Package Insert. OPDIVO (nivolumab) injection. Princeton, NJ: Bristol-Myers Squibb Company; 2015. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/125527s000lbl.pdf. Accessed August 15, 2016.

633. Overman MJ, Kopetz S, McDermott RS, et al. Nivolumab {+/-} ipilimumab in treatment (tx) of patients (pts) with metastatic colorectal cancer (mCRC) with and without high microsatellite instability (MSI-H): CheckMate-142 interim results [abstract]. *ASCO Meeting Abstracts* 2016;34:3501. Available at: <http://meetinglibrary.asco.org/content/166455-176>.

634. Sul J, Blumenthal GM, Jiang X, et al. FDA approval summary: pembrolizumab for the treatment of patients with metastatic non-small cell lung cancer whose tumors express programmed death-ligand 1. *Oncologist* 2016;21:643-650. Available at:

635. Lewis C. Programmed death-1 inhibition in cancer with a focus on non-small cell lung cancer: rationale, nursing implications, and patient management strategies. *Clin J Oncol Nurs* 2016;20:319-326. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27206299>.

636. Hofmann L, Forschner A, Loquai C, et al. Cutaneous, gastrointestinal, hepatic, endocrine, and renal side-effects of anti-PD-1 therapy. *Eur J Cancer* 2016;60:190-209. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27085692>.

637. Zimmer L, Goldinger SM, Hofmann L, et al. Neurological, respiratory, musculoskeletal, cardiac and ocular side-effects of anti-PD-1 therapy. *Eur J Cancer* 2016;60:210-225. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27084345>.

638. Naidoo J, Wang X, Woo KM, et al. Pneumonitis in patients treated with anti-programmed death-1/programmed death ligand 1 therapy. *J*

Clin Oncol 2016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27646942>.

639. Nishino M, Chambers ES, Chong CR, et al. Anti-PD-1 inhibitor-related pneumonitis in non-small cell lung cancer. *Cancer Immunol Res* 2016;4:289-293. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26865455>.

640. Nishino M, Sholl LM, Hodi FS, et al. Anti-PD-1-related pneumonitis during cancer immunotherapy. *N Engl J Med* 2015;373:288-290. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26176400>.

641. Adam R, Miller R, Pitombo M, et al. Two-stage hepatectomy approach for initially unresectable colorectal hepatic metastases. *Surg Oncol Clin N Am* 2007;16:525-536. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17606192>.

642. Boostrom SY, Vassiliki LT, Nagorney DM, et al. Synchronous rectal and hepatic resection of rectal metastatic disease. *J Gastrointest Surg* 2011;15:1583-1588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21748454>.

643. Chen J, Li Q, Wang C, et al. Simultaneous vs. staged resection for synchronous colorectal liver metastases: a metaanalysis. *Int J Colorectal Dis* 2011;26:191-199. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20669024>.

644. Lykoudis PM, O'Reilly D, Nastos K, Fusai G. Systematic review of surgical management of synchronous colorectal liver metastases. *Br J Surg* 2014;101:605-612. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24652674>.

645. Mayo SC, Pulitano C, Marques H, et al. Surgical management of patients with synchronous colorectal liver metastasis: a multicenter international analysis. *J Am Coll Surg* 2013;216:707-716; discussion 716-708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23433970>.



646. Slesser AA, Simillis C, Goldin R, et al. A meta-analysis comparing simultaneous versus delayed resections in patients with synchronous colorectal liver metastases. *Surg Oncol* 2013;22:36-47. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23253399>.

647. Worni M, Mantyh CR, Akushevich I, et al. Is there a role for simultaneous hepatic and colorectal resections? A contemporary view from NSQIP. *J Gastrointest Surg* 2012;16:2074-2085. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22972010>.

648. Reddy SK, Pawlik TM, Zorzi D, et al. Simultaneous resections of colorectal cancer and synchronous liver metastases: a multi-institutional analysis. *Ann Surg Oncol* 2007;14:3481-3491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17805933>.

649. De Rosa A, Gomez D, Brooks A, Cameron IC. "Liver-first" approach for synchronous colorectal liver metastases: is this a justifiable approach? *J Hepatobiliary Pancreat Sci* 2013;20:263-270. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23325126>.

650. Jegatheeswaran S, Mason JM, Hancock HC, Siriwardena AK. The liver-first approach to the management of colorectal cancer with synchronous hepatic metastases: a systematic review. *JAMA Surg* 2013;148:385-391. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23715907>.

651. Lam VW, Laurence JM, Pang T, et al. A systematic review of a liver-first approach in patients with colorectal cancer and synchronous colorectal liver metastases. *HPB (Oxford)* 2014;16:101-108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23509899>.

652. Yoon HI, Koom WS, Kim TH, et al. Upfront systemic chemotherapy and short-course radiotherapy with delayed surgery for locally advanced rectal cancer with distant metastases: outcomes, compliance, and favorable prognostic factors. *PLoS One* 2016;11:e0161475. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27536871>.

653. Bartlett DL, Berlin J, Lauwers GY, et al. Chemotherapy and regional therapy of hepatic colorectal metastases: expert consensus statement. *Ann Surg Oncol* 2006;13:1284-1292. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16955384>.

654. Faron M, Pignon JP, Malka D, et al. Is primary tumour resection associated with survival improvement in patients with colorectal cancer and unresectable synchronous metastases? A pooled analysis of individual data from four randomised trials. *Eur J Cancer* 2015;51:166-176. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25465185>.

655. Karoui M, Roudot-Thoraval F, Mesli F, et al. Primary colectomy in patients with stage IV colon cancer and unresectable distant metastases improves overall survival: results of a multicentric study. *Dis Colon Rectum* 2011;54:930-938. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21730780>.

656. Venderbosch S, de Wilt JH, Teerenstra S, et al. Prognostic value of resection of primary tumor in patients with stage IV colorectal cancer: retrospective analysis of two randomized studies and a review of the literature. *Ann Surg Oncol* 2011;18:3252-3260. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21822557>.

657. McCahill LE, Yothers G, Sharif S, et al. Primary mFOLFOX6 plus bevacizumab without resection of the primary tumor for patients presenting with surgically unresectable metastatic colon cancer and an intact asymptomatic colon cancer: definitive analysis of NSABP trial C-10. *J Clin Oncol* 2012;30:3223-3228. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22869888>.

658. Cirocchi R, Trastulli S, Abraha I, et al. Non-resection versus resection for an asymptomatic primary tumour in patients with unresectable Stage IV colorectal cancer. *Cochrane Database Syst Rev* 2012;8:CD008997. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22895981>.

659. Anwar S, Peter MB, Dent J, Scott NA. Palliative excisional surgery for primary colorectal cancer in patients with incurable metastatic



disease. Is there a survival benefit? A systematic review. *Colorectal Dis* 2012;14:920-930. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21899714>.

660. Yang TX, Billah B, Morris DL, Chua TC. Palliative resection of the primary tumour in patients with Stage IV colorectal cancer: systematic review and meta-analysis of the early outcome after laparoscopic and open colectomy. *Colorectal Dis* 2013;15:e407-419. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23895669>.

661. Joyce DL, Wahl RL, Patel PV, et al. Preoperative positron emission tomography to evaluate potentially resectable hepatic colorectal metastases. *Arch Surg* 2006;141:1220-1226; discussion 1227. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17178965>.

662. Pelosi E, Deandreis D. The role of 18F-fluoro-deoxy-glucose positron emission tomography (FDG-PET) in the management of patients with colorectal cancer. *Eur J Surg Oncol* 2007;33:1-6. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17126522>.

663. Moulton CA, Gu CS, Law CH, et al. Effect of PET before liver resection on surgical management for colorectal adenocarcinoma metastases: a randomized clinical trial. *JAMA* 2014;311:1863-1869. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24825641>.

664. Maffione AM, Lopci E, Bluemel C, et al. Diagnostic accuracy and impact on management of (18)F-FDG PET and PET/CT in colorectal liver metastasis: a meta-analysis and systematic review. *Eur J Nucl Med Mol Imaging* 2015;42:152-163. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25319712>.

665. Gill S, Berry S, Biagi J, et al. Progression-free survival as a primary endpoint in clinical trials of metastatic colorectal cancer. *Curr Oncol* 2011;18 Suppl 2:S5-S10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21969810>.

666. Booth CM, Eisenhauer EA. Progression-free survival: meaningful or simply measurable? *J Clin Oncol* 2012;30:1030-1033. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22370321>.

667. Chibaudel B, Bonnetain F, Shi Q, et al. Alternative end points to evaluate a therapeutic strategy in advanced colorectal cancer: evaluation of progression-free survival, duration of disease control, and time to failure of strategy--an Aide et Recherche en Cancerologie Digestive Group Study. *J Clin Oncol* 2011;29:4199-4204. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21969501>.

668. Shi Q, de Gramont A, Grothey A, et al. Individual patient data analysis of progression-free survival versus overall survival as a first-line end point for metastatic colorectal cancer in modern randomized trials: findings from the analysis and research in cancers of the digestive system database. *J Clin Oncol* 2015;33:22-28. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25385741>.

669. Carrera G, Garcia-Albeniz X, Ayuso JR, et al. Design and endpoints of clinical and translational trials in advanced colorectal cancer. a proposal from GROUP Espanol Multidisciplinar en Cancer Digestivo (GEMCAD). *Rev Recent Clin Trials* 2011;6:158-170. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21241233>.

670. Claret L, Gupta M, Han K, et al. Evaluation of tumor-size response metrics to predict overall survival in Western and Chinese patients with first-line metastatic colorectal cancer. *J Clin Oncol* 2013;31:2110-2114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23650411>.

671. Sharma MR, Gray E, Goldberg RM, et al. Resampling the N9741 trial to compare tumor dynamic versus conventional end points in randomized phase II trials. *J Clin Oncol* 2015;33:36-41. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25349295>.

672. Sargent D, Sobrero A, Grothey A, et al. Evidence for cure by adjuvant therapy in colon cancer: observations based on individual patient data from 20,898 patients on 18 randomized trials. *J Clin Oncol*



2009;27:872-877. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19124803>.

673. Seo SI, Lim SB, Yoon YS, et al. Comparison of recurrence patterns between ≤ 5 years and > 5 years after curative operations in colorectal cancer patients. *J Surg Oncol* 2013;108:9-13. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23754582>.

674. Pietra N, Sarli L, Costi R, et al. Role of follow-up in management of local recurrences of colorectal cancer: a prospective, randomized study. *Dis Colon Rectum* 1998;41:1127-1133. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9749496>.

675. Rodriguez-Moranta F, Salo J, Arcusa A, et al. Postoperative surveillance in patients with colorectal cancer who have undergone curative resection: a prospective, multicenter, randomized, controlled trial. *J Clin Oncol* 2006;24:386-393. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16365182>.

676. Secco GB, Fardelli R, Gianquinto D, et al. Efficacy and cost of risk-adapted follow-up in patients after colorectal cancer surgery: a prospective, randomized and controlled trial. *Eur J Surg Oncol* 2002;28:418-423. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12099653>.

677. Desch CE, Benson AB, Somerfield MR, et al. Colorectal cancer surveillance: 2005 update of an American Society of Clinical Oncology Practice Guideline. *J Clin Oncol* 2005;23:8512-8519. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16260687>.

678. Jeffery M, Hickey BE, Hider PN. Follow-up strategies for patients treated for non-metastatic colorectal cancer. *Cochrane Database Syst Rev* 2007;CD002200. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17253476>.

679. Renehan AG, Egger M, Saunders MP, O'Dwyer ST. Impact on survival of intensive follow up after curative resection for colorectal cancer: systematic review and meta-analysis of randomised trials. *BMJ*

2002;324:813-813. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11934773>.

680. Pita-Fernandez S, Alhayek-Ai M, Gonzalez-Martin C, et al. Intensive follow-up strategies improve outcomes in nonmetastatic colorectal cancer patients after curative surgery: a systematic review and meta-analysis. *Ann Oncol* 2015;26:644-656. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25411419>.

681. Guyot F, Faivre J, Manfredi S, et al. Time trends in the treatment and survival of recurrences from colorectal cancer. *Ann Oncol* 2005;16:756-761. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15790673>.

682. Primrose JN, Perera R, Gray A, et al. Effect of 3 to 5 years of scheduled cea and ct follow-up to detect recurrence of colorectal cancer: The facts randomized clinical trial. *JAMA* 2014;311:263-270. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24430319>.

683. Verberne CJ, Zhan Z, van den Heuvel E, et al. Intensified follow-up in colorectal cancer patients using frequent Carcino-Embryonic Antigen (CEA) measurements and CEA-triggered imaging: results of the randomized "CEAwatch" trial. *Eur J Surg Oncol* 2015;41:1188-1196. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26184850>.

684. Rosati G, Ambrosini G, Barni S, et al. A randomized trial of intensive versus minimal surveillance of patients with resected Dukes B2-C colorectal carcinoma. *Ann Oncol* 2016;27:274-280. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26578734>.

685. Lepage C, Phelip JM, Cany L, et al. Effect of 5 years of imaging and CEA follow-up to detect recurrence of colorectal cancer: The FFCD PRODIGE 13 randomised phase III trial. *Dig Liver Dis* 2015;47:529-531. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25933809>.

686. Locker GY, Hamilton S, Harris J, et al. ASCO 2006 update of recommendations for the use of tumor markers in gastrointestinal



cancer. J Clin Oncol 2006;24:5313-5327. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/17060676>.

687. Macdonald JS. Carcinoembryonic antigen screening: pros and cons. Semin Oncol 1999;26:556-560. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10528904>.

688. Kahi CJ, Boland CR, Dominitz JA, et al. Colonoscopy surveillance after colorectal cancer resection: recommendations of the US Multi-Society Task Force on Colorectal Cancer. Gastroenterology 2016;150:758-768 e711. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26892199>.

689. Green RJ, Metlay JP, Propert K, et al. Surveillance for second primary colorectal cancer after adjuvant chemotherapy: an analysis of Intergroup 0089. Ann Intern Med 2002;136:261-269. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/11848723>.

690. Martin LA, Gross ME, Mone MC, et al. Routine endoscopic surveillance for local recurrence of rectal cancer is futile. Am J Surg 2015;210:996-1002. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26453291>.

691. Pfister DG, Benson AB, 3rd, Somerfield MR. Clinical practice. Surveillance strategies after curative treatment of colorectal cancer. N Engl J Med 2004;350:2375-2382. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/15175439>.

692. Hyder O, Dodson RM, Mayo SC, et al. Post-treatment surveillance of patients with colorectal cancer with surgically treated liver metastases. Surgery 2013;154:256-265. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23889953>.

693. Patel K, Hadar N, Lee J, et al. The lack of evidence for PET or PET/CT surveillance of patients with treated lymphoma, colorectal cancer, and head and neck cancer: a systematic review. J Nucl Med 2013;54:1518-1527. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23776200>.

694. Meyerhardt JA, Mangu PB, Flynn PJ, et al. Follow-up care, surveillance protocol, and secondary prevention measures for survivors of colorectal cancer: american society of clinical oncology clinical practice guideline endorsement. J Clin Oncol 2013;31:4465-4470. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24220554>.

695. Follow-up care, surveillance protocol, and secondary prevention measures for survivors of colorectal cancer. Cancer Care Ontario; 2016. Available at:
<https://www.cancercare.on.ca/common/pages/UserFile.aspx?fileId=124839>. Accessed August 15, 2016.

696. Sargent DJ, Wieand HS, Haller DG, et al. Disease-free survival versus overall survival as a primary end point for adjuvant colon cancer studies: individual patient data from 20,898 patients on 18 randomized trials. J Clin Oncol 2005;23:8664-8670. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/16260700>.

697. Steele SR, Chang GJ, Hendren S, et al. Practice guideline for the surveillance of patients after curative treatment of colon and rectal cancer. Dis Colon Rectum 2015;58:713-725. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26163950>.

698. Butte JM, Gonen M, Allen PJ, et al. Recurrence after partial hepatectomy for metastatic colorectal cancer: potentially curative role of salvage repeat resection. Ann Surg Oncol 2015;22:2761-2771. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25572686>.

699. Litvka A, Cercek A, Segal N, et al. False-positive elevations of carcinoembryonic antigen in patients with a history of resected colorectal cancer. J Natl Compr Canc Netw 2014;12:907-913. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24925201>.

700. Nicholson BD, Shinkins B, Mant D. Blood measurement of carcinoembryonic antigen level for detecting recurrence of colorectal cancer. JAMA 2016;316:1310-1311. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27673308>.



701. Nicholson BD, Shinkins B, Pathiraja I, et al. Blood CEA levels for detecting recurrent colorectal cancer. *Cochrane Database Syst Rev* 2015;CD011134. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26661580>.

702. Lu YY, Chen JH, Chien CR, et al. Use of FDG-PET or PET/CT to detect recurrent colorectal cancer in patients with elevated CEA: a systematic review and meta-analysis. *Int J Colorectal Dis* 2013;28:1039-1047. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23407908>.

703. Martin EW, Minton JP, Carey LC. CEA-directed second-look surgery in the asymptomatic patient after primary resection of colorectal carcinoma. *Ann Surg* 1985;202:310-317. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/4037904>.

704. Yu TK, Bhosale PR, Crane CH, et al. Patterns of locoregional recurrence after surgery and radiotherapy or chemoradiation for rectal cancer. *Int J Radiat Oncol Biol Phys* 2008;71:1175-1180. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18207667>.

705. Hoffman JP, Riley L, Carp NZ, Litwin S. Isolated locally recurrent rectal cancer: a review of incidence, presentation, and management. *Semin Oncol* 1993;20:506-519. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8211198>.

706. Lowy AM, Rich TA, Skibber JM, et al. Preoperative infusional chemoradiation, selective intraoperative radiation, and resection for locally advanced pelvic recurrence of colorectal adenocarcinoma. *Ann Surg* 1996;223:177-185. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8597512>.

707. Dresen RC, Gosens MJ, Martijn H, et al. Radical resection after IORT-containing multimodality treatment is the most important determinant for outcome in patients treated for locally recurrent rectal cancer. *Ann Surg Oncol* 2008;15:1937-1947. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18389321>.

708. Kuehne J, Kleisli T, Biernacki P, et al. Use of high-dose-rate brachytherapy in the management of locally recurrent rectal cancer. *Dis Colon Rectum* 2003;46:895-899. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12847362>.

709. Wang JJ, Yuan HS, Li JN, et al. CT-guided radioactive seed implantation for recurrent rectal carcinoma after multiple therapy. *Med Oncol* 2010;27:421-429. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19415534>.

710. Das P, Delclos ME, Skibber JM, et al. Hyperfractionated accelerated radiotherapy for rectal cancer in patients with prior pelvic irradiation. *Int J Radiat Oncol Biol Phys* 2010;77:60-65. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19695792>.

711. Guren MG, Undseth C, Rekstad BL, et al. Reirradiation of locally recurrent rectal cancer: a systematic review. *Radiother Oncol* 2014;113:151-157. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25613395>.

712. Valentini V, Morganti AG, Gambacorta MA, et al. Preoperative hyperfractionated chemoradiation for locally recurrent rectal cancer in patients previously irradiated to the pelvis: A multicentric phase II study. *Int J Radiat Oncol Biol Phys* 2006;64:1129-1139. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16414206>.

713. Hewitt M, Greenfield S, Stovall E. *From Cancer Patient to Cancer Survivor: Lost in Transition*. 2006. Available at:

<http://www.nap.edu/openbook.php?isbn=0309095956>.

714. El-Shami K, Oeffinger KC, Erb NL, et al. American Cancer Society Colorectal Cancer Survivorship Care Guidelines. *CA Cancer J Clin* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26348643>.

715. Desnoo L, Faithfull S. A qualitative study of anterior resection syndrome: the experiences of cancer survivors who have undergone resection surgery. *Eur J Cancer Care (Engl)* 2006;15:244-251. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16882120>.



716. Downing A, Morris EJ, Richards M, et al. Health-related quality of life after colorectal cancer in England: a patient-reported outcomes study of individuals 12 to 36 months after diagnosis. *J Clin Oncol* 2015;33:616-624. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25559806>.

717. Gami B, Harrington K, Blake P, et al. How patients manage gastrointestinal symptoms after pelvic radiotherapy. *Aliment Pharmacol Ther* 2003;18:987-994. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14616164>.

718. McGough C, Baldwin C, Frost G, Andreyev HJ. Role of nutritional intervention in patients treated with radiotherapy for pelvic malignancy. *Br J Cancer* 2004;90:2278-2287. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15162154>.

719. Schneider EC, Malin JL, Kahn KL, et al. Surviving colorectal cancer : patient-reported symptoms 4 years after diagnosis. *Cancer* 2007;110:2075-2082. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17849466>.

720. Sprangers MA, Taal BG, Aaronson NK, te Velde A. Quality of life in colorectal cancer. Stoma vs. nonstoma patients. *Dis Colon Rectum* 1995;38:361-369. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7720441>.

721. Baxter NN, Habermann EB, Tepper JE, et al. Risk of pelvic fractures in older women following pelvic irradiation. *JAMA* 2005;294:2587-2593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16304072>.

722. Lange MM, Maas CP, Marijnen CA, et al. Urinary dysfunction after rectal cancer treatment is mainly caused by surgery. *Br J Surg* 2008;95:1020-1028. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18563786>.

723. Lange MM, Marijnen CA, Maas CP, et al. Risk factors for sexual dysfunction after rectal cancer treatment. *Eur J Cancer* 2009;45:1578-1588. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19147343>.

724. Jansen L, Herrmann A, Stegmaier C, et al. Health-related quality of life during the 10 years after diagnosis of colorectal cancer: a population-based study. *J Clin Oncol* 2011;29:3263-3269. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21768465>.

725. Lynch BM, Steginga SK, Hawkes AL, et al. Describing and predicting psychological distress after colorectal cancer. *Cancer* 2008;112:1363-1370. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18318044>.

726. Mols F, Beijers T, Lemmens V, et al. Chemotherapy-induced neuropathy and its association with quality of life among 2- to 11-year colorectal cancer survivors: results from the population-based PROFILES registry. *J Clin Oncol* 2013;31:2699-2707. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23775951>.

727. Thong MS, Mols F, Wang XS, et al. Quantifying fatigue in (long-term) colorectal cancer survivors: a study from the population-based patient reported outcomes following initial treatment and long term evaluation of survivorship registry. *Eur J Cancer* 2013;49:1957-1966. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23453750>.

728. Vardy JL, Dhillon HM, Pond GR, et al. Cognitive function in patients with colorectal cancer who do and do not receive chemotherapy: a prospective, longitudinal, controlled study. *J Clin Oncol* 2015;33:4085-4092. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26527785>.

729. Wright P, Downing A, Morris EJ, et al. Identifying social distress: a cross-sectional survey of social outcomes 12 to 36 months after colorectal cancer diagnosis. *J Clin Oncol* 2015. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26282636>.

730. Denlinger CS, Barsevick AM. The challenges of colorectal cancer survivorship. *J Natl Compr Canc Netw* 2009;7:883-893. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19755048>.

731. Faul LA, Shibata D, Townsend I, Jacobsen PB. Improving survivorship care for patients with colorectal cancer. *Cancer Control* 2010;17:35-43. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20010517>.

732. Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol* 2006;24:3535-3541. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16822843>.

733. Meyerhardt JA, Giovannucci EL, Ogino S, et al. Physical activity and male colorectal cancer survival. *Arch Intern Med* 2009;169:2102-2108. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20008694>.

734. Campbell PT, Patel AV, Newton CC, et al. Associations of recreational physical activity and leisure time spent sitting with colorectal cancer survival. *J Clin Oncol* 2013;31:876-885. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23341510>.

735. Kuiper JG, Phipps AI, Neuhauser ML, et al. Recreational physical activity, body mass index, and survival in women with colorectal cancer. *Cancer Causes Control* 2012;23:1939-1948. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23053793>.

736. Arem H, Pfeiffer RM, Engels EA, et al. Pre- and postdiagnosis physical activity, television viewing, and mortality among patients with colorectal cancer in the National Institutes of Health-AARP Diet and Health Study. *J Clin Oncol* 2015;33:180-188. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25488967>.

737. Je Y, Jeon JY, Giovannucci EL, Meyerhardt JA. Association between physical activity and mortality in colorectal cancer: A meta-

analysis of prospective cohort studies. *Int J Cancer* 2013;133:1905-1913. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23580314>.

738. Schmid D, Leitzmann MF. Association between physical activity and mortality among breast cancer and colorectal cancer survivors: a systematic review and meta-analysis. *Ann Oncol* 2014;25:1293-1311. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24644304>.

739. Wu W, Guo F, Ye J, et al. Pre- and post-diagnosis physical activity is associated with survival benefits of colorectal cancer patients: a systematic review and meta-analysis. *Oncotarget* 2016. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27437765>.

740. Dignam JJ, Polite BN, Yothers G, et al. Body mass index and outcomes in patients who receive adjuvant chemotherapy for colon cancer. *J Natl Cancer Inst* 2006;98:1647-1654. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17105987>.

741. Sinicrope FA, Foster NR, Yoon HH, et al. Association of obesity with DNA mismatch repair status and clinical outcome in patients with stage II or III colon carcinoma participating in NCCTG and NSABP adjuvant chemotherapy trials. *J Clin Oncol* 2012;30:406-412. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22203756>.

742. Sinicrope FA, Foster NR, Yothers G, et al. Body mass index at diagnosis and survival among colon cancer patients enrolled in clinical trials of adjuvant chemotherapy. *Cancer* 2013;119:1528-1536. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23310947>.

743. Campbell PT, Newton CC, Dehal AN, et al. Impact of body mass index on survival after colorectal cancer diagnosis: the Cancer Prevention Study-II Nutrition Cohort. *J Clin Oncol* 2012;30:42-52. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22124093>.

744. Lee J, Meyerhardt JA, Giovannucci E, Jeon JY. Association between body mass index and prognosis of colorectal cancer: a meta-analysis of prospective cohort studies. *PLoS One* 2015;10:e0120706. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25811460>.

745. Cespedes Feliciano EM, Kroenke CH, Meyerhardt JA, et al. Metabolic dysfunction, obesity, and survival among patients with early-stage colorectal cancer. *J Clin Oncol* 2016. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27601537>.

746. Daniel CR, Shu X, Ye Y, et al. Severe obesity prior to diagnosis limits survival in colorectal cancer patients evaluated at a large cancer centre. *Br J Cancer* 2016;114:103-109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26679375>.

747. Doleman B, Mills KT, Lim S, et al. Body mass index and colorectal cancer prognosis: a systematic review and meta-analysis. *Tech Coloproctol* 2016;20:517-535. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27343117>.

748. Laake I, Larsen IK, Selmer R, et al. Pre-diagnostic body mass index and weight change in relation to colorectal cancer survival among incident cases from a population-based cohort study. *BMC Cancer* 2016;16:402. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27387027>.

749. Renfro LA, Loupakis F, Adams RA, et al. Body mass index is prognostic in metastatic colorectal cancer: pooled analysis of patients from first-line clinical trials in the ARCAD database. *J Clin Oncol* 2016;34:144-150. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26503203>.

750. Kroenke CH, Neugebauer R, Meyerhardt J, et al. Analysis of body mass index and mortality in patients with colorectal cancer using causal diagrams. *JAMA Oncol* 2016;2:1137-1145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27196302>.

751. Renehan AG, Sperrin M. The obesity paradox and mortality after colorectal cancer: a causal conundrum. *JAMA Oncol* 2016;2:1127-1129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27195485>.

752. Meyerhardt JA, Niedzwiecki D, Hollis D, et al. Association of dietary patterns with cancer recurrence and survival in patients with

stage III colon cancer. *JAMA* 2007;298:754-764. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17699009>.

753. Meyerhardt JA, Sato K, Niedzwiecki D, et al. Dietary glycemic load and cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Natl Cancer Inst* 2012;104:1702-1711. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23136358>.

754. Fuchs MA, Sato K, Niedzwiecki D, et al. Sugar-sweetened beverage intake and cancer recurrence and survival in CALGB 89803 (Alliance). *PLoS One* 2014;9:e99816. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24937507>.

755. Kushi LH, Byers T, Doyle C, et al. American Cancer Society Guidelines on Nutrition and Physical Activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin* 2006;56:254-281; quiz 313-254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17005596>.

756. Hawkes AL, Chambers SK, Pakenham KI, et al. Effects of a telephone-delivered multiple health behavior change intervention (CanChange) on health and behavioral outcomes in survivors of colorectal cancer: a randomized controlled trial. *J Clin Oncol* 2013;31:2313-2321. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23690410>.

757. Rock CL, Doyle C, Demark-Wahnefried W, et al. Nutrition and physical activity guidelines for cancer survivors. *CA Cancer J Clin* 2012;62:242-274. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22539238>.

758. Sun V, Grant M, Wendel CS, et al. Dietary and behavioral adjustments to manage bowel dysfunction after surgery in long-term colorectal cancer survivors. *Ann Surg Oncol* 2015;22:4317-4324. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26159443>.

759. Cai H, Zhang G, Wang Z, et al. Relationship between the use of statins and patient survival in colorectal cancer: a systematic review and



meta-analysis. PLoS One 2015;10:e0126944. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/26030771>.

760. Cardwell CR, Hicks BM, Hughes C, Murray LJ. Statin use after colorectal cancer diagnosis and survival: a population-based cohort study. J Clin Oncol 2014;32:3177-3183. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25092779>.

761. Bains SJ, Mahic M, Myklebust TA, et al. Aspirin as secondary prevention in patients with colorectal cancer: an unselected population-based study. J Clin Oncol 2016;34:2501-2508. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/27247217>.

762. Bastiaannet E, Sampieri K, Dekkers OM, et al. Use of aspirin postdiagnosis improves survival for colon cancer patients. Br J Cancer 2012;106:1564-1570. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/22454078>.

763. Chan AT, Ogino S, Fuchs CS. Aspirin use and survival after diagnosis of colorectal cancer. JAMA 2009;302:649-658. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19671906>.

764. Goh CH, Leong WQ, Chew MH, et al. Post-operative aspirin use and colorectal cancer-specific survival in patients with stage I-III colorectal cancer. Anticancer Res 2014;34:7407-7414. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25503181>.

765. Li P, Wu H, Zhang H, et al. Aspirin use after diagnosis but not prediagnosis improves established colorectal cancer survival: a meta-analysis. Gut 2015;64:1419-1425. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25239119>.

766. McCowan C, Munro AJ, Donnan PT, Steele RJ. Use of aspirin post-diagnosis in a cohort of patients with colorectal cancer and its association with all-cause and colorectal cancer specific mortality. Eur J Cancer 2013;49:1049-1057. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/23182687>.

767. Ng K, Meyerhardt JA, Chan AT, et al. Aspirin and COX-2 inhibitor use in patients with stage III colon cancer. J Natl Cancer Inst 2015;107:345. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25432409>.

768. Domingo E, Church DN, Sieber O, et al. Evaluation of PIK3CA mutation as a predictor of benefit from nonsteroidal anti-inflammatory drug therapy in colorectal cancer. J Clin Oncol 2013;31:4297-4305. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24062397>.

769. Elwood PC, Morgan G, Pickering JE, et al. Aspirin in the treatment of cancer: reductions in metastatic spread and in mortality: a systematic review and meta-analyses of published studies. PLoS One 2016;11:e0152402. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/27096951>.

770. Liao X, Lochhead P, Nishihara R, et al. Aspirin use, tumor PIK3CA mutation, and colorectal-cancer survival. N Engl J Med 2012;367:1596-1606. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23094721>.

771. Nan H, Hutter CM, Lin Y, et al. Association of aspirin and NSAID use with risk of colorectal cancer according to genetic variants. JAMA 2015;313:1133-1142. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25781442>.

772. Reimers MS, Bastiaannet E, Langley RE, et al. Expression of HLA class I antigen, aspirin use, and survival after a diagnosis of colon cancer. JAMA Intern Med 2014;174:732-739. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/24687028>.

773. Whitlock EP, Burda BU, Williams SB, et al. Bleeding risks with aspirin use for primary prevention in adults: a systematic review for the U.S. Preventive Services Task Force. Ann Intern Med 2016;164:826-835. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/27064261>.